



## PRIOR AUTHORIZATION POLICY

- POLICY:** Antiseizure Medications – Clobazam Products Prior Authorization Policy
- Onfi® (clobazam tablets and oral suspension – Lundbeck, generic)
  - Sympazan® (clobazam oral soluble film – Aquestive Therapeutics)

**REVIEW DATE:** 11/13/2024

### INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

## CIGNA NATIONAL FORMULARY COVERAGE:

### OVERVIEW

All forms of clobazam are indicated for the adjunctive treatment of seizures associated with **Lennox-Gastaut syndrome (LGS)** in patients  $\geq 2$  years of age.<sup>1,2</sup>

### Disease Overview

LGS, a severe epileptic and developmental encephalopathy, is associated with a high rate of morbidity and mortality.<sup>3,4</sup> LGS most often begins between 3 and 5 years of age and comprises approximately 4% to 10% of childhood epilepsies; the prevalence is 0.26 per 1,000 people.<sup>3-6</sup> Affected children experience several different types of seizures, most commonly atonic seizures (sudden loss of muscle tone and limpness, also called drop seizures) and tonic seizures (increased muscle tone and muscle stiffness).<sup>3,6</sup> Seizures associated with LGS are usually resistant to treatment.<sup>6</sup> The three main forms of treatment of LGS are antiseizure medications (ASMs), dietary therapy (typically the ketogenic diet), and device/surgery (e.g., vagus nerve stimulation, corpus callosotomy).<sup>6</sup> None of the therapies are effective in all cases of LGS and the disorder has proven particularly resistant to most therapeutic options. The choice of treatment should take into consideration the patient's age and other associated conditions.

### Other Uses with Supportive Evidence

Dravet syndrome is a rare genetic epileptic encephalopathy (dysfunction of the brain) marked with frequent and/or prolonged seizures.<sup>7,8</sup> It has been estimated that 1 out of 20,000 to 40,000 infants born in the US are affected with Dravet syndrome. The seizures generally begin in the first year of life in an otherwise healthy infant. Affected individuals can develop many seizure types: myoclonic, tonic-clonic, absence, atypical absence, atonic, focal aware or impaired awareness (previously called partial seizures), and status epilepticus.<sup>8</sup> As the seizures continue, most of the children develop some level of developmental disability and other conditions associated with the syndrome. Two or more ASMs are often needed to control the seizures; most of the seizures are refractory to medications. The goals of treatment are cessation of prolonged convulsions, reductions in overall seizure frequency, and minimization of treatment side effects.<sup>9,10</sup> Some patients respond to the ketogenic diet and/or vagus nerve stimulation.

## **Guidelines/Recommendations**

The American Academy of Neurology and the American Epilepsy Society published a guideline update for treatment-resistant epilepsy (2018) stating that clobazam is probably effective as add-on therapy for LGS and is possibly effective as add-on therapy for treatment-resistant adult focal epilepsy.<sup>13</sup> Adjunctive therapy with clobazam has been effective in the treatment of uncontrolled or refractory epilepsy in adults and children.<sup>14</sup> If first-line treatment is ineffective or not tolerated, clobazam has been used as adjunctive treatment of refractory focal seizures (partial seizure and localization-related seizure) in children, young adults, and adults; adjunctive treatment of generalized tonic-clonic seizures in children, young adults, and adults; and adjunctive treatment of children and young adults with benign epilepsy with centrotemporal spikes, Panayiotopoulos syndrome or late-onset childhood occipital epilepsy (Gastaut type).

## ***Lennox-Gastaut Syndrome***

Currently, the FDA-approved drugs for this condition are Epidiolex® (cannabidiol oral solution), felbamate, lamotrigine, rufinamide tablets and oral suspension, topiramate, clobazam, and Fintepla® (fenfluramine oral solution).<sup>11,14</sup> Despite the lack of level I or level II evidence, valproic acid remains a mainstay in treatment.<sup>5,6,12</sup> If valproic acid does not provide adequate seizure control, which is almost always the case, lamotrigine should be added as the first adjunctive therapy.<sup>4</sup> If the combination regimen of valproic acid and lamotrigine does not provide adequate control, then rufinamide should be initiated and either valproic acid or lamotrigine should be discontinued. If seizure control is still not achieved, the next adjunctive therapies to consider are topiramate, clobazam, and felbamate. There is limited evidence for the use of levetiracetam, zonisamide, and Fycompa® (perampanel tablet, oral suspension). Where possible, no more than two ASMs should be used concomitantly; use of multiple ASMs raises the risk of adverse effects and/or drug-drug interactions.

## ***Dravet Syndrome***

Valproic acid and clobazam are considered to be the first-line treatment for Dravet syndrome.<sup>7,9,10</sup> If seizure control is suboptimal, Diacomit® (stiripentol capsules), Epidiolex, Fintepla, and topiramate are treatment options. If control is still inadequate, other therapies to consider are clonazepam, levetiracetam, and zonisamide. Drugs that should be avoided in Dravet syndrome include sodium channel blockers (e.g., carbamazepine, oxcarbazepine, lamotrigine, and phenytoin), Sabril® (vigabatrin tablet, oral packet), and tiagabine.

## **POLICY STATEMENT**

Prior Authorization is recommended for prescription benefit coverage of clobazam. Because of the specialized skills required for evaluation and diagnosis of patients treated with

clobazam as well as the monitoring required for adverse events and long-term efficacy, initial approval requires clobazam to be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals are provided for the duration noted below.

- **Onfi® (clobazam tablets and oral suspension – Lundbeck, generic)**
- **Sympazan® (clobazam oral soluble film – Aquestive Therapeutics)**

**is(are) covered as medically necessary when the following criteria is(are) met for fda-approved indication(s) or other uses with supportive evidence (if applicable):**

### **FDA-Approved Indication**

- 1. Lennox-Gastaut Syndrome.** Approve for 1 year if the patient meets ONE of the following (A or B):
  - A) Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, and iii):
    - i. Patient is  $\geq 2$  years of age; AND
    - ii. Patient has tried and/or is concomitantly receiving ONE of the following (a or b):
      - a) At least two other antiseizure medications; OR  
*Note:* Examples of other antiseizure medications include valproic acid, levetiracetam, zonisamide, Fycompa (perampanel), vigabatrin, others.
      - b) One of lamotrigine, topiramate, rufinamide, felbamate, Fintepla (fenfluramine oral solution), or Epidiolex (cannabidiol oral solution); AND
    - iii. Clobazam is prescribed by or in consultation with a neurologist.
  - B) Patient is Currently Receiving Clobazam.** Approve if the patient is responding to therapy, as determined by the prescriber.  
*Note:* Examples of therapy response include reduced seizure severity, frequency, and/or duration from baseline (prior to initiation of clobazam).

### **Other Uses with Supportive Evidence**

- 2. Dravet Syndrome.** Approve for 1 year if the patient meets ONE of the following (A or B):
  - A) Initial Therapy.** Approve if the patient meets BOTH of the following (i and ii):
    - i. Patient is  $\geq 2$  years of age; AND
    - ii. Clobazam is prescribed by or in consultation with a neurologist.
  - B) Patient is Currently Receiving Clobazam.** Approve if the patient is responding to therapy, as determined by the prescriber.  
*Note:* Examples of therapy response include reduced seizure severity, frequency, and/or duration from baseline (prior to initiation of clobazam).
- 3. Treatment-Refractory Seizures/Epilepsy.** Approve for 1 year if the patient meets ONE of the following (A or B):
  - A) Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, and iii):
    - i. Patient is  $\geq 2$  years of age; AND
    - ii. Patient has tried and/or is concomitantly receiving at least two other antiseizure medications; AND  
*Note:* Examples of other antiseizure medications are valproic acid, lamotrigine, topiramate, clonazepam, levetiracetam, zonisamide, rufinamide, felbamate.
    - iii. Clobazam is prescribed by or in consultation with a neurologist.
  - B) Patient is Currently Receiving Clobazam.** Approve if the patient is responding to therapy, as determined by the prescriber.

Note: Examples of therapy response include reduced seizure severity, frequency, and/or duration from baseline (prior to initiation of clobazam).

## CONDITIONS NOT COVERED

- **Onfi® (clobazam tablets and oral suspension – Lundbeck, generic)**
- **Sympazan® (clobazam oral soluble film – Aquestive Therapeutics)**

**is(are) considered experimental, investigational, or unproven for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):**

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

## REFERENCES

1. Onfi® tablets and oral suspension [prescribing information]. Deerfield, IL: Lundbeck; March 2024.
2. Sympazan® oral film [prescribing information]. Warren, NJ: Aquestive Therapeutics; March 2024.
3. Sirven JI, Shafer PO. Epilepsy Foundation – Lennox-Gastaut Syndrome. Updated February 2020. Available at: <https://www.epilepsy.com/learn/types-epilepsy-syndromes/lennox-gastaut-syndrome-lgs/>. Accessed on October 29, 2024.
4. Cross JH, Auvin S, Falip M, et al. Expert opinion on the management of Lennox-Gastaut syndrome: treatment algorithms and practical considerations. *Front Neurol*. 2017; 8:505.
5. Ostendorf AP, Ng YT. Treatment-resistant Lennox-Gastaut syndrome: therapeutic trends, challenges, and future directions. *Neuropsych Dis Treatment*. 2017; 13:1131-1140.
6. Wheless JW. National Organization for Rare Diseases – Lennox-Gastaut syndrome. Updated May 20, 2024. Available at: <https://rarediseases.org/rare-diseases/lennox-gastaut-syndrome/>. Accessed on October 29, 2024.
7. Dravet Foundation – Dravet Syndrome. Available at: <https://www.dravetfoundation.org/what-is-dravet-syndrome/>. Accessed on October 29, 2024.
8. Shafer PO. Epilepsy Foundation – Dravet Syndrome. Updated August 2020. Available at: <https://www.epilepsy.com/learn/types-epilepsy-syndromes/dravet-syndrome/>. Accessed on October 29, 2024.
9. Wirrell EC, Hood V, Knupp KG, et al. International consensus on diagnosis and management of Dravet syndrome. *Epilepsia*. 2022;63(7):1761-1777.
10. Knupp KG, Wirrell EC. Treatment Strategies for Dravet Syndrome. *CNS Drugs*. 2018;32(4):335-350.
11. Lennox-Gastaut Syndrome Foundation – Lennox-Gastaut Syndrome. Updated October 25, 2024. Available at: <https://www.lgsfoundation.org/about-lgs-2/what-is-lennox-gastaut-syndrome/>. Accessed on October 29, 2024.
12. Cherian KA. Lennox-Gastaut syndrome treatment & management. Updated September 10, 2024. Available at: <https://emedicine.medscape.com/article/1176735-treatment/>. Accessed on October 29, 2024.
13. Kanner AM, Ashman E, Gloss D, et al. Practice guideline update summary: Efficacy and tolerability of the new antiepileptic drugs II: Treatment-resistant epilepsy. Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the

American Academy of Neurology and the American Epilepsy Society. *Neurology*. 2018; 91:82-90.

14. Merative Micromedex®. Merative 2024. Available at: <https://www.micromedexsolutions.com/>. Accessed on October 29, 2024. Search terms: clobazam, Fintepla.

## HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	The policy name was changed from Antiepileptics – Clobazam Prior Authorization Policy to Antiseizure Medications – Clobazam Prior Authorization Policy. Throughout the criteria, reference to antiepileptic medications was changed to antiseizure medications. <b>Lennox-Gastaut Syndrome:</b> The addition of Fintepla to the list of “one of the following” antiseizure medications that a patient has tried and/or is concomitantly receiving. Examples of other antiseizure medications were moved to a Note.	11/15/2023
Annual Revision	No criteria changes.	11/13/2024

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.