

PRIOR AUTHORIZATION POLICY

POLICY: Oncology – Valchlor Prior Authorization Policy

Valchlor® (mechlorethamine topical gel – Helsinn)

REVIEW DATE: 12/13/2023

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Valchlor, a nitrogen mustard, is indicated for the topical treatment of Stage IA and IB **mycosis fungoides-type cutaneous T-cell lymphoma** in patients who have received prior skin-directed therapy.¹

Guidelines

Valchlor is addressed in National Comprehensive Cancer Network guidelines:

- Histiocytic neoplasms: Guidelines (version 1.2023 August 11, 2023) recommend Valchlor for the topical treatment of unifocal Langerhans cell histiocytosis with isolated skin disease.^{2,5}
- Primary cutaneous lymphomas: Guidelines (version 1.2023 January 5, 2023) recommend Valchlor for the topical treatment of primary cutaneous B-cell lymphoma, mycosis fungoides/Sezary syndrome, and primary cutaneous CD30+ T-cell lymphoproliferative disorders.^{2,3}
- **T-cell lymphomas:** Guidelines (version 1.2023 January 5, 2023) recommend Valchlor for the topical treatment of adult T-cell leukemia/lymphoma chronic/smoldering subtype.^{2,4}

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Valchlor. All approvals are provided for the duration noted below.

• Valchlor® (mechlorethamine topical gel – Helsinn)

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indication

 Cutaneous Lymphomas. Approve for 1 year if the patient is ≥ 18 years of age. <u>Note</u>: Includes mycosis fungoides/Sezary syndrome, primary cutaneous B-cell lymphoma, primary cutaneous CD30+ T-cell lymphoproliferative disorders.

Other Uses with Supportive Evidence

- **2. Adult T-Cell Leukemia/Lymphoma**. Approve for 1 year if the patient has chronic/smoldering subtype of adult T-cell leukemia/lymphoma.
- **3. Langerhans Cell Histiocytosis**. Approve for 1 year if, according to the prescriber, patient has unifocal Langerhans cell histiocytosis with isolated skin disease.

CONDITIONS NOT COVERED

Valchlor® (mechlorethamine topical gel – Helsinn)

is(are) considered experimental, investigational, or unproven for ANY other use(s).

REFERENCES

- 1. Valchlor® topical gel [prescribing information]. Iselin, NJ: Helsinn; January 2020.
- 2. The NCCN Drugs and Biologics Compendium. © 2023 National Comprehensive Cancer Network. Available at: http://www.nccn.org. Accessed on December 8, 2023. Search term: mechlorethamine.
- 3. The NCCN Primary Cutaneous Lymphomas Clinical Practice Guidelines in Oncology (version 1.2023 January 5, 2023). © 2023 National Comprehensive Cancer Network. Available at: http://www.nccn.org. Accessed on December 8, 2023.
- The NCCN T-Cell Lymphomas Clinical Practice Guidelines in Oncology (version 1.2023 January 5, 2023). © 2023 National Comprehensive Cancer Network. Available at: http://www.nccn.org. Accessed on December 8, 2023.
- 5. The NCCN Histiocytic Neoplasms Clinical Practice Guidelines in Oncology (version 1.2023 August 11, 2023). © 2023 National Comprehensive Cancer Network. Available at: http://www.nccn.org. Accessed December 8, 2023.

HISTORY

Type of	Summary of Changes	Review
Revision		Date

Annual	No criteria change.	11/30/2022
Revision		
Annual	No criteria change.	12/13/2023
Revision		

[&]quot;Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. © 2023 Cigna