



PRIOR AUTHORIZATION POLICY

POLICY: Inflammatory Conditions – Sotyktu Prior Authorization Policy

- Sotyktu™ (deucravacitinib tablets – Bristol Myers Squibb)

REVIEW DATE: 09/13/2023; selected revision 03/27/2027

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Sotyktu, a tyrosine kinase 2 (TYK2) inhibitor, is indicated for treatment of moderate to severe **plaque psoriasis** in adults who are candidates for systemic therapy or phototherapy.¹ Limitation of use: Sotyktu is not recommended in combination with potent immunosuppressants.

Guidelines

Guidelines have not been updated to address Sotyktu. Joint guidelines from the American Academy of Dermatology and National Psoriasis Medical Board (2019) have been published for management of psoriasis with biologics.² These guidelines list all the biologics approved at the time of publication as agents that may be used as monotherapy for adults with moderate to severe psoriasis. Guidelines from the European Dermatology Forum (2015) recommend biologics (i.e., etanercept, adalimumab, infliximab, Stelara [ustekinumab subcutaneous injection]) as second-line therapy for induction and long-term treatment if phototherapy and conventional systemic agents have failed, are contraindicated, or are not tolerated.³

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Sotyktu. Because of the specialized skills required for evaluation and diagnosis of patients treated with Sotyktu as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Sotyktu to be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals are for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

- **Sotyktu™ (deucravacitinib tablets (Bristol Myers Squibb))**

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indication

1. Plaque Psoriasis. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 3 months if the patient meets ALL of the following (i, ii, and iii):

i. Patient is \geq 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) Patient has tried at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant; OR

Note: Examples of one traditional systemic agent include methotrexate, cyclosporine, acitretin tablets, or psoralen plus ultraviolet A light (PUVA). An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the patient has already had a 3-month trial or previous intolerance to at least one biologic other than the requested drug. A biosimilar of the requested biologic does not count. Refer to [Appendix](#) for examples of biologics used for plaque psoriasis. A patient who has already tried a biologic for psoriasis is not required to “step back” and try a traditional systemic agent for psoriasis.

b) Patient has a contraindication to methotrexate, as determined by the prescriber; AND

iii. The medication is prescribed by or in consultation with a dermatologist.

B) Patient is Currently Receiving Sotyktu. Approve for 1 year meets ALL of the following (i, ii, and iii):

i. Patient has been established on therapy for at least 3 months; AND

Note: A patient who has received < 3 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).

ii. Patient experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating the requested drug) in at least one of the following: estimated body surface area, erythema, induration/thickness, and/or scale of areas affected by psoriasis; AND

iii. Compared with baseline (prior to initiating the requested drug), patient experienced an improvement in at least one symptom, such as decreased pain, itching, and/or burning.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

- **Sotyktu™ (deucravacitinib tablets (Bristol Myers Squibb))**

is(are) considered experimental, investigational or unproven for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Concurrent Use with other Biologics or with Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs).** Data are lacking evaluating concomitant use of Sotyktu with another biologic or with a targeted synthetic DMARD for an inflammatory condition (see [Appendix](#) for examples). Combination therapy with biologics and/or biologics + targeted synthetic DMARDs has a potential for a higher rate of adverse effects and lack controlled trial data in support of additive efficacy.⁴

Note: This does NOT exclude the use of methotrexate (a traditional systemic agent used to treat psoriasis) in combination with Sotyktu.

- 2. Concurrent use with Other Potent Immunosuppressants, Including Methotrexate.**¹ Co-administration with other potent immunosuppressive drugs has the risk of added immunosuppression and has not been evaluated.

REFERENCES

1. Sotyktu™ tablets [prescribing information]. Princeton, NJ: Bristol Myers Squibb; September 2022.
2. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
3. Nast A, Gisondi P, Ormerod AD, et al. European S3-Guidelines on the systemic treatment of psoriasis vulgaris – Update 2015 – Short version – EDF in cooperation with EADV and IPC. *J Eur Acad Dermatol Venereol.* 2015;29(12):2277-2294.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	09/14/2022
Annual Revision	No criteria changes.	09/13/2023
Selected Revision	Plaque Psoriasis: For a patient currently taking Sotyktu, the timeframe for established on therapy was changed from 90 days to 3 months.	03/27/2024

APPENDIX

	Mechanism of Action	Examples of Inflammatory Indications*
Biologics		
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA

Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
Simponi®, Simponi® Aria™ (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
Actemra® (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA
		IV formulation: JIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA [^] , RA
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC
Siliq™ (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection)	Inhibition of IL-17A	AS, ERA, nr-axSpA, PsO, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Ilumya™ (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PSA, PsO
		IV formulation: CD
Tremfya™ (guselkumab SC injection)	Inhibition of IL-23	PsO
Entyvio™ (vedolizumab IV infusion)	Integrin receptor antagonist	CD, UC
Oral Therapies/Targeted Synthetic DMARDs		
Otezla® (apremilast tablets)	Inhibition of PDE4	PsO, PsA
Cibinqo™ (abrocitinib tablets)	Inhibition of JAK pathways	AD
Olumiant® (baricitinib tablets)	Inhibition of JAK pathways	RA
Rinvoq® (upadacitinib extended-release tablets)	Inhibition of JAK pathways	AD, AS, nr-axSpA, RA, PsA, UC
Sotyktu™ (deucravacitinib tablets)	Inhibition of TYK2	PsO
Xeljanz® (tofacitinib tablets)	Inhibition of JAK pathways	RA, PJIA, PsA, UC
Xeljanz® XR (tofacitinib extended-release tablets)	Inhibition of JAK pathways	RA, PsA, UC

* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; [^] Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; TYK2 – Tyrosine kinase 2.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health

Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.