



PRIOR AUTHORIZATION POLICY

POLICY: Growth Disorders – Ngenla Prior Authorization Policy

- Ngenla® (somatrogen-ghla subcutaneous injection – Pfizer)

REVIEW DATE: 07/26/2023; selected revision 11/01/2023

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Ngenla, a human growth hormone (hGH) product, is indicated for the treatment **growth failure due to inadequate secretion of growth hormone (GH)** in pediatric patients ≥ 3 years of age.¹

Disease Overview

Ngenla is a hGH analog which is made up of the amino acid sequence of hGH with an added three copies of the C-terminal peptide of human chorionic gonadotropin.¹ The addition of the C-terminal peptides lead to a longer half-life. Ngenla binds to the GH receptor which initiates changes in growth and metabolism. In children with GH deficiency (GHD), somatropin is effective for increasing final adult height.² Somatropin therapy is recommended to normalize adult height and prevent extreme shortness in children and adolescents with GHD.² In addition to congenital causes, hypopituitarism may also be caused by radiation therapy; somatropin may be used to improve final height of children who have undergone radiation.^{3,4}

Guidelines

Current guidelines do not specifically address Ngenla. Neither the Pediatric Endocrine Society guidelines for children and adolescents with GHD² (2016) nor the GH Research Society guidelines on children with short stature¹¹ (2019) recommend a

specific GH product for GHD. Guidelines recommend the use of GH to normalize adult height and prevent extreme shortness in pediatric patients with GHD.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Ngenla. All reviews will be directed to a clinician (i.e., pharmacist) for verification of criteria. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Ngenla as well as monitoring required for adverse events and long-term efficacy, initial approval requires the patient to be evaluated by a physician who specializes in the condition being treated. hGH is FDA-approved for treatment of a limited number of conditions. The FDA has not approved the use of hGH as therapy for anti-aging, longevity, or cosmetic or performance enhancement. Federal law prohibits the dispensing of hGH for non-approved purposes. A pharmacy's failure to comply with that law could result in significant criminal penalties to the pharmacy and its employees. Accordingly, a pharmacy may decline to dispense prescriptions for hGH when written by a physician or other authorized prescribers who they believe may be involved in or affiliated with the fields of anti-aging, longevity, rejuvenation, cosmetic, performance enhancement, or sports medicine.

- **Ngenla® (somatrogon-ghla subcutaneous injection (Pfizer) is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):**

FDA-Approved Indication

- 1. Growth Hormone Deficiency in a Pediatric Patient (≥ 3 years of age to < 18 years of age).** Approve for 1 year if the patient meets the following (A or B):
 - A) Initial Therapy with any Growth Hormone Agent.** Approve if the patient meets one of the following (i, ii, iii, iv, or v):
 - i. Patient meets BOTH of the following a (a and b):**
 - a) Patient meets at least ONE of the following (1 or 2):**
 - (1)** Patient has had two growth hormone stimulation tests performed with any of the following agents: levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon AND the peak growth hormone response to both tests are < 10 ng/mL; OR
 - (2)** Patient meets BOTH of the following (a and b):
 - (a)** Patient has had at least one growth hormone stimulation test performed with any of the following agents: levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon AND the peak growth hormone response to at least one test is < 10 ng/mL; AND
 - (b)** Patient has at least one risk factor for growth hormone deficiency (e.g., the height for age curve has deviated downward across two major height percentiles [e.g., from

above the 25th percentile to below the 10th percentile]; the child's growth rate is less than the expected normal growth rate based on age and gender; low insulin-like growth factor [IGF]-1 and/or IGFBP-3 levels; the child has a very low peak growth hormone level on provocative testing as defined by the prescribing physician; the child's growth velocity is less than the 10th percentile for age and gender [height velocity percentile is NOT the same as height-for-age percentile]; the patient is status post craniopharyngioma resection; the patient has optic nerve hypoplasia; the patient has a growth hormone gene deletion); AND

- b)** Patient has been evaluated by an endocrinologist.
- ii.** Patient has undergone brain radiation or tumor resection AND meets BOTH of the following (a and b):
 - a)** Patient meets at least ONE of the following (1 or 2):
 - (1)** Patient meets BOTH of the following (i and ii):
 - (i)** Patient has had one growth hormone stimulation test with any of the following agents: levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon; AND
 - (ii)** The peak growth hormone response to at least one test is < 10 ng/mL; OR
 - (2)** Patient has a deficiency in at least one other pituitary hormone (i.e., adrenocorticotrophic hormone, thyroid-stimulating hormone, gonadotropin [luteinizing hormone and/or follicle stimulating hormone deficiency are counted as one deficiency], or prolactin); AND
 - b)** Patient has been evaluated by an endocrinologist.
- iii.** Patient has congenital hypopituitarism AND meets BOTH of the following (a and b):
 - a)** Patient meets at least ONE of the following (1, 2, or 3):
 - (1)** Patient meets BOTH of the following (i and ii):
 - (i)** Patient has had one growth hormone stimulation test with any of the following agents: levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon; AND
 - (ii)** The peak growth hormone response to at least one test is < 10 ng/mL; OR
 - (2)** Patient has a deficiency in at least one other pituitary hormone (i.e., adrenocorticotrophic hormone, thyroid-stimulating hormone, gonadotropin [luteinizing hormone and/or follicle stimulating hormone deficiency are counted as one deficiency], or prolactin); OR
 - (3)** Patient has the imaging triad of ectopic posterior pituitary and pituitary hypoplasia with abnormal pituitary stalk; AND
 - b)** Patient has been evaluated by an endocrinologist.
- iv.** Patient has multiple pituitary hormone deficiencies and meets BOTH of the following (a and b):

Note: Growth hormone deficiency may occur in combination with other pituitary hormone deficiencies and is referred to as hypopituitarism, panhypopituitarism, or multiple pituitary hormone deficiency.

a) Patient meets at least ONE of the following (1 or 2):

(1) Patient has three or more of the following pituitary hormone deficiencies: somatotropin (growth hormone), adrenocorticotrophic hormone, thyroid-stimulating hormone, gonadotropin (luteinizing hormone and/or follicle stimulating hormone deficiency are counted as one deficiency), and prolactin; OR

(2) Patient meets BOTH of the following (i and ii):

(i) Patient has had one growth hormone stimulation test with any of the following agents: levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon; AND

(ii) The peak growth hormone response to at least one test is < 10 ng/mL; AND

b) Patient has been evaluated by an endocrinologist.

v. Patient has had a hypophysectomy (surgical removal of pituitary gland).

B) Patient is Currently Receiving Ngenla or is switching to Ngenla from another Growth Hormone Agent (Patient has been established on either therapy for ≥ 10 months). Approve if the patient meets one of the following (i or ii):

i. Patient is < 12 years of age: Patient's height has increased by ≥ 2 cm/year in the most recent year; OR

ii. Patient is ≥ 12 years of age and < 18 years of age: Patient meets both of the following (a and b):

a) Patient's height has increased by ≥ 2 cm/year in the most recent year; AND

b) Patient's epiphyses are open.

CONDITIONS NOT COVERED

• **Ngenla® (somatrogon-ghla subcutaneous injection (Pfizer)) is(are) considered experimental, investigational or unproven for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):**

1. Athletic Ability Enhancement.⁵ Somatotropin and related agents are not FDA-approved for athletic performance enhancement or for body building in non-athletes. Federal law prohibits the distribution or dispensing of somatotropin or related agents for non-FDA approved uses.

2. Central Precocious Puberty. Children with precocious puberty are often treated with gonadotropin releasing hormone (GnRH) agonists (Lupron® [leuprolide acetate injection]) to suppress pituitary gonadal activity, to slow the advancement of bone age (prevent premature fusion of the epiphyseal growth plates), and to improve adult height. In some patients, GnRH agonist therapy may result in marked deceleration of bone growth and may result in adult height that is less than the midparental height. Somatotropin has been used in girls when growth

velocity decreases or if it appears that the targeted adult height will not be attained.⁶ There are no large well-controlled trials on the efficacy and safety of adding somatropin to GnRH agonist therapy in these children or the effect on final height.^{6,7}

3. Congenital Adrenal Hyperplasia (CAH).^{8,9} The Endocrine Society clinical practice guidelines on CAH due to steroid 21-hydroxylase deficiency recommend against the use of experimental treatment approaches outside of formally approved clinical trials.⁹ Children with predicted adult height standard deviation ≤ -2.25 may be considered for growth-promoting treatments in appropriately controlled trials.

4. Constitutional Delay of Growth and Puberty. These children have delayed skeletal maturation and pubertal development. Administering somatropin does not increase adult height (which is usually normal).¹⁰ Short-term androgen therapy accelerates growth and the rate of pubertal advancement in boys.

REFERENCES

1. Ngenla® subcutaneous injection [prescribing information]. New York, NY: Pfizer; June 2023.
2. Grimberg A, DiVall SA, Polychronakos C, et al; Drug and Therapeutics Committee and Ethics Committee of the Pediatric Endocrine Society. Guidelines for growth hormone and insulin-like growth factor-I treatment in children and adolescents: growth hormone deficiency, idiopathic short stature, and primary insulin-like growth factor-I deficiency. *Horm Res Paediatr.* 2016;86(6):361-397.
3. Melmed S. Idiopathic adult growth hormone deficiency. *J Clin Endocrinol Metab.* 2013;98:2187-2197.
4. Isfan F, Kanold J, Merlin E, et al. Growth hormone treatment impact on growth rate and final height of patients who received HSCT with TBI or/and cranial irradiation in childhood: a report from the French Leukaemia Long-Term Follow-Up Study (LEA). *Bone Marrow Transplant.* 2012;47:684-693.
5. Clemmons DR, Molitch M, Hoffman AR, et al. Growth hormone should be used only for approved indications. *J Clin Endocrinol Metab.* 2014;99:409-411.
6. Carel JC, Eugster EA, Rogol A, et al on behalf of the members of the ESPE-LWPES GnRH Analogs Consensus Conference Group. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. *Pediatrics.* 2009;123:e752-762.
7. Carel JC, Leger J. Precocious puberty. *N Engl J Med.* 2008;358:2366-2377.
8. Lin-Su K, Harbison MD, Lekarev O, et al. Final adult height in children with congenital adrenal hyperplasia treated with growth hormone. *J Clin Endocrinol Metab.* 2011;96:1710-1717.
9. Speiser PW, Arlt W, Auchus RJ, et al. Congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2018;103(11):4043-4088.
10. De Luca F, Argente J, Cavallo L, et al; International Workshop on Management of Puberty for Optimum Auxological Results. Management of puberty in constitutional delay of growth and puberty. *Pediatr Endocrinol Metab.* 2001;14 Suppl 2:953-957.
11. Collett-Solberg PF, Ambler G, Backelijaw PF, et al. Diagnosis, genetics, and therapy of short stature in children: A growth hormone research society international perspective. *Horm Res Paediatr.* 2019;92(1):1-14.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	07/26/2023

Selected Revision	Growth Hormone Deficiency in a Pediatric Patient (≥ 3 years of age to < 18 years of age): Criterion that specified the result of a growth hormone stimulation test to be < 10 ng/mL was added. Criterion that the stimulation test show an inadequate response as defined by a peak response below the normal reference range as determined by the testing laboratory was removed.	11/01/2023
-------------------	---	------------

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. © 2023 Cigna