



Medical Coverage Policy

Effective Date..... 4/15/2021
Next Review Date..... 3/15/2022
Coverage Policy Number 0178

Breast Reconstruction Following Mastectomy or Lumpectomy

Table of Contents

Overview	1
Coverage Policy.....	1
General Background.....	4
Medicare Coverage Determinations	20
Coding/Billing Information.....	20
References	24

Related Coverage Resources

- [Botulinum Therapy](#)
- [Breast Implant Removal](#)
- [Complex Lymphedema Therapy \(Complete Decongestive Therapy\)](#)
- [Injectable Fillers](#)
- [Panniculectomy and Abdominoplasty](#)
- [Pneumatic Compression Devices and Compression Garments](#)
- [Reduction Mammoplasty](#)
- [Redundant Skin Surgery](#)
- [Scar Revision](#)
- [Surgical Treatment of Chest Wall Deformities](#)
- [Surgical Treatments for Lymphedema and Lipedema](#)
- [Tissue-Engineered Skin Substitutes](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses reconstructive breast surgery and external breast prostheses and mastectomy bras following mastectomy or lumpectomy.

Coverage Policy

Coverage for breast reconstruction* and breast prostheses following mastectomy or lumpectomy is governed by federal and/or state mandates.

Breast Reconstruction

***Please note:** Coverage for breast reconstruction services following mastectomy and lumpectomy is available to both females and males. In addition, a diagnosis of breast cancer is not required for breast reconstruction services to be covered, and the timing of reconstructive services is not a factor in coverage.

Breast reconstruction following mastectomy or lumpectomy is considered medically necessary for EITHER of the following:

- **breast reconstruction procedures performed on the diseased/affected breast (i.e., breast on which the mastectomy/lumpectomy was performed), including:**
 - areolar and nipple reconstruction
 - areolar and nipple tattooing
 - autologous fat transplant (i.e., liposuction, lipoinjection, lipofilling, lipomodeling)
 - breast implant removal and subsequent reimplantation
 - capsulectomy
 - capsulotomy
 - implantation of tissue expander
 - implantation of U.S. Food and Drug Administration (FDA)-approved internal breast prosthesis
 - oncoplastic reconstruction
 - reconstructive surgical revisions
 - tissue/muscle reconstruction procedures (e.g., flaps), including, but not limited to, the following:
 - deep inferior epigastric perforator (DIEP) flap
 - latissimus dorsi (LD) myocutaneous flap
 - Ruben's flap
 - superficial inferior epigastric perforator/artery (SIEP/SIEA) flap
 - superior or inferior gluteal free flap
 - thoracodorsal artery perforator (TDAP) flap
 - transverse rectus abdominus myocutaneous (TRAM) flap
 - transverse upper gracilis (TUG) flap

- **breast reconstruction procedures performed on the nondiseased/unaffected/contralateral breast, in order to produce a symmetrical appearance, including:**
 - areolar and nipple reconstruction
 - areolar and nipple tattooing
 - augmentation mammoplasty
 - augmentation with implantation of FDA-approved internal breast prosthesis when the unaffected breast is smaller than the smallest available internal prosthesis
 - autologous fat transplant (i.e., liposuction, lipoinjection, lipofilling, lipomodeling)
 - breast implant removal and subsequent reimplantation when performed to produce a symmetrical appearance
 - breast reduction by mammoplasty or mastopexy
 - capsulectomy
 - capsulotomy
 - reconstructive surgery revisions to produce a symmetrical appearance

Intraoperative assessment of tissue perfusion is considered an integral part of the breast reconstruction procedure and not separately reimbursable.

The following products* are considered medically necessary when used in association with a medically necessary breast reconstruction procedure:

- AlloDerm™
- AlloMax™
- Cortiva™
- DermACELL™
- FlexHD® Acellular Hydrated Dermis
- NeoForm™ Dermis

The following products* when used in association with a breast reconstruction procedure are considered experimental, investigational or unproven (this list may not be all-inclusive):

- ARTIA™ Reconstructive Tissue Matrix
- BellaDerm® Acellular Hydrated Dermis
- Biodesign® Nipple Reconstruction Cylinder
- DermaMatrix Acellular Dermis
- GalaFLEX® Surgical Scaffold
- GalaFORM™ 3D
- GalaSHAPE 3D
- hMatrix®
- Juvederm®
- Permacol®
- Phasix™ Mesh
- Radiesse®
- Repriza®
- SERI™ Surgical Scaffold
- Strattice™ Reconstructive Tissue Matrix
- SurgiMend®
- Veritas Collagen Matrix

***Note: Refer to the table in Appendix A for a list of products and the associated CPT and HCPCS codes.**

The following breast reconstruction procedures are considered experimental, investigational or unproven for this indication:

- the use of adipose-derived stem cells in autologous fat transplantation
- xenograft cartilage grafting

Suction lipectomy or ultrasonically-assisted suction lipectomy (liposuction) for correction of surgically-induced donor site asymmetry (e.g., trunk or extremity) that results from one or more flap breast reconstruction procedures is considered cosmetic in nature and not medically necessary.

Removal of either a saline-filled OR silicone gel-filled breast implant when associated with breast reconstruction following mastectomy or lumpectomy for ANY indication, including for the purpose of producing a symmetrical appearance of the nondiseased breast is considered medically necessary. Refer to the Breast Implant Removal Medical Coverage Policy for additional information on breast implant removal.

Following removal of a breast implant, the subsequent surgical implantation of a new U.S. Food and Drug Administration (FDA)-approved breast implant is considered medically necessary for EITHER of the following:

- breast reconstruction of a diseased or affected breast following mastectomy or lumpectomy
- creation of a symmetrical appearance in the contralateral/nondiseased breast following mastectomy or lumpectomy in the opposite breast

External Breast Prostheses and Mastectomy Bras

External breast prostheses and mastectomy bras following mastectomy or lumpectomy are covered under the core medical benefits of the plan.

An external breast prosthesis or mastectomy bra for any other indication is considered not medically necessary.

This Medical Coverage Policy does not address treatments for lymphedema. For information on these treatments, refer to the separate Cigna Medical Coverage Policies Pneumatic Compression Devices and Compression Garments, Surgical Treatments for Lymphedema and Lipedema, and Complex Lymphedema Therapy (Complete Decongestive Therapy).

General Background

Breast reconstruction was originally designed to reduce post-mastectomy complications and to establish symmetry between the surgical breast and the contralateral breast. Surgical procedures that are performed to establish symmetry can include: breast reduction; breast augmentation with an FDA-approved breast implant; and/or areola-with-nipple reconstruction and nipple-area tattooing. Breast reconstruction after mastectomy has evolved over the last century to become an integral component of therapy for patients with breast cancer. Reconstruction can occur immediately after a mastectomy, or be delayed for weeks or years until a patient undergoes radiation, chemotherapy, or determines whether they want breast reconstruction.

Prosthetic Reconstruction

Breast Implants: Breast implants can be inserted at the same time as the mastectomy (e.g., direct-to-implant breast reconstruction or one-stage immediate breast reconstruction) or in two stages, using an implanted tissue expander in the first stage followed by removal of the expander and insertion of a permanent breast implant (e.g., two-stage reconstruction or two-stage delayed reconstruction). The FDA-approved implant is placed either deep in the breast on the pectoralis fascia (submammary) or beneath the pectoralis major. The advantages of tissue expander implant reconstruction are the reliability, simplicity, and avoidance of donor-site morbidity. Complications associated with the use of breast implants can occur in the immediate perioperative period or years later. Such complications include exposure, extrusion, or infection of the implants. Longer term problems also include asymmetry, capsular contracture, malposition of the implant, rupture, and pain. These conditions, when they become clinically significant, may require removal of the implant (Roehl, et al., 2012; American Cancer Society [ACS], 2019; Roostaeian, et al., 2012).

Indications for implant reconstruction include: bilateral reconstruction; individuals requiring augmentation in addition to reconstruction; individuals not suited for long surgery; a lack of abdominal tissue; individual unwilling to have additional scars on either their back or abdomen; and a small breast mound with minimal ptosis. Relative contraindications to implant reconstruction include: young age (i.e., may need implant replaced multiple times); individual unwilling to follow up; very large or ptotic breast. The contraindications to implant reconstruction include: silicone allergy; fear of implants; previously failed implants; or need for adjuvant radiation therapy (Roehl, et al., 2012).

Surgical complications associated with breast implantation are similar to those encountered with other breast surgeries: infection, bleeding, change in nipple sensation (e.g., hypersensitivity or hyposensitivity), malposition, delayed healing, and anesthetic accidents.

Although implantable breast prostheses may be inserted for either reconstructive or cosmetic reasons, clinically significant post-implant complications may occur, necessitating removal of the implants. Local complications associated with implanted breast prostheses include: capsular contracture, persistent infection, silicone implant extrusion, tissue necrosis and silicone implant rupture. These conditions, when they become clinically significant, may require removal of the implant. Additionally, the presence of an implant may interfere with the diagnosis or treatment of breast cancer. Infections that may occur in or around an implant include wound infections, as well

as infections within a capsular contracture or as a result of a ruptured implant. Removal of the implant may be necessary when the infection does not respond to antibiotics. Unstable or weakened tissue and/or interruption in wound healing may result in the implant breaking through the skin or extrusion. Necrotic tissue may form around the implant, requiring implant removal. Silicone gel-filled implant rupture may cause the contents to leak into the surrounding tissues.

U.S. Food and Drug Administration (FDA): In the FDA labeling for approved breast implants, Mentor Corp., Santa Barbara, CA; Ideal Implant[®], Inc., Dallas, TX; Allergan Corp. (formerly Inamed), Irvine, CA and Sientra, Inc., Santa Barbara, CA are listed as manufacturers of silicone and saline breast implants.

FDA-approved saline-filled implants:

- Allergan Medical RTV Saline-Filled Breast Implant
- Ideal Implant Saline-Filled Breast Implant
- Mentor Saline-Filled and Spectrum™ Breast Implants

The FDA approved saline-filled breast implants for breast augmentation in women age 18 or older and for breast reconstruction in women of any age. They are also used in revision surgeries, which correct or improve the result of an original surgery.

FDA-approved silicone gel-filled breast implants:

- Allergan Natrelle[®]
- Allergan Natrelle[®] 410 Highly Cohesive Anatomically Shaped Silicone-Filled Breast Implant
- Mentor MemoryGel[®]
- Mentor MemoryShape™ Silicone Gel-Filled Breast Implant
- Sientra[®] Silicone Gel Breast Implant

The FDA labeling for silicone and saline breast implantation states breast implant surgery should not be performed in women with: an active infection, existing cancer or precancer of a breast that has not been adequately treated, or who are pregnant or nursing (FDA, 2020a).

In June 2011 (updated 2018), the FDA released a report updating the clinical and scientific information for silicone gel-filled breast implants, including preliminary safety data from studies conducted by the manufacturers as a condition of their November 2006 approval. The conclusion in the report states that, “Based on the totality of the evidence, the FDA believes that silicone gel-filled breast implants have a reasonable assurance of safety and effectiveness when used as labeled. Despite frequent local complications and adverse outcomes, the benefits and risks of breast implants are sufficiently well understood for women to make informed decisions about their use.” Manufacturers and physicians should continue to provide balanced and up-to-date information to women considering breast implants to help inform their decisions (FDA, 2018). On July 24, 2019, the FDA requested that Allergan, the manufacturer of a specific type of textured implant, recall specific models of its textured breast implants from the U.S. market due to the risk of breast implant-associated anaplastic large cell lymphoma (BIA-ALCL). The FDA's analysis was attributed to a new worldwide reported total of 573 unique BIA-ALCL cases including 33 patient deaths. Of the 573 cases of BIA-ALCL, 481 are reported to have Allergan breast implants at the time of diagnosis. (FDA, 2019b; FDA, 2019c).

Tissue Expanders

Following mastectomy, some individuals have inadequate elasticity in the remaining tissue to accommodate and support a breast implant. For these individuals, tissue expanders can be inserted under the chest muscle or skin. The expander is an empty balloon-like container that, over time, is injected with saline. This inflation causes the tissue to expand. The tissue expander is surgically removed once an adequate pocket has been established, and the permanent implant is then inserted. The most appropriate patients for this type of reconstruction are individuals who do not qualify for autogenous reconstruction, individuals who do not want additional scars from other donor sites, individuals who prefer a typically quicker postoperative recovery period, and individuals who have relatively small breasts. Contraindication for this type of reconstruction are mastectomy flaps that are too thin for adequate implant coverage and the completed or planned use of adjuvant radiation therapy because of higher implant complication rates (ACS, 2019; Hu, et al., 2007).

Tissue Flap Procedures

Autologous breast reconstruction procedures are safe and effective and are a well-established standard of care include tissue/muscle reconstruction procedures (e.g., flaps). Methods of autologous tissue breast reconstruction include local flaps and distant flaps. Local flaps rely on transposition of muscle, subcutaneous tissue, and skin into the mastectomy defect based on the attached native blood supply of the muscle (e.g., latissimus dorsi myocutaneous (LD) flap and the pedicled transverse rectus abdominus myocutaneous (TRAM) flap). Distant flap breast reconstruction requires the use of microvascular free-tissue transfer (e.g., free TRAM flap, deep inferior epigastric perforator [DIEP] flap, superficial inferior epigastric artery perforator [SIEP] flap, inferior or superior gluteal flap, superior gluteal artery perforator flap, Reubens flap, and the transverse upper gracilis (TUG) flap). Breast reconstruction using these donor sites relies on harvesting the flap with its vascular pedicle, which is anastomosed using microsurgical technique to appropriate recipient vessels in the mastectomy site. The two most common types of tissue flap procedures are the TRAM flap and the LD flap. Other tissue flap surgeries are more specialized, and may not be available everywhere. The choice of procedure for a given individual is affected by her age, her health, her contralateral breast size and shape, her personal preference, and the expertise of the reconstructive surgeon (ACS, 2019, Roehl, et al., 2012; Spear et al., 2007; Mehrara et al., 2006; Alderman et al., 2006; Garvey et al., 2006; Bajaj et al., 2006; Wechselberger, et al., 2004; Behnam et al., 2003).

Deep Inferior Epigastric Perforator (DIEP) Flap: A modification of the free TRAM flap is the deep inferior epigastric perforator (DIEP) flap. This flap does not harvest any muscle or fascia from the abdomen, and reportedly has significantly less donor-site morbidity than the usual TRAM flap. Patients are thought to have reduced postoperative pain, a lower risk of abdominal bulge or hernia, and less postoperative abdominal donor-site weakness. In reducing the amount of disturbance to the abdominal wall donor site, however, use of the DIEP flap unavoidably reduces the number of perforators supplying blood to the flap. This could potentially lead to a reduced supply of blood to the flap, thereby causing an increase in partial flap loss and fat necrosis (American Society of Plastic Surgeons [ASPS], 2017; Kroll, 2000).

Latissimus Dorsi Myocutaneous (LD) Flap: The LD flap moves muscle and skin from the back to reconstruct the breast. The LD flap is ideally suited for single-stage reconstruction for individuals with small breasts and a moderate degree of ptosis and for patients with no available abdominal donor site due to scars or lack of tissue. The LD flap can be used to correct lumpectomy defects which require a smaller implant or no implant. Some individuals may have weakness in their back, shoulder, or arm after this surgery. Relative contraindications to the LD flap include: planned postoperative radiation therapy, bilateral reconstruction, and significant breast ptosis. Contraindications to the LD flap include: previous lateral thoracotomy and individuals with large breast volume who do not desire reduction (Roehl, et al., 2012).

Rubens Flap: The Rubens flap is based on the circumflex iliac vessels and is an option for individuals who have an excess of soft tissue over the hips. Because this reconstructive procedure is limited in bulk and skin envelope, and often requires a balancing procedure on the contralateral hip, it is not usually considered as a first option for breast reconstruction (Roehl, et al., 2012).

Superficial Inferior Epigastric Perforator/Artery (SIEP/SIEA) Flap: The skin and fat of the lower abdomen are supplied by perforators (vessels that perforate the rectus abdominis muscle), including the superficial inferior epigastric artery (SIEA). For this type of reconstruction, an elliptical flap of tissue is transferred from the lower abdomen to the chest while still allowing a tension-free closure of the donor site in the abdomen. The apex of the triangular flap becomes the tail of the reconstructed breast. The internal mammary artery perforators or thoracodorsal vessels are often a good size match for the SIEA; these are anastomosed to the perforators of the graft using microsurgical technique. Construction of an SIEA flap presents several technical challenges and cannot be used in all cases (Hayes, 2014, reviewed 2016).

Superior or Inferior Gluteal Free Flap: The superior or inferior gluteal free flap requires skin, fat, blood vessels, and muscle is removed from the gluteus maximus to reconstruct the breast. This technique is an option for when the abdomen is no longer an alternative for flap transfer. This flap is technically complex and has complications including: seroma, sciatica, unfavorable scar location, and asymmetrical buttock contour (Roehl, et al., 2012).

Thoracodorsal Artery Perforator (TDAP) Flap: The TDAP flap is a rarely chosen source for autogenous tissue breast reconstruction. The TDAP flap is an evolution of the LD flap. The TDAP flap allows for collection of skin and soft tissue from the upper back without sacrifice of muscle tissue. The flap is based on proximal perforating vessels that originate from the thoracodorsal artery and vein. These vessels pass through the latissimus dorsi muscle and into the overlying skin and fat (DellaCroce, 2015).

Transverse Rectus Abdominus Myocutaneous (TRAM) Flap: The TRAM flap is the most commonly performed autologous reconstructive procedure and is considered the gold standard in breast reconstruction because of the lower abdominal tissue's similarities in consistency with breast tissue. There are three types of TRAM flaps: unipedicle, bipedicle, or free. Pedicle flaps involve leaving the flap attached to its original blood supply and tunneling it under the skin to the breast area. Free flap involves cutting the flap free of skin, fat, blood vessels, and muscle from its original location and attaching the flap to blood vessels in the chest area. The advantage of these procedures lies in the consistency of the reconstructed breast and its aesthetic appearance. These procedures are indicated for individuals with (Zenn, 2018, Roehl, et al., 2012):

- large tissue requirement after a radical mastectomy
- history of radiation to the chest wall
- small or large opposite breast that is difficult to match with an implant
- previous failure of implant reconstruction
- excess lower abdominal tissue

Abdominal complications resulting from this surgery include loss of abdominal strength, abdominal bulge and hernia formations. It is recommended that reconstruction be delayed when adjuvant chemotherapy is planned, as complications of the reconstruction can be detrimental in beginning the individual's therapy.

Numerous factors place an individual at higher risk for complications and are therefore considered relative contraindications to TRAM flap surgery (e.g., cardiac and/or pulmonary disease, diabetes, history of pulmonary embolus or deep venous thrombosis) (Zenn, 2018; ASPS, 2017; Roehl, et al., 2012):

Transverse Upper Gracilis (TUG) Flap: The TUG flap is taken from the upper inner thigh area. Part or all of the gracilis muscle is included with the flap to ensure the most reliable blood supply. This is a breast reconstructive method for those individuals who have limited flap donor sites. Candidates for TUG flap breast reconstruction include individuals desiring autogenous breast reconstruction with sufficient upper inner thigh tissue but who have had a previous abdominoplasty or a flap previously taken from their abdomen. Very thin or athletic individuals who may have insufficient abdominal donor tissue may be candidates for the TUG flap. This flap may be referred to as the TUG Perforator Flap which, as a perforator flap, it is a flap made of skin and fat only (no muscle). The TUG Myocutaneous Flap includes skin, fat, a portion of the gracilis muscle and the blood vessels associated with it to keep it alive. It is not usually considered as a first option for breast reconstruction.

Intraoperative Assessment of Tissue Perfusion

One of the reported causes of early complications following breast reconstructive procedures is considered to be inadequate tissue perfusion. Accurate and reliable intraoperative evaluation of tissue perfusion is needed to reduce complications and improve clinical outcomes. Besides clinical judgement, several technologies to assess tissue vascularity have been evaluated in studies and are used clinically (e.g., intraoperative laser angiography using indocyanine green (ICG); fluorescein, doppler) (Gurtner, et al., 2013). One device that is used is the SPY® Fluorescent Imaging System (Stryer, Kalamazoo, MI; formerly Novadaq Technologies Inc., Mississauga, Ontario) (FDA, 2008). Intra-operative assessment of tissue perfusion is considered an integral part of a breast reconstruction procedure.

Reconstruction of the Nipple-Areolar Complex

This portion of the breast reconstruction is usually performed as a second or third stage after the breast mound has been constructed. The recreation of the nipple-areolar complex involves various proposed techniques such as skin grafts, autologous and xenograft cartilage grafts, local tissue flaps, tissue-engineered structures, and tattooing and/or transplantation of nipple-areolar tissue from the opposite breast. It has been reported that within 12 months, most reconstructed nipples undergo a 50% reduction in projection. Therefore, the nipple should be made larger than desired during the initial surgery. The rebuilding of the nipple-areolar area is conducted first,

and the tattooing procedure is done when swelling has subsided, usually 3–6 weeks after nipple creation. Successful nipple-areola reconstruction is expected to maintain nipple projection and areola size; however, longevity of this reconstruction is highly variable and is influenced by factors such as tissue thickness, scar contracture, trauma and radiation. Tattooing is commonly repeated (Chun, 2017; Beckenstein, 2014; Roehl, et al., 2012; Heitland, et al., 2006; Guerra, et al., 2003).

Local tissue flaps are the most frequently performed methods of reconstruction. A common donor site is the medial thigh skin. Nipple reconstruction with local flaps is achieved with various techniques, each with its own proponents and benefits. These include the skate flap, bell flap, double opposing tab flap, star flap, top-hat flap, twin flap, propeller flap, S flap, rolled dermal-fat flap, and autologous cartilage. Acellular dermal matrices used alone or in conjunction with local flaps are being proposed as well as injectable materials for nipple reconstruction. Some have also advocated creating a more stable de-epithelialized skin base for the reconstructed nipple to minimize loss of projection (Chun, 2017; Beckenstein, 2014).

Loss of nipple projection commonly occurs a few years after reconstruction. This problem may be reduced with the use of bell and double opposing tab flaps. Secondary nipple reconstruction has been proposed by various procedures such as re-elevating the flap; inserting autologous dermal tissue, autologous or banked cartilage; and using filler injection or AlloDerm. Discoloration and uneven pigment distribution may occur over time and can usually be corrected with tattooing (Chun, 2017).

In a systematic review, Winocour et al. (2016) reported the efficacy, projection, and complication rates of different materials used in nipple reconstruction. A total of 31 retrospective and prospective studies with controlled and uncontrolled conditions reporting on outcomes of autologous, allogeneic, and synthetic grafts in nipple reconstruction were included. The authors reported heterogeneity in the type of material used within each category and inconsistent methodology used in outcomes assessment in nipple reconstruction. Overall, the quality of evidence is low. Synthetic materials have higher complication rates and allogeneic grafts have nipple projection comparable to that of autologous grafts. The authors reported that further investigation with high-level evidence is necessary to determine the optimal material for nipple reconstruction.

Xenograft Cartilage Grafting: The use of cartilage is another method of nipple reconstruction, particularly in prosthetic reconstruction where there might be a soft-tissue deficiency. The procedure is applicable to both unilateral and bilateral nipple reconstruction, is reported to be an easy procedure to perform, does not involve a donor site, and maintains long-lasting projection. A reported disadvantage of donated cartilage is that the resulting nipple is firm and unnatural in feel. If the grafts are placed too superficially and do not have a smooth contour, they can extrude through the skin, warranting revision and/or removal. Caution is recommended with thin skin flaps or irradiated tissue which can also make extrusion more likely. The use of simple nipple–areola tattooing is recommended for these patients. During the informed consent, it is recommended that the patient be warned that the cartilage is from an organ donor and there is a minor risk of infectious diseases (Beckenstein, 2014). Autologous cartilage grafting in breast reconstruction procedures is the standard of care. There is a lack of evidence in the peer reviewed published literature on the long-term outcomes, safety and efficacy of Xenograft cartilage use in breast reconstructive procedures.

Juvederm®: Juvederm (Allergan, Irvin, TX) Voluma® XC hyaluronic acid filler has been proposed to reshape nipples after reconstruction of the breast following mastectomy. On October 22, 2013, Juvederm Voluma XC received FDA premarket approval (PMA). The FDA indications for use state that the device is indicated for deep (subcutaneous and/or supraperiosteal) injection for cheek augmentation to correct age-related volume deficit in the midface in adults over the age of 21. Breast reconstruction is not specifically mentioned as an approved FDA indication in the FDA PMA approval order or in the PMA supplements (FDA, 2013). Evidence in the published, peer-reviewed scientific literature supporting the use of this product in breast reconstructive procedures is lacking and its role is unclear.

Radiesse®: Radiesse (BioForm Medical, Inc., San Mateo, CA) has been proposed to reshape nipples after reconstruction of the breast following mastectomy. Radiesse injections consist of very small, smooth calcium hydroxylapatite (CaHA) microspheres that are suspended in a water-based gel carrier. Radiesse has received PMA approval by the FDA as a medical device for subdermal implantation for two indications: correction of moderate to severe facial wrinkles and folds such as nasolabial folds and the correction of facial fat loss in

people with human immunodeficiency virus (FDA, 2006c). There remains a lack of evidence in the peer reviewed published literature on the long-term outcomes, safety and efficacy of Radiesse in breast reconstructive procedures.

Cook Biodesign® Nipple Reconstruction Cylinder: The Cook Biodesign Nipple Reconstruction Cylinder (Cook Biotech Incorporated, West Lafayette, IN) is a porcine non-cross-linked, non-dermis-based biologic graft material that is marketed for breast procedures including breast reconstruction, breast revision and mastopexy. It may be used in combination with Biodesign® Tissue Generation Matrix. On June 20, 2011, the Cook Biodesign Nipple Reconstruction Cylinder received FDA 510(k) approval. The FDA indications for use for the Biodesign Nipple Reconstruction Cylinder states it is intended for implantation to reinforce soft tissue where weakness exists, in plastic and reconstructive surgery of the nipple. The cylinder is supplied sterile and is intended for one-time use. The Biodesign Nipple Reconstruction Cylinder is a rolled Small Intestinal Submucosa (SIS) mesh and available in sizes from 0.7 cm to 1.0 cm in diameter and 1.0 cm to 2.5 cm in length. The cylinder is a scaffold which becomes infiltrated by the host cells during the body's natural repair process. The device is implanted using a skin flap procedure that prevents migration of the device. The clinical performance of the Biodesign Nipple Reconstruction Cylinder was assessed in two case studies and anecdotal evidence of 186 device implants. Of the 188 implants, complications included device extrusion (number of extrusions not given). Follow-up periods ranged from 2 to 12 months. The clinical studies showed the Biodesign Nipple Reconstruction Cylinder as substantially equivalent to its predicates in its application (FDA, 2011).

In the first multi-center prospective study, Collins et al. (2016) reported on the use of the Biodesign Nipple Reconstruction Cylinder (NRC) during reconstruction of the nipple after mastectomy in patients with a history of breast cancer and mastectomy. Unilateral or bilateral nipple reconstruction was performed. Skin flaps were raised, the NRC was placed beneath the flaps as a stent, and the site was protected for up to four weeks with a nipple shield. Nipple projection was measured for 12 months after surgery. Patient satisfaction was measured and adverse events were recorded. Follow-up examinations were performed at one week, and then at one, three, six, and 12 months after surgery. A total of 82 nipple reconstructions were performed in 50 patients. Related postoperative adverse events were minor, but reported in eight reconstructions (9.8%) representing seven patients (14.0%). Average projection at six and 12 months was 4.1 ± 1.6 mm and 3.8 ± 1.5 mm, respectively, compared with 10.5 ± 2.2 mm one week after surgery. Of patients completing the satisfaction questionnaire at 12 months, 70/75 (93.3%) of reconstructions were rated "pleased" or "very pleased" with the overall outcome. Overall, 45/46 (97.8%) patients would recommend nipple reconstruction to other women. This study is limited by the small homogenous sample size, lack of a control group and short term follow-up.

There is a lack of evidence in the peer reviewed published literature regarding the long-term outcomes and efficacy of the Cook Biodesign Nipple Reconstruction Cylinder for use in breast reconstruction or for any other indication.

Contralateral Breast

Although the goal of breast reconstruction is to restore symmetry, the process may leave the opposite or contralateral breast larger or smaller than the surgical breast. To correct this asymmetry, a mastopexy or reduction mammoplasty may be performed on the contralateral breast. If the reconstructed breast is larger, then an augmentation mammoplasty with implant may be performed on the nondiseased breast (Roehl, et al., 2012).

Oncoplastic Reconstruction

Oncoplastic procedures are performed immediately or one to two weeks after lumpectomy, once final pathology is available. They include rearrangement of the remaining breast tissue through a variety of techniques, often adhering to breast reduction principles. In addition, more tissue can be brought into the breast to correct the volume deficit, often in the form of a latissimus dorsi flap. Indications for these procedures depend on the patient's preoperative breast size, available remaining breast tissue, and overall goals for ultimate breast size and shape. All these procedures are done prior to radiation to prevent the contracture of the lumpectomy defect and distortion of the nipple-areolar complex (Roehl, et al., 2012).

Radiation Tattoo Markers: Ink markers are tattooed as landmarks before radiotherapy of breast cancer with the purpose of obtaining a precise radiation field. These tattoos are permanent, but are the size of a freckle.

Individuals may have these tattoo markers removed via laser or punch biopsy excision as a part of the overall breast reconstruction procedure (Bregnhøj, et al., 2010).

Nonsurgical Options

Some women may choose not to have breast reconstruction or are poor candidates for reconstruction. For these women, an external breast prosthesis and mastectomy bras are additional options (Hu, et al., 2007).

Skin Substitutes

During breast reconstruction, acellular dermal skin substitutes (i.e., AlloDerm®, AlloMax™, DermACELL® and FlexHD®) are primarily used in the setting of tissue expander and breast implant reconstruction. Patients should be in overall good health and have no underlying condition that would restrict blood flow or interfere with the normal healing process (e.g., uncontrolled diabetes, hypertension, previous surgery). These matrixes may be indicated when there is insufficient tissue expander or implant coverage by the pectoralis major muscle and additional coverage is required, as may be the case in a very thin patient; if there is viable but compromised or thin post-mastectomy skin flaps that are at risk of dehiscence or necrosis; or if there is a need to re-establish the inframammary fold and lateral mammary fold landmarks. When used in appropriate candidates, these skin substitutes are proposed to improve control over placement of the inframammary fold and final breast contour, enhance use of available mastectomy skin, reduce the number of expander fills necessary, reduce time to complete expansion and eventual implant exchange, potential improved management of a threatened implant, reduce the need for explantation and the potential for reduction in the incidence of capsular contracture. However, there are ongoing concerns regarding the increased risk of seroma and infection, a higher risk of an implant having to be removed, and tissue flap death (ACS, 2019; Nguyen, et al., 2011; Sbitany and Serletti, 2011).

There is a paucity of data comparing the skin substitute products directly. The products vary in many aspects, including the source of tissue, processing, storage, surgical preparation and available sizes. The most familiar product to most plastic surgeons, AlloDerm, was the first human dermis product available in 1994 (Cheng, et al., 2012).

U.S. Food and Drug Administration (FDA)

Depending on the purpose of the product and how it functions, skin substitutes are regulated by the FDA premarket approval (PMA) process, 510(k) premarket notification process, or the FDA regulations for banked human tissue.

Products that are classified by the FDA under the PMA process as a class III, high-risk device require clinical data to support their claims for use. These devices may be used as a long-term skin substitute or a temporary synthetic skin substitute. They actively promote healing by interacting directly or indirectly with the body tissues.

Other wound care devices are approved by the 510(k) process, and their primary purpose is to protect the wound and provide a scaffold for healing. They may or may not be integrated into the body tissue. Some devices are rejected by the body after approximately ten days to several weeks and removed prior to definitive wound therapy or skin grafting.

The following is a list of skin substitutes, not derived from human tissue, that are FDA approved (this list may not be all-inclusive):

- Permacol
- Strattice
- SurgiMend

Donated skin that requires minimal processing and is not significantly changed in structure from its natural form is classified by the FDA as banked human tissue, is not considered a medical device, and does not require PMA or 510(k) approval. Donated skin (human cells or tissue) intended for implantation, transplantation, infusion, or transfer into a human recipient is regulated as a human cell, tissue, and cellular and tissue-based product (HCT/P) by the Center for Biologics Evaluation and Research (CBER) (FDA, 2021). The American Association of Tissue Banks (AATB) sets standards for the safety and use of donated human tissue (AATB, 2019). AATB

oversees a voluntary accreditation program and the FDA focuses on preventing the transmission of communicable diseases by requiring donor screening and testing. Tissue establishments must register with the FDA and list each cell or tissue produced. An example of a banked human tissue product is AlloDerm, an acellular dermal matrix (FDA, 2004).

The following skin substitutes are derived from human tissue and therefore subject to the rules and regulations for banked human tissue developed by the American Association of Tissue Banks (AATB) (this list may not be all-inclusive):

- Alloderm
- Allomax
- Cortiva
- DermACell
- DermaMatrix
- FlexHD
- hMatrix
- Repriza

The safety and efficacy of the skin substitutes listed below are supported by the evidence in the published peer-reviewed scientific literature and/or are established treatment options for post-mastectomy breast reconstruction.

AlloDerm™: AlloDerm (Allergan™, Parsippany, NJ [formerly LifeCell™ Corporation, Branchburg, NJ]) is an acellular dermal matrix allograft classified as banked human tissue by the FDA because it is minimally processed and not significantly changed in structure from the natural material. AlloDerm is an established treatment option and is supported by the evidence in the published peer-reviewed scientific literature for tissue repair during postmastectomy breast reconstruction (Lee, et al., 2018; McCarthy, et al., 2012; Cheng, et al., 2012; Vardanian, et al., 2012; Jansen and Macadam, 2011; Salzberg, et al., 2011; Joanna, et al., 2011; Antony, et al., 2010; Haddock, et al., 2010; Spear, et al., 2008; Bindingnavele, et al., 2007; Breuing and Colwell, 2007; Zienowicz, et al., 2007; Gamboa-Bobadilla, 2006; Glasberg, et al., 2006; Salzberg, 2006; Breuing, et al., 2005; Nahabedian, 2005).-Various forms of AlloDerm are available including AlloDerm™ Regenerative Tissue Matrix, AlloDerm Select™ Tissue Matrix and AlloDerm Select Duo™ Tissue Matrix Bilateral Pair (Allergan, 2020; Hayes, 2019, reviewed 2020).

AlloMax™: AlloMax Surgical Graft (Bard Davol, Inc. Warwick, RI) is an acellular non-cross-linked human dermis allograft. Because AlloMax is a natural human product it is classified as banked human tissue and does not require FDA approval. It is regulated by the American Association of Tissue Banks and the FDA guidelines for banked human tissue. AlloMax Surgical Graft is available in multiple sizes. The AlloMax Surgical Graft for Breast Reconstruction (previously marketed as NeoForm™) is proposed for post-mastectomy breast reconstruction and is an established skin substitute for this indication (Venturi, et al., 2013; Bard, 2017).

Cortiva®

Cortiva (RTI Surgical, Alachua, FL) is a non-crosslinked, cadaveric human acellular dermal matrix processed by Tutoplast technology using low-dose gamma irradiation. The matrix is FDA regulated as human cell, tissue, and cellular and tissue-based product (361 HCT/P) and proposed for the repair, replacement, reconstruction or augmentation of soft tissue, including supplemental support and reinforcement of soft tissue in breast reconstruction and hernia repair. There are three products: Cortiva, Cortiva 1.0 mm and Cortiva 1 mm tailored allograft dermis. The matrixes are offered in regular and 1 mm thicknesses and supplied in a range of sizes from 2x4 cm to 16x20 cm (RTI, Inc., 2021; Centers for Medicare and Medicaid Services [CMS], 2015). Cortiva has evolved into an acceptable tissue substitute for breast reconstruction and a randomized controlled trial with short-term follow-up reported that outcomes with Cortiva were not inferior to outcomes using AlloDerm.

Parikh, et al. (2018) reported the outcomes of a phase 2 randomized controlled trial that compared outcomes following breast reconstruction surgery using Cortiva 1 mm allograft or AlloDerm Ready to Use (RTU) regenerative tissue matrix. The 16x8 cm graft was used as a sling to support tissue expanders placed in the submuscular location in one study arm, and prepectoral reconstructions with tissue expanders (TEs) or direct-to-

implants (DTI) in a second study arm. The interim analysis of the submuscular reconstruction group is reported herein. Breasts reconstructed with AlloDerm RTU (n=17 patients; 28 breasts) or Cortiva 1 mm (n=17 patients; 31 breasts) submuscular TE, completed the interim analysis. During the study a significant shift to prepectoral reconstructions was noted and the prepectoral arm of the study was added to optimize enrollment rates. Patients who underwent prepectoral breast reconstruction with either DTI or TE supported by a 20x16 cm ADM sheet were compared in a separate study arm. The decision to proceed with prepectoral or submuscular reconstruction with either a TE or DTI was determined preoperatively. Female patients, aged 22–70 years old, undergoing immediate prosthetic reconstruction following therapeutic or prophylactic skin- or nipple-sparing mastectomy with a body mass index (BMI) less than 36 kg/m² were included. Excluded patients were those who were pregnant or breastfeeding immediately before mastectomy. The primary outcome measure was premature explantation of the TE before exchange, or unintended explantation of a DTI reconstruction during the first three months postoperatively. Secondary outcome measures included other complications (e.g., seroma, cellulitis, wound or ADM dehiscence, skin flap necrosis). Patients undergoing TE placement in either study arm were followed until there was TE exchange with an implant, flap, or both, or there was premature removal of the device. Patients undergoing DTI reconstruction were followed for at least three months following surgery. Patients undergoing reoperation of the surgical site without device exchange or removal were kept in the study. Patients undergoing planned exchange of TEs for implants or flaps within 145.6 ± 51.6 days in the AlloDerm group and 167.0 ± 61.5 days in the Cortiva 1 mm group (p=0.27), not statistically significant. Most patients were exchanged with breast implant alone, but 14.3% in the AlloDerm group and 26.6% in the Cortiva group (p=0.25) received an autologous flap, not statistically significant. There was no significant difference between the groups in integration of the ADM to the mastectomy flap (p=0.69), in drain removal between the groups or in physical well-being, or satisfaction with information or plastic surgeon. A significant difference was seen in detectable seroma in the AlloDerm (n=3) vs. the Cortiva group (n=0). Premature explantation was performed in no AlloDerm breast vs. one breast with Cortiva. The initial size of the TE selected was significantly larger in patients reconstructed with Cortiva 1 mm (p=0.02). The AlloDerm RTU group was comprised of a significantly higher proportion of patients who had never smoked (p=0.009). This interim analysis of submuscular reconstructions revealed no evidence of inferiority of outcomes of AlloDerm vs. Cortiva. Limitations of the study include the small patient population and short-term follow-up.

DermACELL™: DermACELL (LifeNet Health®, Virginia Beach, VA) is an acellular human dermis allograft collagen scaffold proposed for the treatment of breast reconstruction. LifeNet Health is registered with the FDA as an establishment producing tissue- and cellular-based products. MatrACELL® is a patented process that removes > 97% of donor DNA that renders Demacell acellular. Terminal sterilization is performed by low dose gamma irradiation. The use of DermACELL for breast reconstruction has evolved into an accepted standard of practice. Although the evidence supporting DermACELL for breast reconstruction is primarily in the form of case series and retrospective reviews, outcomes reported a significant improvement in time to drainage removal and fewer “red breast episodes” compared to AlloDerm (Pittman, et al., 2016). Zenn et al. (2016) reported that DermACELL was as good as AlloDerm RTU in the occurrence of postoperative infection, implant loss, seroma and hematoma. Other studies have also reported favorable outcomes with DermACELL (Bullocks, et al., 2014; Vashi, 2014). Therefore, DermACELL has evolved into an accepted skin substitute for breast reconstruction (Ortiz, 2017; Chang and Liu, 2017; Pittman, et al., 2016; Zenn, et al., 2016; Bullocks, et al., 2014; Vashi, 2014).

FlexHD® Acellular Hydrated Dermis: FlexHD Acellular Hydrated Dermis (Musculoskeletal Transplant Foundation, Edison, NJ) is a matrix derived from donated human allograft skin. The product is regulated by the American Association of Tissue Banks and the FDA guidelines for banked human tissue. FlexHD is indicated for the replacement of damaged or inadequate integumental tissue or for the repair, reinforcement or supplemental support of soft tissue defects. Flex HD is available in multiple sizes. Results of case series and retrospective reviews in the peer-reviewed literature support the safety and efficacy of FlexHD for use during postmastectomy breast reconstruction. FlexHD is an established skin substitute for this indication (Lee, et al., 2018; Liu, et al., 2014; Seth, et al., 2013; Seth, et al., 2012; Brooke, et al., 2012; Rawlani, et al., 2011; Cahan, et al., 2011; Topol, et al., 2008).

NeoForm™ Dermis: Neoform Dermis (Mentor Corp., Santa Barbara, CA) is a solvent-dehydrated, gamma-irradiated preserved human allograft dermis indicated for use as a soft tissue graft for horizontal and vertical soft tissue augmentation of thickness and length, such as breast reconstruction. NeoForm is classified as banked human tissue by the FDA. Although evidence in the published, peer-reviewed scientific literature supporting the

use of this product in breast reconstruction is limited, Neoform Dermis is an established skin substitute used for tissue expansion in breast reconstruction following a mastectomy. Neoform is no longer available for distribution.

Other Skin Substitutes

Additional skin substitutes have been proposed as treatment options in breast reconstruction as discussed below, but the evidence in the published peer-reviewed scientific literature does not support the safety and efficacy of the use of these substitutes. The number of available studies is limited and involves small, heterogeneous patient populations, short-term follow-ups, minimal comparisons to the established treatment method for the condition, and/or lack of a control group. In some cases, reported outcomes are inconsistent, and a consensus on patient selection criteria and the appropriate surgical approach and techniques that should be used have not been established.

ARTIA™ Reconstructive Tissue Matrix: ARTIA Reconstructive Tissue Matrix (Artia Tissue Matrix)/ ARTIA Tissue Matrix-Perforated (Allergan™, Parsippany, NJ [formerly LifeCell™ Corporation, Branchburg, NJ]) is a surgical mesh that is derived from porcine skin that is processed and preserved in a patented phosphate buffered aqueous solution containing matrix stabilizers. ARTIA was originally developed by LifeCell Corporation and is currently distributed by Allergan. According to the FDA 510(k) approval, ARTIA Tissue Matrix is intended for use as a soft tissue patch to reinforce soft tissue where weakness exists and for the surgical repair of damaged or ruptured soft tissue membranes which require the use of reinforcing or bridging material to obtain the desired surgical outcome. The implant is intended for reinforcement in plastic and reconstructive surgery (FDA, 2017). There is insufficient evidence to support the safety and efficacy of ARTIA Reconstructive Tissue Matrix as a skin substitute for breast reconstructive surgery.

BellaDerm® Acellular Hydrated Dermis: BellaDerm Acellular Hydrated Dermis (Musculoskeletal Transplant Foundation, Edison, NJ) is human allograft skin minimally processed to remove epidermal and dermal cells and is packaged in an ethanol solution. The process utilized preserves the extracellular matrix of the dermis. The resulting allograft serves as a framework to support cellular repopulation and vascularization at the surgical site. BellaDerm is processed to remove cells while maintaining the integrity of the matrix with the intent to address the issues of the specific and nonspecific inflammatory responses. It is used for the replacement of damaged or inadequate integumental tissue or for the repair, reinforcement or supplemental support of soft tissue defects. Grafts of BellaDerm can range between 0.8mm and 1.7mm. It is used for breast augmentation revision procedures, including correction of symmastia, capsular contracture, bottoming out and malposition. There is insufficient evidence to support the safety and efficacy of BellaDerm Acellular Hydrated Dermis as a skin substitute for breast reconstructive surgery.

DermaMatrix Acellular Dermis: DermaMatrix (formerly manufactured by Synthes Inc., West Chester, PA) is an allograft derived from human skin and is classified by the FDA as banked human tissue. DermaMatrix is proposed for use for breast reconstruction postmastectomy. Per the manufacturer, as of June 2014, DermaMatrix is no longer available for distribution (FDA K142326).

GalaFlex® Surgical Scaffold/GalaFLEX Mesh: This surgical scaffold/mesh (Tepha, Inc., Lexington, MA) is a sterile, knitted, resorbable mesh, constructed of non-dyed monofilament fibers made from poly-4-hydroxybutyrate (P4HB). P4HB is produced from a naturally occurring monomer and is processed into monofilament fibers and knitted into a surgical fold. It is provided in single sheets of varying widths, lengths and shapes, and may also be cut to the shape or size desired for a specific application. According to the FDA 510(k) approval GalaFLEX Mesh is indicated for use “as a transitory scaffold for soft tissue support and to repair, elevate and reinforce deficiencies where weakness or voids exist that require the addition of material to obtain the desired surgical outcome. This includes reinforcement of soft tissue in plastic and reconstructive surgery, and general soft tissue reconstruction”. In November 2017, Hayes published a Clinical Research Response for this product concluding that “Based on a review of published abstracts, there is insufficient evidence to inform conclusions regarding the efficacy of GalaFLEX Surgical Scaffold for use as reinforcement in soft tissue reconstruction”. There is insufficient evidence to support the safety and efficacy of GalaFLEX Mesh as a skin substitute for breast reconstructive surgery (Hayes, 2017; Williams, et al., 2016; FDA, 2014).

GalaFORM™ 3D: GalaFORM 3D (Tepha, Inc., Lexington, MA) is a bioresorbable surgical mesh manufactured from P4HB. According to the FDA 510(k) approval, GalaFORM 3D scaffold is indicated for use as a

bioresorbable scaffold for soft tissue support and to repair, elevate and reinforce deficiencies where weakness or voids exist that require the addition of material to obtain the desired surgical outcome. This includes reinforcement of soft tissue in plastic and reconstructive surgery, and general soft tissue reconstruction. GalaFORM 3D scaffold is also indicated for the repair of fascial defects that require the addition of a reinforcing or bridging material to obtain the desired surgical result. The predicate device is GalaFLEX™ Scaffold (FDA, 2017). There is insufficient evidence to support the safety and efficacy of GalaFORM 3D scaffold as a skin substitute for breast reconstructive surgery.

GalaSHAPE 3D, GalaFlex and GalaForm 3D: GalaSHAPE 3D (Tepha, Inc., Lexington, MA) is a 3-dimensional monofilament scaffold made from poly-4-hydroxybutyrate (P4HB). After implantation, the scaffold is proposed to gradually bioresorb over 18-24 months providing a lattice for new tissue ingrowth and regeneration. The FDA 510(k) approval indications for use state that “GalaSHAPE™ 3D is indicated for use as a bioresorbable scaffold for soft tissue support and to repair, elevate and reinforce deficiencies where weakness or voids exist that require the addition of material to obtain the desired surgical outcome. This includes reinforcement of soft tissue in plastic and reconstructive surgery, and general soft tissue reconstruction. GalaSHAPE™ 3D is also indicated for the repair of fascial defects that require the addition of a reinforcing or bridging material to obtain the desired surgical result”. GalaFLEX scaffold and GalaFORM 3D are also approved for the same indication. (FDA, 2017; FDA, 2016; FDA 2014; Galatea Surgical, 2021). The Galatea products are available in various sizes in oval, rectangular, triangular, circular shapes and can be custom made. There is insufficient evidence to support the safety and efficacy of Galatea products as a skin substitute for breast reconstructive surgery.

hMatrix®: hMatrix Acellular Dermis (Bacterin International Holdings Inc., Belgrade, MT) is an acellular dermal scaffold processed from donated human skin. The dermis is processed using a proprietary method and the matrix is packaged and sterilized using low-dose gamma irradiation. hMatrix is regulated by the American Association of Tissue Banks and the FDA guidelines for banked human tissue. The product is stored and supplied frozen. Bacterin hMatrix PR for breast augmentation (Bacterin, 2013). hMatrix is available in four sizes. There is insufficient evidence to support the safety and efficacy of hMatrix as a skin substitute for breast reconstruction.

Permacol®: The Permacol Crosslinked Porcine Dermal Collagen Surgical Mesh (Tissue Sciences Laboratories PLC, Hants, United Kingdom), a xenograft, is a fibrous flat sheet comprised of acellular porcine dermal collagen and elastin. According to the FDA 510(k) approval, Permacol® is intended for use as a soft tissue patch to reinforce soft tissue where weakness exists and for the surgical repair of damaged or ruptured soft tissue membranes (FDA, 2000). Breast reconstruction is not specifically mentioned as an approved FDA indication. However, muscle flap reinforcement is an FDA-approved indication for use. Permacol is available in multiple sizes. Evidence in the published, peer-reviewed scientific literature supporting the use of this product in breast reconstruction is lacking and its role is unclear (Knabben, et al., 2017; Ramsden, et al., 2009).

Phasix™ Mesh: Phasix Mesh (Bard Davol, Inc. Warwick, RI), is a fully resorbable P4HB polymer material proposed for use in breast reconstructive procedures. The P4HB is produced from a naturally occurring monomer and is processed into monofilament fiber then knitted into a surgical mesh. Phasix Mesh received FDA 510k approval on March 31, 2015 to reinforce soft tissue where weakness exists in patients undergoing plastic and reconstructive surgery, or for use in procedures involving soft tissue repair, such as the repair of hernia or other fascial defects that require the addition of a reinforcing or bridging material to obtain the desired surgical result (FDA, 2015). The FDA approved indication for Phasix has no specific language related to breast reconstruction. Phasix is available in multiple sizes as round, rectangular, and square implants (Hayes, 2019, reviewed 2020). Evidence in the published, peer-reviewed scientific literature supporting the use of this product in breast reconstruction is lacking and its role is unclear.

Repriza® Acellular Dermal Matrix: Repriza (Promethean Lifesciences Inc., Pittsburg, PA) is a human skin, acellular dermal matrix. The product is regulated by the American Association of Tissue Banks and the FDA guidelines for banked human tissue. Repriza is membrane free and proposed for implantation for reconstructive surgery including breast reconstruction, abdominal wall reconstruct and augmentation of soft tissue irregularities. The matrix is provided in 4X12 cm and 6X16 cm sheets. It can also be custom made. There is insufficient evidence to support the safety and efficacy of Repriza for breast reconstructive surgery.

SERI™ Surgical Scaffold: SERI Surgical Scaffold (Sofregen Medical Inc., Cambridge, MA formerly Allergan, Medford, MA) has been proposed as a skin substitute for breast reconstruction (Jewell, et al., 2015). Breast reconstruction is not specifically mentioned as an approved FDA indication. The FDA 510(k) summary states that SERI is a “knitted, multi-filament, bioengineered, long-term bioresorbable scaffold. It is derived from silk that has been BIOSILK™ purified to yield ultra-pure fibroin. The device is a mechanically strong and biocompatible bioprotein” (FDA, 2013). The FDA indications for use state, “SERI Surgical Scaffold is indicated for use as a transitory scaffold for soft tissue support and repair to reinforce deficiencies where weakness or voids exist that require the addition of material to obtain the desired surgical outcome. This includes reinforcement of soft tissue in plastic and reconstructive surgery, and general soft tissue reconstruction” (FDA, 2013). There is insufficient evidence in the published peer-reviewed scientific literature supporting the efficacy of SERI Surgical Scaffold for breast reconstruction.

In May 2015 the FDA issued a warning letter to Allergan stating that the FDA approval of SERI Surgical Scaffold did not include the use of SERI Surgical Scaffold for breast reconstruction. Per the FDA, this indication falls outside of the intended use “because surgical mesh has not been cleared or approved for use in breast reconstruction using a tissue expander or implant”. The FDA requested Allergan “immediately cease activities that result in the misbranding or adulteration of the SERI Surgical Scaffold” for breast reconstruction” (FDA, 2015).

Strattice™ Reconstructive Tissue Matrix or LTM Surgical Mesh: Strattice Reconstructive Tissue Matrix or LTM Surgical Mesh (Allergan™, Parsippany, NJ [formerly LifeCell™ Corporation, Branchburg, NJ]) is an acellular, xenographic tissue matrix derived from porcine dermis (FDA, 2007). It is FDA 510(k) approved as LTM-RC surgical mesh “for use as a soft tissue patch to reinforce soft tissue where weakness exists and for the surgical repair of damaged or ruptured soft tissue membranes. The implant is intended for the reinforcement of soft tissues repaired by sutures or suture anchors, during rotator cuff surgery. Indications for use also include the repair of hernias and/or body wall defects which require the use of reinforcing or bridging material to obtain the desired surgical outcome” (FDA, 2007). Strattice is proposed for use during postmastectomy breast reconstruction to support medial repair for breast pocket size and position. Strattice is available in seven sizes with various shapes. There is insufficient evidence in the published peer-reviewed scientific literature supporting the efficacy of Strattice. Studies are primarily in the form of retrospective reviews (Dikmans, et al., 2016; Barber, et al., 2015; Lardi, et al., 2014; Kilchenmann, et al., 2014; Maxwell, et al., 2014; Glasberg and Light, 2012).

In June 2015 the FDA issued a warning letter to LifeCell Corporation stating that the FDA approval for Strattice did not include the use of Strattice for breast reconstruction. Per the FDA, this indication falls outside of the intended use “because surgical mesh has not been cleared or approved for use in breast reconstructive surgery applications”. The FDA requested that LifeCell “immediately cease activities that result in the misbranding or adulteration of the Strattice Reconstructive Tissue Matrix” for breast reconstruction.

SurgiMend® Collagen Matrix: SurgiMend or SurgiMend Collagen Matrix (Integra® LifeSciences Corp., Plainsboro, NJ formerly TEI Biosciences Inc., Boston, MA) is an acellular dermal tissue matrix derived from fetal or neonatal bovine dermis. The matrix acts as a scaffold that is progressively integrated, remodeled, and replaced by the functional host tissue. Approved as a Class II, FDA 510(k) device, SurgiMend is “intended for implantation to reinforce soft tissue where weakness exists and for the surgical repair of damage or ruptured soft tissue membranes” specifically for plastic and reconstructive surgery, muscle flap reinforcement, and hernia repair (e.g., abdominal, inguinal, femoral, diaphragmatic, scrotal, umbilical, incisional) (FDA, 2009). SurgiMend is available in multiple sizes and thicknesses. SurgiMend PRS, a pure collagen product, is designed for plastic and reconstructive surgery and is available in multiple shapes, sizes and thicknesses (Integra LifeSciences Corp, 2017; Butterfield, et al., 2013, Gaster, et al., 2013, Ohkuma, et al., 2013; Endress, et al., 2012; Craft, et al., 2011; Cromwell, et al., 2009).

TEI has historically marketed SurgiMend for breast reconstruction. In May 2015, the FDA issued TEI a warning letter stating that TEI did not have FDA clearance or approval to market SurgiMend for breast reconstruction. Per the FDA, this indication falls outside of the intended use “because surgical mesh has not been cleared or approved for use in breast reconstructive surgery applications”. The FDA requested that TEI “immediately cease activities that result in the misbranding or adulteration of SurgiMend” for breast reconstruction (FDA, 2015).

Veritas Collagen Matrix: Veritas Collagen Matrix (Synovis® Surgical Innovations, St. Paul, MN) is an implantable noncrosslinked biologic mesh made from bovine pericardium. Veritas is FDA approved as a surgical mesh under the 510(k) process for use as an implant for surgical repair of soft tissue deficiencies including: muscle flap reinforcement. There is also a Veritas Collagen Matrix Dry product that is FDA approved as a predicate device for the conventional Collagen Matrix (FDA, 2008; FDA, 2006). There is insufficient evidence to support the use of Veritas Collagen Matrix. The limited number of published studies investigating is primarily in the form of retrospective reviews.

Systematic Review and Meta-Analysis: Ho et al. (2012) conducted a systematic review and meta-analysis to determine an aggregate estimate of risks associated with acellular dermal matrix (ADM)-assisted breast reconstruction. AlloDerm was used in the majority of studies. ADMs other than AlloDerm were used in one study (i.e., FlexHD, Strattice). Seven complications were studied including seroma, cellulitis, infection, hematoma, skin flap necrosis, capsular contracture, and reconstructive failure. Sixteen studies met the inclusion criteria. The pooled complication rates were seroma 6.9%, cellulitis 2.0%, infection 5.7%, skin flap necrosis 10.9%, hematoma 1.3%, capsular contracture 0.6% and reconstructive failure 5.1%. Five studies reported findings for both the ADM and non-ADM patients and were used in the meta-analysis to calculate pooled odds ratio (OR). ADM-assisted breast reconstructions had a higher likelihood of seroma, infection, and reconstructive failure than breast reconstructions without the use of ADM. The relation of ADM use to hematoma, cellulitis, and skin flap necrosis was inconclusive. In the studies evaluated, ADM-assisted breast reconstructions exhibited a higher likelihood of seroma, infection, and reconstructive failure than prosthetic-based breast reconstructions using traditional musculofascial flaps. ADM is associated with a lower rate of capsular contracture. The authors reported that given the relatively low quality evidence that currently exists in the literature, additional randomized-controlled studies are needed to further evaluate the safety and efficacy of ADM in implant-based breast reconstruction.

Kim et al. (2012) conducted a systematic review and meta-analysis to evaluate complication outcomes from human acellular dermal matrix (ADM) used as an adjunct to traditional submuscular tissue expander/implant breast reconstruction. Forty eight uncontrolled cohort studies met inclusion criteria. Thirteen studies had information only on human ADM matrix-based reconstruction, 29 had information only on submuscular-based reconstructions, and six reported complications on human ADM and submuscular techniques. A total of 2037 human ADM reconstructions and 12,847 submuscular reconstructions were included in the meta-analysis. A total of 877 human ADM and 3464 muscular reconstructions from six studies were used to calculate relative risks. Average follow-up time was 13.8 months in the human ADM group and 28.3 months in the submuscular cohort. There was an increased rate of total complications, 15.4% vs. 14.0%; seroma, 4.8% vs. 3.5%; infection, 5.3% vs. 4.7%; and flap necrosis, 6.9% vs. 4.9% in the human ADM cohort compared to the submuscular reconstruction cohort. The rate of hematomas was greater in the submuscular cohort 1.5% vs. 1% in the human ADM. Meta-analysis revealed an increase in the risk of total complications (relative risk, 2.05; 95 percent confidence interval, 1.55 to 2.70), seroma (relative risk, 2.73; 95 percent confidence interval, 1.67 to 4.46), infection (relative risk, 2.47; 95 percent confidence interval, 1.71 to 3.57), and reconstructive failure (relative risk, 2.80; 95 percent confidence interval, 1.76 to 4.45) in the human ADM cohort. There was a trend toward increased risk of hematoma (relative risk, 2.06; 95 percent confidence interval, 0.86 to 4.95) and flap necrosis (relative risk, 1.56; 95 percent confidence interval, 0.85 to 2.85) in the human acellular dermal matrix cohort, but the results were not statistically significant. The majority of pooled complication analyses showed significant heterogeneity. The meta-analysis suggested that the use of human ADM increases complication rates compared to submuscular approach.

Nguyen et al (2011) conducted a systematic review of the literature to assess the quality and quantity of evidence regarding the use of acellular dermal matrices (ADM) (e.g., AlloDerm) in prosthetic breast reconstruction. Eighteen articles in the form of case reports, prospective cohort studies and retrospective reviews met inclusion criteria. The authors analyzed the evidence for the following proposed advantages of ADM: decrease or eliminate the need for expanders to create a tissue pocket for an implant; reduction in postoperative pain; decrease in operative time; increased initial fill volumes; fewer expansions; precise control of over the lateral and inframammary fold; ability to use more of the mastectomy skin flaps; faster time to completion of reconstruction; improved lower pole expansion; decreased incidence of capsular contractures; decreased rate of revisions; improved aesthetic outcome of the breast. The authors concluded that there was insufficient data to

support any of the above proposed benefits of ADM for breast reconstruction due to the paucity of actual data and conflicting data. Some studies did not formally quantify or report applicable data; evidence was inconsistent due to the variations in size of the matrix, type of mastectomy, size and viability of the mastectomy flaps and surgeon judgment; and data was conflicting due to variations in surgical technique, patients' physical characteristics, and number of expansions used.

Use Outside of the US

Skin substitutes are offered outside the US by several companies. Several products have received CE Mark approval (e.g., Integra® SurgiMend PRS Meshed, Strattice, Veritas Collagen Matrix). Products approved in the US may not be approved for use outside of the US and products approved outside the US may not be approved for use in the US. Also, the approved indications for the products may not be the same within and outside of the US. Integra LifeSciences has CE Mark approval for Surgimend PRS Mesh for pre- and sub-pectoral breast reconstruction in Europe. Strattice has CE Mark approval by the Netherlands-based notified body, KEMA, for its Strattice® Reconstructive Tissue Matrix. The CE Mark allows Strattice to be marketed in 27 European Union member states. Strattice is proposed for use in breast reconstruction. Native® (mbp, Germany) porcine acellular dermal matrix is proposed for use in Europe for breast reconstruction.

Autologous Fat Transplant (Lipoinjection, Lipofilling, Lipomodelling)

Despite various techniques of reconstruction with autologous tissues or prostheses, autologous fat transplant (i.e., lipoinjection, lipofilling, lipomodelling) has been proposed to replace volume after breast reconstruction or to fill defects in the breast following breast conservation surgery. It has been proposed as a stand-alone procedure or as an adjunct to other breast reconstruction techniques. The use of autologous fat transplant for primary breast augmentation is controversial due to a lack of clarity regarding its safety and efficacy. It has been proposed that injection of fat into a previous tumor site may create a new environment for cancer and adjacent cells. Additionally, fat transplant to the breast has been discouraged since it has been reported that calcifications secondary to fat necrosis may mimic breast cancer or that radiological changes post fat grafting would obscure and delay the diagnosis of subsequent breast cancer. In breast reconstruction, unlike elsewhere in the body, fat is implanted in a poorly vascularized and loose space which may attribute to the poor results.

Autologous fat transplant generally involves the transfer of fat from the abdomen or thighs into the breast under local anesthesia. The harvested fat is injected into the breast, usually in small parcels and thin strips, at different levels in the subcutis. It has been reported that a certain amount of fat resorption is expected in all cases of fat grafting. Clinically, volume loss has been reported between 40%-60% and usually within the first 4-6 months. Patients usually have 2-4 sessions of lipomodelling depending on their condition. The proposed benefit of the procedure is that it can restore volume to the breast without the morbidity associated with other reconstruction techniques. Although refinement in technique has aided reproducibility of favorable results, a standardized method of fat harvest, preparation, and injection is needed. Results are dependent on technique and surgeon expertise. It is recommended that breast reconstruction using autologous fat transfer be carried out by surgeons with specialist expertise and training in the procedure.

Literature Review: The available literature consists mostly of case reports, case series and expert opinion and describes autologous fat transplant for various breast indications, both cosmetic and reconstructive. Although the published evidence supporting the role of autologous fat transplant as a breast reconstruction procedure is not robust, limited data from several small studies indicate that autologous fat transplant raises no major safety concerns and may improve outcomes in a carefully selected subset of patients. Additionally, autologous fat transplant is widely used and accepted in clinical practice as a breast reconstruction procedure (Hayes, 2020; De Decker, et al., 2016; Claro, et al., 2012; Parikh, et al., 2012; Saint-Cyr, et al., 2012; Rosing, et al., 2011; de Blacam, et al., 2011; Losken, et al., 2011; Petit, et al., 2011a; Petit, et al., 2011b; Illouz, et al., 2009; Hyakusoku, et al., 2009; Kanchwala, et al., 2009; Chan, et al., 2008; American Society for Dermatologic Surgery (ASDS); 2008; Coleman, et al., 2007; Spear, et al., 2005). Research is ongoing to distinguish benign from malignant lesions after fat grafting (Parikh, et al, 2012).

Professional Societies/Organizations: The 2008 American Society for Dermatologic Surgery (ASDS) guidelines of care on injectable fillers states that, "One significant concern is the safety of fat transfer into the female breast. Calcifications and nodularity may develop and require the patient to undergo numerous tests and repeated evaluations to rule out an underlying malignancy" (Alam, et al., 2008).

Use Outside the US: In 2012 The National Institute of Excellence (NICE) (United Kingdom).published an interventional procedure guidance and overview of breast reconstruction using lipomodelling after breast cancer treatment. The guidance states that the “current evidence on the efficacy of breast reconstruction using lipomodelling after breast cancer treatment is adequate and the evidence raises no major safety concerns. Therefore this procedure may be used provided that normal arrangements are in place for clinical governance, consent and audit. There is a theoretical concern about any possible influence of the procedure on recurrence of breast cancer in the long-term, although there is no evidence of this in published reports. NICE therefore encourages long-term data collection on this procedure. Patient selection should be carried out by a breast cancer multidisciplinary team. Breast reconstruction using lipomodelling after breast cancer treatment should only be carried out by surgeons with specialist expertise and training in the procedure” (NICE, 2012).

The Australian Safety and Efficacy Register of New Interventional Procedures –Surgical (ASERNIP-S) published a systematic review on autologous fat transfer for cosmetic and reconstructive breast augmentation. The authors concluded that “the evidence base in this review is rated as poor, limited by the quality of the available evidence. Specific limitations of the evidence include absence of studies comparing autologous fat transfer to the nominated comparator procedures, as well as a lack of standardized reporting of outcomes. Autologous fat transfer for cosmetic and reconstructive breast augmentation is considered to be at least as safe as the nominated comparator procedures. It is important to note that this rating is based on indirect comparisons that have been made using overall complication rates. Important safety data examining the effect of microcalcifications following autologous fat transfer on subsequent breast cancer detection were not reported in the studies included in this review; therefore, safety in regards to this outcome cannot be determined. The efficacy of autologous fat transfer cannot be determined from the literature included in this review. Efficacy outcomes reported in the included autologous fat transfer studies varied from those reported for the nominated comparator procedures; therefore, it was not possible to compare efficacy. However, the inability of autologous fat transfer to achieve a volume increase comparable to that of prostheses or autologous tissue augmentation suggests that it is less efficacious than these comparator procedures. There is a need for controlled trials (ideally randomized), assessing the effects of microcalcifications following autologous fat transfer on immediate and long-term breast cancer detection, to be conducted. Studies to determine the maximal breast volume increase reliably achieved by autologous fat transfer would also be useful in order to define the patient population who would benefit most from the procedure, as well as which breast indications should be treated using autologous fat transfer” (Leopardi, et al., 2010).

Autologous Fat Transplant with the use of Adipose-Derived Stem Cells

Research has indicated that subcutaneous fat contains many stem and regenerative cells including cells important in tissue survival and vascularization. It is proposed that autologous adipose-derived regenerative cells (ADRCs) may increase graft survival. With the growing use of autologous fat grafting, published preclinical and clinical data describing cell enriched adipose tissue transplants in breast defect repairs and breast augmentation are increasing. After adipose harvesting using syringe liposuction, the tissue is processed with a system such as the Celution 800 System®, (Cytori Therapeutics, Inc., San Diego, CA) which washes the graft and isolates ADRCs (Kamakura, et al., 2011).

Literature Review: The available literature is limited and consists mostly of case reports and case series, both cosmetic and reconstructive (Ito, et al., 2017; Pérez-Cano, et al., 2012; Kamakura, et al., 2011). Optimal patient selection criteria have not been established through well-designed comparative clinical trials with long-term outcomes data (Kamakura, et al., 2011; Yoshimura, et al., 2008).

Pérez-Cano et al. (2012) conducted a single-arm, prospective, multicenter clinical trial of 71 women who underwent breast conserving surgery for breast cancer and autologous adipose-derived regenerative cell (ADRC)-enriched fat grafting for reconstruction of defects =150 mL (the RESTORE-2 trial). Endpoints included patient and investigator satisfaction with functional and cosmetic results and improvement in overall breast deformity at 12 months post-procedure. Females presenting with partial mastectomy defects and without breast prosthesis were eligible. The RESTORE-2 protocol allowed for up to two treatment sessions and 24 patients elected to undergo a second procedure following the six-month follow-up visit. Of the 67 patients treated, 50 reported satisfaction with treatment results through 12 months. Sixty-one patients underwent radiation therapy as part of their treatment; two patients did not receive radiation and the status of radiation treatment was not known

for the other four patients. Using the same metric, investigators reported satisfaction with 57 out of 67 patients. There were no serious adverse events associated with the ADRC-enriched fat graft injection procedure. There were no reported local cancer recurrences. The investigators reported improvement from baseline through 12 months in the degree of retraction or atrophy in 29 out of 67 patients, while 34 patients had no change and four patients reported worse symptoms. Post-radiation fibrosis at 12 months was reported as improved in 29 patients, while 35 patients had no change and three patients had worse symptoms. Management of atrophy was reported as improved in 17 patients, with 48 patients having no change and two patients reporting worse symptoms. The authors reported that future comparative studies are needed to determine the benefit of ADRC-enriched fat grafting as compared to traditional fat grafting in various clinical circumstances.

In a case series study, Kamakura et al. (2011), reported on the use of autologous adipose-derived stem cell (ADSC) enriched fat grafting for breast augmentation (n=20) After the adipose tissue was harvested by liposuction, it was processed in the Celution 800 System® to wash and isolate the adipose-derived regenerative cells and produce a fat graft enriched with the regenerative cells. The average number of cells per gram of harvested adipose tissue was 3.4×10^5 , and the mean cell viability as measured with an automated cell counting system before graft delivery was 85%. Clinical outcomes measured included improvement in circumferential breast measurement from baseline state. There was improvement in circumferential breast measurement in all patients, and breast measurements were stable by three months after grafting. At nine months, the mean breast measurement had increased 3.3cm from preoperative measurements. Through nine months, overall patient satisfaction was 75%, and physician satisfaction 69%. The procedure was well-tolerated without any serious adverse events. Postoperative cyst formation was seen in two patients. This study was limited by small sample size, no control group and lack of long-term outcomes. The authors reported that to date a, a randomized, controlled study has not been performed to compare the outcomes of cell-enriched fat transfer with those of traditional fat transfer.

Suction-Assisted Lipectomy

Suction-assisted lipectomy of the trunk or extremity area is a procedure in which excess fat deposits are removed using a liposuction cannula with the goal of recontouring the body, thereby improving appearance. This procedure may be performed alone or as one component of the flap breast reconstruction procedure. Suction-assisted lipectomy, when performed alone and not as part of a medically necessary flap breast reconstruction procedure is considered cosmetic in nature. When the procedure is performed as part of a medically necessary flap breast reconstruction procedure, suction-assisted lipectomy of the trunk or extremity area is considered incidental to the primary procedure.

External Breast Prostheses and Mastectomy Bras Following Mastectomy or Lumpectomy

Breast reconstruction has become an integral component of the treatment of patients with breast cancer who have undergone a mastectomy or lumpectomy. External breast prostheses are available for women who have uneven- or unequal-sized breasts and who decide not to, or are waiting to, undergo surgical breast reconstruction. They may choose to wear a breast prosthesis and mastectomy bra, or elect to wear a mastectomy garment that has the prosthesis already inserted in it.

Prostheses can attach to the skin with a fabric backing and adhesive or may be worn unattached with a mastectomy bra. Prefabricated prostheses come in various shapes, sizes and skin tones. Custom-fabricated prostheses are custom-designed and special ordered for the individual. These are usually patterned after a mold that is taken of the breast and chest wall prior to surgery. In general, prefabricated prostheses can adequately meet the external prosthetic needs of most individuals. Therefore, custom-fabricated prosthetic garments would not generally be considered medically necessary. Reusable external nipple prostheses are available to cover flat or missing nipples.

U.S. Food and Drug Administration (FDA)

The FDA classifies external aesthetic restoration prostheses as Class I devices that are exempt from premarket notification (FDA, 2013).

Federal Mandate

The Women's Health and Cancer Rights Act of 1998 (WHCRA) was enacted as a federal mandate in October 1998. The federal mandate defines coverage for breast reconstruction following mastectomy as:

- reconstruction of the breast on which the mastectomy was performed
- surgery and reconstruction on the other breast to produce symmetrical appearance
- prostheses and treatment of physical complications in all stages of mastectomy, including lymphedemas

Under this mandate, benefits for breast reconstruction services following mastectomy or lumpectomy must be provided to both men and women; a diagnosis of breast cancer cannot be required; and timing of breast reconstruction services is not a factor in coverage. In addition, the mandate prohibits any limitations to the number of prostheses or the length of time from the date of the mastectomy.

Appendix A Product/CPT/HCPCS Code Crosswalk

Product	CPT® Code	HCPCS Code
Considered Medically Necessary		
AlloDerm®	15777	Q4116
AlloMax™	15777	C1781, Q4100
Cortiva™	15777	Q4100, C9399
Dermacell™	15777	Q4122
FlexHD®	15777	Q4128
NeoForm™ Dermis	15777	C1781, Q4100
Experimental, Investigational or Unproven and Not Medically Necessary		
ARTIA™ Reconstructive Tissue Matrix	15777	C1763
BellaDerm® Acellular Hydrated Dermis	15777	C1781, Q4100
Biodesign® Nipple Reconstruction Cylinder	19350	C1763
DermaMatrix Acellular Dermis	15777	C1781, Q4100
GalaFLEX® Surgical Scaffold/Mesh	15777	C1781, Q4100
GalaFORM™ 3D	15777	C1781, Q4100
GalaSHAPE 3D	15777	C1781, Q4100
hMatrix	15777	Q4134
Juvederm®	19350, 11950	C9399
Permacol™	15777	C9364
Phasix™ Mesh	15777	C1781
Repriza®	15777	Q4143
SERI™ Surgical Scaffold	15777	C1781, Q4100
Strattice™ Reconstructive Tissue Matrix	15777	Q4130
SurgiMend® Collagen Matrix	15777	C9358, C9360
Veritas Collagen Matrix	15777	Q4100, C9354

Medicare Coverage Determinations

	Contractor	Policy Name/Number	Revision Effective Date
NCD	National	Breast Reconstruction Following Mastectomy (140.2)	1/1/1997
LCD	CGS	External Breast Prostheses (L33317)	1/01/2020

Note: Please review the current Medicare Policy for the most up-to-date information.

Coding/Billing Information

- Note:** 1) This list of codes may not be all-inclusive.
 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Breast Reconstruction

Not covered when used to report intraoperative assessment of tissue perfusion as it considered integral to the primary procedure and not separately reimbursable.

CPT®* Codes	Description
15860	Intravenous injection of agent (eg fluorescein) to test vascular flow in flap or graft

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
11920	Tattooing, intradermal introduction of insoluble opaque pigments to correct color defects of skin, including micropigmentation; 6.0 sq cm or less
11921	Tattooing, intradermal introduction of insoluble opaque pigments to correct color defects of skin, including micropigmentation; 6.1 to 20.0 sq cm
11970	Replacement of tissue expander with permanent implant
11971	Removal of tissue expander without insertion of implant
13100	Repair, complex, trunk; 1.1 cm to 2.5 cm
13101	Repair, complex, trunk; 2.6 cm to 7.5 cm
13102	Repair, complex, trunk; each additional 5 cm or less (List separately in addition to code for primary procedure)
15734	Muscle, myocutaneous, or fasciocutaneous flap; trunk
15771†	Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; 50 cc or less injectate
15772†	Grafting of autologous fat harvested by liposuction technique to trunk, breasts, scalp, arms, and/or legs; each additional 50 cc injectate, or part thereof (List separately in addition to code for primary procedure)
15777	Implantation of biologic implant (eg, acellular dermal matrix) for soft tissue reinforcement (eg, breast, trunk) (List separately in addition to code for primary procedure)
15839	Excision, excessive skin and subcutaneous tissue (includes lipectomy); other area
19316	Mastopexy
19318	Breast reduction
19324	Mammoplasty, augmentation; without prosthetic implant (Code deleted 12/31/2020)
19325	Breast augmentation with implant
19328	Removal of intact breast implant
19330	Removal of ruptured breast implant, including implant contents (eg, saline, silicone gel)
19340	Insertion of breast implant on same day of mastectomy (ie, immediate)
19342	Insertion or replacement of breast implant on separate day from mastectomy
19350††	Nipple/areola reconstruction
19357	Tissue expander placement in breast reconstruction, including subsequent expansion
19361	Breast reconstruction; with latissimus dorsi flap
19364	Breast reconstruction; with free flap (eg, fTRAM, DIEP, SIEA, GAP flap)
19366	Breast reconstruction with other technique (Code deleted 12/31/2020)
19367	Breast reconstruction; with single-pedicled transverse rectus abdominis myocutaneous (TRAM) flap
19368	Breast reconstruction; with single-pedicled transverse rectus abdominis myocutaneous (TRAM) flap, requiring separate microvascular anastomosis (supercharging)
19369	Breast reconstruction; with bipedicled transverse rectus abdominis myocutaneous (TRAM) flap
19370	Revision of peri-implant capsule, breast, including capsulotomy, capsulorrhaphy, and/or partial capsulectomy
19371	Peri-implant capsulectomy, breast, complete, including removal of all intracapsular contents

CPT®* Codes	Description
19380	Revision of reconstructed breast (eg, significant removal of tissue, re-advancement and/or re-inset of flaps in autologous reconstruction or significant capsular revision combined with soft tissue excision in implant-based reconstruction)
19499†††	Unlisted procedure, breast

†**Note:** Considered medically necessary when used to report autologous fat transplant without the use of adipose-derived stem cells or xenograft cartilage grafting.

††**Note:** Considered experimental/investigational/unproven when used to report nipple reconstruction with Juvederm, Radiesse, or Cook Biodesign® Nipple Reconstruction Cylinder.

†††**Note:** Considered medically necessary when used to report thoracodorsal artery perforator (TDAP) flap with a breast reconstruction procedure performed on the diseased/affected breast (i.e., breast on which the mastectomy/lumpectomy was performed).

HCPCS Codes	Description
C1789	Prosthesis, breast (implantable)
L8600	Implantable breast prosthesis, silicone or equal
Q4116	Alloderm, per square centimeter
Q4122	Dermacell, per square centimeter
Q4128	Flex hd or allopatch hd, or matrix hd per square centimeter
S2066	Breast reconstruction with gluteal artery perforator (GAP) flap, including harvesting of the flap, microvascular transfer, closure of donor site and shaping the flap into a breast, unilateral
S2067	Breast reconstruction of a single breast with "stacked" deep inferior epigastric perforator (DIEP) flap(s) and/or gluteal artery perforator (GAP) flap(s), including harvesting of the flap(s), microvascular transfer, closure of donor site(s) and shaping the flap into a breast, unilateral
S2068	Breast reconstruction with deep inferior epigastric perforator (DIEP) flap or superficial inferior epigastric artery (SIEA) flap, including harvesting of the flap, microvascular transfer, closure of donor site and shaping the flap into a breast, unilateral

Considered medically necessary when used to report AlloMax™ or NeoForm™ Dermis used in association with a medically necessary breast reconstruction procedure:

CPT®* Codes	Description
C1781	Mesh (implantable)
Q4100	Skin substitute, not otherwise specified

Experimental/Investigational/Unproven when used to report a product as not covered as listed in the policy statement above:

CPT®* Codes	Description
11950††††	Subcutaneous injection of filling material (eg, collagen); 1 cc or less
11951††††	Subcutaneous injection of filling material (eg, collagen); 1.1 to 5.0 cc
11952††††	Subcutaneous injection of filling material (eg, collagen); 5.1 to 10.0 cc
11954††††	Subcutaneous injection of filling material (eg, collagen); over 10.0 cc

††††**Note:** Experimental/Investigational/Unproven when used to report autologous fat transplant with the use of adipose-derived stem cells or xenograft cartilage grafting.

HCPCS Codes	Description
C9358	Dermal substitute, native, non-denatured collagen, fetal bovine origin, (Surgimend collagen matrix), per 0.5 square centimeters
C9360	Dermal substitute, native, non-denatured collagen, neonatal bovine origin (Surgimend collagen matrix), per 0.5 square centimeters
C9364	Porcine implant Permacol, per square cm
Q2026	Injection, Radiesse, 0.1 ml
Q4134	Hmatrix, per square centimeter
Q4143	Repriza, per square centimeter

Experimental/Investigational/Unproven when used to report Biodesign® Nipple Reconstruction Cylinder:

HCPCS Codes	Description
C1763	Connective tissue, non-human (includes synthetic)

Experimental/Investigational/Unproven when used to report ARTIA™ Reconstructive Tissue Matrix, BellaDerm® Acellular Hydrated Dermis, DermaMatrix Acellular Dermis, GalaFLEX® Surgical Scaffold, GalaFORM™ 3D, Juvederm®, Phasix™ Mesh, or SERI™ Surgical Scaffold:

HCPCS Codes	Description
C1781	Mesh (implantable)
C9399	Unclassified drugs or biologicals
Q4100	Skin substitute, not otherwise specified

Experimental/Investigational/Unproven when used to report Strattice™ :

HCPCS Codes	Description
Q4130	Strattice™, per square centimeter

Experimental/Investigational/Unproven when used to report Veritas Collagen Matrix:

HCPCS Codes	Description
C9354	Acellular pericardial tissue matrix of non-human origin (Veritas), per square centimeter
Q4100	Skin substitute, not otherwise specified

Experimental/Investigational/Unproven when used to report breast reconstruction procedures using adipose-derived stem cells in autologous fat transplantation or xenograft cartilage grafting:

CPT®*	Description
19366	Breast reconstruction with other technique
19499	Unlisted procedure, breast

Cosmetic in nature/Not Medically Necessary when used to report correction of surgically-induced donor site asymmetry that results from flap breast reconstruction procedures:

CPT®*	Description
15877††††	Suction assisted lipectomy; trunk
15879††††	Suction assisted lipectomy; lower extremity

++++**Note:** Considered incidental to the primary procedure when used to report suction-assisted lipectomy of the trunk as part of a medically necessary flap breast reconstruction procedure

External Breast Prostheses and Mastectomy Bras Following Mastectomy or Lumpectomy

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

HCPCS Codes	Description
L8000	Breast prosthesis; mastectomy bra, without integrated breast prosthesis form, any size, any type
L8001	Breast prosthesis, mastectomy bra, with integrated breast prosthesis form, unilateral, any size, any type
L8002	Breast prosthesis, mastectomy bra, with integrated breast prosthesis form, bilateral, any size, any type
L8015	External breast prosthesis garment, with mastectomy form, post mastectomy
L8020	Breast prosthesis, mastectomy form
L8030	Breast prosthesis, silicone or equal, without integral adhesive
L8031	Breast prosthesis, silicone or equal, with integral adhesive
L8032	Nipple prosthesis, prefabricated, reusable, any type, each
L8033	Nipple prosthesis, custom fabricated, reusable, any material, any type, each
L8035	Custom breast prosthesis, post mastectomy, molded to patient model
L8039	Breast prosthesis, not otherwise specified

***Current Procedural Terminology (CPT®) ©2020 American Medical Association: Chicago, IL.**

References

1. Adams WP Jr, Toriumi DM, Van Natta BW. Clinical Use of GalaFLEX in Facial and Breast Cosmetic Plastic Surgery. *Aesthet Surg J.* 2016 Nov;36(suppl 2):S23-S32.
2. Alam M, Gladstone H, Kramer EM, Murphy JP Jr, Nouri K, Neuhaus IM, et al; American Society for Dermatologic Surgery. ASDS guidelines of care: injectable fillers. *Dermatol Surg.* 2008 Jun;34 Suppl 1:S115-48.
3. Alderman AK, Kuzon WM Jr, Wilkins EG. A two-year prospective analysis of trunk function in TRAM breast reconstructions. *Plast Reconstr Surg.* 2006 Jun;117(7):2131-8.
4. Allergan. Juvederm®. Accessed Jan 25, 2021. Available at URL address: <https://www.juvederm.com/>
5. Allergan. Our products. 2020. Accessed Jan 25, 2021. Available at URL address: <https://www.allergan.com/products>
6. American Association of Tissue Banks (AATB). 2019. Accessed Jan 25, 2021. Available at URL address: <https://www.aatb.org/>
7. American Cancer Society (ACS): Breast reconstruction surgery. Last revised Sept 18, 2019. Accessed Jan 25, 2021. Available at URL address: <http://www.cancer.org>
8. American Society of Plastic Surgeons (ASPS). Evidence-Based Clinical Practice Guideline: Autologous Breast Reconstruction with DIEP or Pedicled TRAM Abdominal Flaps. 2017. Accessed Jan 27, 2021. Available at URL address: <http://www.plasticsurgery.org/for-medical-professionals/legislation-and-advocacy/health-policy-resources/evidence-based-guidelinespractice-parameters.html>

9. Antony AK, McCarthy CM, Cordeiro PG, Mehrara BJ, Pusic AL, Teo EH, et al. Acellular human dermis implantation in 153 immediate two-stage tissue expander breast reconstructions: determining the incidence and significant predictors of complications. *Plast Reconstr Surg.* 2010 Jun;125(6):1606-14.
10. Bacterin International Holdings Inc., Belgrade, MT. hMatrix®. 2015. Accessed Jan 27, 2021. Available at URL address: http://xtantmedical.com/app/uploads/2016/09/hMatrix-PR-ADM-Brochure_5258D_Digital.pdf
11. Bajaj AK, Chevray PM, Chang DW. Comparison of donor-site complications and functional outcomes in free muscle-sparing TRAM flap and free DIEP flap breast reconstruction. *Plast Reconstr Surg.* 2006 Mar;117(3):737-46; discussion 747-50.
12. Baker BG, Irri R, MacCallum V, Chattopadhyay R, Murphy J, Harvey JR. A Prospective Comparison of Short-Term Outcomes of Subpectoral and Prepectoral Strattice-Based Immediate Breast Reconstruction. *Plast Reconstr Surg.* 2018 May;141(5):1077-1084.
13. Ball JF, Sheena Y, Tarek Saleh DM, Forouhi P, Benyon SL, Irwin MS, Malata CM. A direct comparison of porcine (Strattice™) and bovine (Surgimend™) acellular dermal matrices in implant-based immediate breast reconstruction. *J Plast Reconstr Aesthet Surg.* 2017 Aug;70(8):1076-1082.
14. Barber MD, Williams L, Anderson ED, Neades GT, Raine C, Young O, et al. Outcome of the use of acellular-dermal matrix to assist implant-based breast reconstruction in a single centre. *Eur J Surg Oncol.* 2015 Jan;41(1):100-5.
15. Bard Davol, Inc. AlloMax™ surgical graft for breast reconstruction. Accessed Jan 27, 2021. Available at URL address: <https://www.crbard.com/Davol/en-US/Ordering-Info>
16. Bard Davol, Inc. Phasix™ Mesh. Accessed Jan 27, 2021. Available at URL address: <https://www.bd.com/en-us/offerings/capabilities/hernia-repair-and-fixation/hernia-repair-mesh/bioresorbable-mesh>
17. Baxter RA. Intracapsular allogenic dermal grafts for breast implant-related problems. *Plast Reconstr Surg.* 2003 Nov;112(6):1692-6; discussion 1697-8.
18. Beckenstein MS. Nipple Reconstruction. In: Grabb and Smith Plastic Surgery. 7th ed. Lippincott, Williams and Wilkins. Philadelphia, PA; 2014. Ch 63.
19. Becker S, Saint-Cyr M, Wong C, Dauwe P, Nagarkar P, Thornton JF, Peng Y. AlloDerm versus DermaMatrix in immediate expander-based breast reconstruction: a preliminary comparison of complication profiles and material compliance. *Plast Reconstr Surg.* 2009 Jan;123(1):1-6; discussion 107-8.
20. Behnam AB, Nguyen D, Moran SL, Serletti JM. TRAM flap breast reconstruction for patients with advanced breast disease. *Ann Plast Surg.* 2003 Jun;50(6):567-71.
21. Bindingavele V, Gaon M, Ota KS, Kulber DA, Lee DJ. Use of acellular cadaveric dermis and tissue expansion in postmastectomy breast reconstruction. *J Plast Reconstr Aesthet Surg.* 2007;60(11):1214-8.
22. Borgognone A, Anniboletti T, De Vita F. Does Veritas® play a role in breast reconstruction? a case report. *Breast Cancer (Dove Med Press).* 2011 Dec 20;3:175-7.
23. Bregnhøj A, Haedersdal M. Q-switched YAG laser vs. punch biopsy excision for iatrogenic radiation tattoo markers--a randomized controlled trial. *J Eur Acad Dermatol Venereol.* 2010 Oct;24(10):1183-6.
24. Breuing KH, Colwell AS. Inferolateral AlloDerm hammock for implant coverage in breast reconstruction. *Ann Plast Surg.* 2007 Sep;59(3):250-5.

25. Breuing KH, Warren SM. Immediate bilateral breast reconstruction with implants and inferolateral AlloDerm slings. *Ann Plast Surg.* 2005 Sep;55(3):232-9.
26. Brooke S, Mesa J, Uluer M, Michelotti B, Moyer K, Neves RI, et al. Complications in Tissue Expander Breast Reconstruction: A Comparison of AlloDerm, DermaMatrix, and FlexHD Acellular Inferior Pole Dermal Slings. *Ann Plast Surg.* 2012 Aug 3.
27. Bullocks JM. DermACELL: a novel and biocompatible acellular dermal matrix in tissue expander and implant-based breast reconstruction. *Eur J Plast Surg.* 2014;37(10):529-538.
28. Butterfield JL. 440 Consecutive immediate, implant-based, single-surgeon breast reconstructions in 281 patients: a comparison of early outcomes and costs between SurgiMend fetal bovine and AlloDerm human cadaveric acellular dermal matrices. *Plast Reconstr Surg.* 2013 May;131(5):940-51.
29. Cahan AC, Palaia DA, Rosenberg M, Bonanno PC. The aesthetic mastectomy utilizing a non-nipple-sparing portal approach. *Ann Plast Surg.* 2011 May;66(5):424-8.
30. Carlson GW, Styblo TM, Lyles RH, Jones G, Murray DR, Staley CA, Wood WC. The use of skin sparing mastectomy in the treatment of breast cancer: The Emory experience. *Surgical Oncology.* 2003 Dec;12(4):265-9.
31. Caviglioli F, Maione L, Forcellini D, Klinger F, Klinger M. Autologous fat graft in postmastectomy pain syndrome. *Plast Reconstr Surg.* 2011 Aug;128(2):349-52
32. Centers for Medicare and Medicaid Services (CMS). Healthcare Common Procedure Coding System (HCPCS). May 2015. Accessed Jan 27, 2021. Available at URL address: <https://www.cms.gov/Medicare/Coding/MedHCPCSGenInfo/Downloads/May-7-2015-DrugAgenda.pdf>
33. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed Jan 27, 2021. Available at URL address: https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx?Cntrctr=373&ContrVer=1&CntrctrSelected=373*1&DocType=Active%7cFuture&s=All&bc=AggAAAQAAAA&
34. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for breast reconstruction following mastectomy (140.2), Jan 1,1997. Accessed Jan 27, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/ncd-alphabetical-index.aspx>
35. Centers for Medicare & Medicaid Services (CMS). The Women's Health and Cancer Rights Act. Accessed Jan 27, 2021. Available at URL address: http://www.cms.gov/CCIIO/Programs-and-Initiatives/Other-Insurance-Protections/whcra_factsheet.html
36. Chan CW, McCulley SJ, Macmillan RD. Autologous fat transfer--a review of the literature with a focus on breast cancer surgery. *J Plast Reconstr Aesthet Surg.* 2008 Dec;61(12):1438-48.
37. Chang EI, Liu J. Prospective unbiased experience with three acellular dermal matrices in breast reconstruction. *J Surg Oncol.* 2017 Sep;116(3):365-370.
38. Cheng A, Saint-Cyr M. Comparison of different ADM materials in breast surgery. *Clin Plast Surg.* 2012 Apr;39(2):167-75.
39. Cheng MH, Rodriguez ED, Smartt JM, Cardenas-Mejia A. Nipple reconstruction using the modified top hat flap with banked costal cartilage graft: long-term follow-up in 58 patients. *Ann Plast Surg.* 2007 Dec;59(6):621-8.

40. Chun YS. Nipple-Areola Reconstruction. Updated Jan 31, 2019. Accessed Jan 27, 2021. Available at URL address: <http://www.emedicine.com/plastic/topic144.htm>
41. Claro F Jr, Figueiredo JC, Zampar AG, Pinto-Neto AM. Applicability and safety of autologous fat for reconstruction of the breast. *Br J Surg*. 2012 Jun;99(6):768-80.
42. ClinicalTrials.org. Compare Outcomes Between Two Acellular Dermal Matrices. NCT02891759. First posted: Sept 7, 2016. Last updated: May 13, 2020. Accessed Jan 27, 2021. Available at URL address: <https://clinicaltrials.gov/show/NCT02891759>
43. Coleman SR, Saboeiro AP. Fat grafting to the breast revisited: safety and efficacy. *Plast Reconstr Surg*. 2007 Mar;119(3):775-85; discussion 786-7.
44. Collins B, Williams JZ, Karu H, Hodde JP, Martin VA, Gurtner GC. Nipple Reconstruction with the Biodesign Nipple Reconstruction Cylinder: A Prospective Clinical Study. *Plast Reconstr Surg Glob Open*. 2016 Aug 9;4(8):e832.
45. Colwell AS, Damjanovic B, Zahedi B, Medford-Davis L, Hertl C, Austen WG Jr. Retrospective review of 331 consecutive immediate single-stage implant reconstructions with acellular dermal matrix: indications, complications, trends, and costs. *Plast Reconstr Surg*. 2011 Dec;128(6):1170-8.
46. Craft RO, May JW Jr. Staged nipple reconstruction with vascularized SurgiMend acellular dermal matrix. *Plast Reconstr Surg*. 2011 Jun;127(6):148e-9e.
47. de Blacam C, Momoh AO, Colakoglu S, Tobias AM, Lee BT. Evaluation of clinical outcomes and aesthetic results after autologous fat grafting for contour deformities of the reconstructed breast. *Plast Reconstr Surg*. 2011 Nov;128(5):411e-418e.
48. De Decker M, De Schrijver L, Thiessen F, Tondu T, Van Goethem M, Tjalma WA. Breast cancer and fat grafting: efficacy, safety and complications-a systematic review. *Eur J Obstet Gynecol Reprod Biol*. 2016 Dec;207:100-108.
49. DellaCroce FJ. Perforator Flap Reconstruction Treatment and Management. Updated Feb 28, 2019. Accessed Jan 27, 2021. Available at URL address: <http://emedicine.medscape.com/article/1276406-treatment>
50. Deutsch MF, Robb GL, Talavera F, Shenaq SM, Slenkovich N, Aly A. Free Flap Breast Reconstruction. Updated Jan 7, 2021. Accessed Jan 27, 2021. Available at URL address: <http://www.emedicine.com/plastic/topic140.htm>
51. Dikmans RE, El Morabit F, Ottenhof MJ, Tuinder SM, Twisk JW, Moues C, et al. Single-stage breast reconstruction using Strattice™: A retrospective study. *J Plast Reconstr Aesthet Surg*. 2016 Feb;69(2):227-33.
52. Dikmans RE, Negenborn VL, Bouman MB, Winters HA, Twisk JW, Ruhé PQ, et al. Two-stage implant-based breast reconstruction compared with immediate one-stage implant-based breast reconstruction augmented with an acellular dermal matrix: an open-label, phase 4, multicentre, randomised, controlled trial. *Lancet Oncol*. 2017 Feb;18(2):251-258.
53. Draper LB, Bui DT, Chiu ES, Mehrara BJ, Pusic AL, Cordeiro PG, Disa JJ. Nipple-Areola Reconstruction Following Chest-Wall Irradiation for Breast Cancer: Is It Safe? *Ann Plast Surg*. 2005 Jul;55(1):12-5.
54. Evans KK, Rasko Y, Lenert J, Olding M. The use of calcium hydroxylapatite for nipple projection after failed nipple-areolar reconstruction: early results. *Ann Plast Surg*. 2005 Jul;55(1):25-9; discussion 29.

55. Fansa H, Schirmer S, Warnecke IC, Cervelli A, Frerichs O. The transverse myocutaneous gracilis muscle flap: a fast and reliable method for breast reconstruction. *Plast Reconstr Surg.* 2008 Nov;122(5):1326-33.
56. Fine NA, Lehfeldt M, Gross JE, Downey S, Kind GM, Duda G, et al. SERI surgical scaffold, prospective clinical trial of a silk-derived biological scaffold in two-stage breast reconstruction: 1-year data. *Plast Reconstr Surg.* 2015 Feb;135(2):339-51.
57. Gabbay JS, Eby JB, Kulber DA. The midabdominal TRAM flap for breast reconstruction in morbidly obese patients. *Plast Reconstr Surg.* 2005 Mar;115(3):764-70.
58. Galatea Surgical. GalaShape® 3D Surgical Scaffold. 2021. Accessed Jan 27, 2021. Available at URL address: <https://www.galateasurgical.com/surgical-scaffolds/now-available-3d/>
59. Gamboa-Bobadilla GM. Implant breast reconstruction using acellular dermal matrix. *Ann Plast Surg.* 2006 Jan;56(1):22-5.
60. Garramone CE, Lam B. Use of AlloDerm in primary nipple reconstruction to improve long-term nipple projection. *Plast Reconstr Surg.* 2007 May;119(6):1663-8.
61. Gaster RS, Berger AJ, Monica SD, Sweeney RT, Endress R, Lee GK. Histologic Analysis of Fetal Bovine Derived Acellular Dermal Matrix in Tissue Expander Breast Reconstruction. *Ann Plast Surg.* 2013 Mar 11.
62. Garvey PB, Buchel EW, Pockaj BA, Casey WJ 3rd, Gray RJ, Hernandez JL, Samson TD. DIEP and pedicled TRAM flaps: a comparison of outcomes. *Plast Reconstr Surg.* 2006 May;117(6):1711-9; discussion 1720-1.
63. Glasberg SB, D'Amico RA. Use of regenerative human acellular tissue (AlloDerm) to reconstruct the abdominal wall following pedicle TRAM flap breast reconstruction surgery. *Plast Reconstr Surg.* 2006 Jul;118(1):8-15.
64. Glasberg SB, Light D. AlloDerm and Stratice in breast reconstruction: a comparison and techniques for optimizing outcomes. *Plast Reconstr Surg.* 2012 Jun;129(6):1223-33.
65. Guerra AB, Khoobehi K, Metzinger SE, Allen RJ. New technique for nipple areola reconstruction: arrow flap and rib cartilage graft for long-lasting nipple projection. *Ann Plast Surg.* 2003 Jan;50(1):31-7.
66. Gurtner GC, Jones GE, Neligan PC, Newman MI, Phillips BT, Sacks JM, Zenn MR. Intraoperative laser angiography using the SPY system: review of the literature and recommendations for use. *Ann Surg Innov Res.* 2013 Jan 7;7(1):1.
67. Gutowski KA; ASPS Fat Graft Task Force. Current applications and safety of autologous fat grafts: a report of the ASPS fat graft task force. *Plast Reconstr Surg.* 2009 Jul;124(1):272-80.
68. Haddock N, Levine J. Breast reconstruction with implants, tissue expanders and AlloDerm: predicting volume and maximizing the skin envelope in skin sparing mastectomies. *Breast J.* 2010 Jan-Feb;16(1):14-9.
69. Hayes, Inc. Hayes Medical Technology Directory Report. Human Acellular Dermal Matrix Grafts for Breast Reconstruction. Lansdale, PA: Hayes, Inc.; published Apr 2014; annual review Apr 2015; Apr 2016; Apr 2017; Oct 2018; archived Jan 28, 2019.
70. Hayes, Inc. Medical Technology Directory. Comparative effectiveness review of human acellular dermal matrix for breast reconstruction. Lansdale, PA: Hayes, Inc.; published Jan 28, 2019; annual review May 11, 2020.

71. Hayes, Inc. Hayes Technology Brief. Superficial Inferior Epigastric Artery (SIEA) Flap Procedure for Postmastectomy Breast Reconstruction. Lansdale, PA: Hayes, Inc.; Nov 2014; annual review Oct 2015; Oct 2016; archived Dec 19, 2017.
72. Hayes, Inc. Hayes Technology Brief. Autologous Fat Grafting for Breast Reconstruction After Breast Cancer Surgery. Lansdale, PA: Hayes, Inc.; Aug 27, 2015; annual review Jul 2016; Jul 2017; archived Sept 26, 2018.
73. Hayes, Inc. Health Technology Assessment. Autologous Fat Grafting for Breast Reconstruction After Breast Cancer Surgery. Lansdale, PA: Hayes, Inc.; Oct 21, 2020.
74. Hayes, Inc. Hayes Technology Brief. SurgiMend (Integra LifeSciences) for Postmastectomy Breast Reconstruction. Lansdale, PA: Hayes, Inc.; Apr 27, 2017; review May 2, 2019; archived May 26, 2020.
75. Hayes, Inc. Clinical Research Response. Product Comparison. Silicone Gel Breast Implants. Lansdale, PA: Hayes, Inc.; Jan 26, 2017; archived Apr 17, 2018.
76. Hayes, Inc. Clinical Research Response. Product Comparison. Biological Versus Resorbable Mesh for Breast Reconstruction. Lansdale, PA: Hayes, Inc.; May 4, 2017; archived Aug 5, 2018.
77. Hayes, Inc. Clinical Research Response. GalaFLEX Surgical Scaffold (Galatea Surgical). Lansdale, PA: Hayes, Inc.; Nov 20, 2017; archived Dec 19, 2018.
78. Heitland A, Markowicz M, Koellensperger E, Allen R, Pallua N. Long-term nipple shrinkage following augmentation by an autologous rib cartilage transplant in free DIEP-flaps. *J Plast Reconstr Aesthet Surg.* 2006;59(10):1063-7.
79. Ho G, Nguyen TJ, Shahabi A, Hwang BH, Chan LS, Wong AK. A systematic review and meta-analysis of complications associated with acellular dermal matrix-assisted breast reconstruction. *Ann Plast Surg.* 2012 Apr;68(4):346-56.
80. Hu E, Alderman AK. Breast reconstruction. *Surg Clin North Am.* 2007 Apr;87(2):453-67, x.
81. Humphreys K. Autologous fat injection for breast reconstruction. August 2008. Australia and New Zealand Horizon Scanning Network. Accessed Jan 27, 2021. Available at URL address: <http://www.horizonscanning.gov.au/internet/horizon/publishing.nsf/Content/technologies-assessed-lp-2>
82. Hyakusoku H, Ogawa R, Ono S, Ishii N, Hirakawa K. Complications after autologous fat injection to the breast. *Plast Reconstr Surg.* 2009 Jan;123(1):360-70; discussion 371-2.
83. Ibrahim AM, Shuster M, Koolen PG, Kim K, Taghinia AH, Sinno HH, et al. Analysis of the National Surgical Quality Improvement Program database in 19,100 patients undergoing implant-based breast reconstruction: complication rates with acellular dermal matrix. *Plast Reconstr Surg.* 2013 Nov;132(5):1057-66.
84. Illouz YG, Sterodimas A. Autologous fat transplantation to the breast: a personal technique with 25 years of experience. *Aesthetic Plast Surg.* 2009 Sep;33(5):706-15.
85. Integra® LifeSciences Corp. SurgiMend®. Accessed Jan 27, 2021. Available at URL address: <http://www.surgimend.com/>
86. Ito S, Kai Y, Masuda T, Tanaka F, Matsumoto T, Kamohara Y, et al. Long-term outcome of adipose-derived regenerative cell-enriched autologous fat transplantation for reconstruction after breast-conserving surgery for Japanese women with breast cancer. *Surg Today.* 2017 Dec;47(12):1500-1511.

87. Jansen LA, Macadam SA. The use of AlloDerm in postmastectomy alloplastic breast reconstruction: part I. A systematic review. *Plast Reconstr Surg*. 2011 Jun;127(6):2232-44.
88. Jewell M1, Daunch W2, Bengtson B3, Mortarino E2. The development of SERI® Surgical Scaffold, an engineered biological scaffold. *Ann N Y Acad Sci*. 2015 Sep 16.
89. Joanna Nguyen T, Carey JN, Wong AK. Use of human acellular dermal matrix in implant based breast reconstruction: Evaluating the evidence. *J Plast Reconstr Aesthet Surg*. 2011 Mar 7.
90. Kamakura T, Ito K. Autologous cell-enriched fat grafting for breast augmentation. *Aesthetic Plast Surg*. 2011 Dec;35(6):1022-30.
91. Kanchwala SK, Glatt BS, Conant EF, Bucky LP. Autologous fat grafting to the reconstructed breast: the management of acquired contour deformities. *Plast Reconstr Surg*. 2009 Aug;124(2):409-18.
92. Karp N, Choi M, Kulber DA, Downey S, Duda G, Kind GM, et al. SERI Surgical Scaffold in 2-Stage Breast Reconstruction: 2-Year Data from a Prospective, Multicenter Trial. *Plast Reconstr Surg Glob Open*. 2017 May 16;5(5):e1327.
93. Keifer OP Jr, Page EK, Hart A, Rudderman R, Carlson GW, Losken A. A Complication Analysis of 2 Acellular Dermal Matrices in Prosthetic-based Breast Reconstruction. *Plast Reconstr Surg Glob Open*. 2016 Jul 13;4(7):e800.
94. Kilchenmann AJ, Lardi AM, Ho-Asjoe M, Junge K, Farhadi J. An evaluation of resource utilisation of single stage porcine acellular dermal matrix assisted breast reconstruction: A comparative study. *Breast*. 2014 Dec;23(6):876-82
95. Kim EK, Lee TJ. Use of lyophilized allogeneic costal cartilage: is it effective to maintain the projection of the reconstructed nipple? *Ann Plast Surg*. 2011 Feb;66(2):128-30.
96. Kim JY, Davila AA, Persing S, Connor CM, Jovanovic B, Khan SA, et al. A meta-analysis of human acellular dermis and submuscular tissue expander breast reconstruction. *Plast Reconstr Surg*. 2012 Jan;129(1):28-41.
97. Kim J, Long JN. Breast reconstruction with acellular dermis. Updated Jan 10, 2020. Accessed Jan 27, 2021. Available at URL address: <http://emedicine.medscape.com/article/1851090-overview#a1>
98. Kroll SS. Fat necrosis in free transverse rectus abdominis myocutaneous and deep inferior epigastric perforator flaps. *Plast Reconstr Surg*. 2000 Sep;106(3):576–83.
99. Lardi AM, Ho-Asjoe M, Mohanna PN, Farhadi J. Immediate breast reconstruction with acellular dermal matrix: factors affecting outcome. *J Plast Reconstr Aesthet Surg*. 2014 Aug;67(8):1098-105.
100. Lee KT, Mun GH. Updated Evidence of Acellular Dermal Matrix Use for Implant-Based Breast Reconstruction: A Meta-analysis. *Ann Surg Oncol*. 2016 Feb;23(2):600-10.
101. Lee KT, Mun GH. A Meta-analysis of Studies Comparing Outcomes of Diverse Acellular Dermal Matrices for Implant-Based Breast Reconstruction. *Ann Plast Surg*. 2017 Jul;79(1):115-123.
102. Leopardi D, et al. Systematic review of autologous fat transfer for cosmetic and reconstructive breast augmentation. ASERNIP-S Report No.70. Adelaide, South Australia: ASERNIP-S, September 2010. Accessed Dec 27, 2019. Available at URL address: <http://www.surgeons.org>
103. LifeCell Corporation. AlloDerm Tissue Matrix Defined [product information]. Accessed Jan 27, 2021. Available at URL address: <https://hcp.alloderm.com/>

104. LifeCell Corporation. Strattice™. Accessed Jan 27, 2021. Available at URL address: <https://www.allergan.com/products/key-products/product-prescribing/labeling/strattice>
105. LifeNet Health. Wound Management and Surgical Reconstruction. Dermacell® Soft Tissue Reconstruction. 2021. Accessed Jan 25, 2021. Available at URL address: <https://www.lifenethealth.org/wound-management-and-surgical-reconstruction>
106. Liu DZ, Mathes DW, Neligan PC, Said HK, Louie O. Comparison of outcomes using AlloDerm versus FlexHD for implant-based breast reconstruction. *Ann Plast Surg.* 2014 May;72(5):503-7.
107. Losken A, Carlson GW, Bostwick J III, Jones GE, Culbertson JH, Schoemann M. Trends in unilateral breast reconstruction and management of the contralateral breast: the Emory experience. *Plast Reconstr Surg.* 2002 Jul;110(1):89–97.
108. Losken A. Early Results Using Sterilized Acellular Human Dermis (Neoform) in Post-Mastectomy Tissue Expander Breast Reconstruction. *Plast Reconstr Surg.* 2009 Mar 23.
109. Losken A, Pinell XA, Sikoro K, Yezhelyev MV, Anderson E, Carlson GW. Autologous fat grafting in secondary breast reconstruction. *Ann Plast Surg.* 2011 May;66(5):518-22.
110. Martin L, O'Donoghue JM, Horgan K, Thrush S, Johnson R, Gandhi A; Association of Breast Surgery and the British Association of Plastic, Reconstructive and Aesthetic Surgeons. Acellular dermal matrix (ADM) assisted breast reconstruction procedures: joint guidelines from the Association of Breast Surgery and the British Association of Plastic, Reconstructive and Aesthetic Surgeons. *Eur J Surg Oncol.* 2013 May;39(5):425-9.
111. Maxwell GP, Gabriel A. Non-cross-linked porcine acellular dermal matrix in revision breast surgery: long-term outcomes and safety with neopectoral pockets. *Aesthet Surg J.* 2014 May 1;34(4):551-9.
112. Mazari FAK, Wattoo GM, Kazzazi NH, Kolar KM, Olubowale OO, Rogers CE, Azmy IA. The Comparison of Strattice and SurgiMend in Acellular Dermal Matrix-Assisted, Implant-Based Immediate Breast Reconstruction. *Plast Reconstr Surg.* 2018 Feb;141(2):283-293.
113. McCarthy CM, Lee CN, Halvorson EG, Riedel E, Pusic AL, Mehrara BJ, Disa JJ. The use of acellular dermal matrices in two-stage expander/implant reconstruction: a multicenter, blinded, randomized controlled trial. *Plast Reconstr Surg.* 2012 Nov;130(5 Suppl 2):57S-66S.
114. Mehrara BJ, Santoro TD, Arcilla E, Watson JP, Shaw WW, Da Lio AL. Complications after microvascular breast reconstruction: experience with 1195 flaps. *Plast Reconstr Surg.* 2006 Oct;118(5):1100-9; discussion 1110-1.
115. Michelotti BF, Brooke S, Mesa J, Wilson MZ, Moyer K, Mackay DR, et al. Analysis of clinically significant seroma formation in breast reconstruction using acellular dermal grafts. *Ann Plast Surg.* 2013 Sep;71(3):274-7.
116. Moyer HR, Hart AM, Yeager J, Losken A. A Histological Comparison of Two Human Acellular Dermal Matrix Products in Prosthetic-Based Breast Reconstruction. *Plast Reconstr Surg Glob Open.* 2017 Dec 27;5(12):e1576.
117. Musculoskeletal Transplant Foundation. FlexHD® Acellular Hydrated Dermis. Accessed Jan 27, 2021. Available at URL address: <https://www.mtfbionics.org/our-products>
118. Musculoskeletal Transplant Foundation. BellaDerm® Acellular Hydrated Dermis. Accessed Dec 27, 2019. Available at URL address: <https://www.mtfbionics.org/our-products>

119. Nahabedian MY. Secondary nipple reconstruction using local flaps and AlloDerm. *Plast Reconstr Surg*. 2005 Jun;115(7):2056-61.
120. Nahabedian MY, Davies M. Implant-based breast reconstruction and augmentation. Last updated Jan 13, 2020. In: *UpToDate*, Collins KA (Ed), *UpToDate*, Waltham, MA.
121. National Comprehensive Cancer Network (NCCN). NCCN® practice guidelines in oncology. Breast Cancer. V.1.2021-Jan 15, 2021. Accessed Jan 27, 2021. Available at URL address: https://www.nccn.org/professionals/physician_gls/default.aspx
122. National Institute for Health and Clinical Excellence (NICE). Interventional procedure overview IPG417 Interventional procedure overview of breast reconstruction using lipomodelling after breast cancer treatment. Issued January 2012. Accessed Jan 27, 2021. Available at URL address: <http://www.nice.org.uk/>
123. National Library of Medicine. Breast Reconstruction. Reviewed June 21, 2016. Topic last updated Dec 4, 2020. Accessed Jan 27, 2021. Available at URL address: <http://www.nlm.nih.gov/medlineplus/breastreconstruction.html>
124. Nguyen TJ, Carey JN, Wong AK. Use of human acellular dermal matrix in implant-based breast reconstruction: evaluating the evidence. *Jour Plas Recon* (2011):64, 1553-1561.
125. Nguyen AT, Chang EI, Suami H, Chang DW. An algorithmic approach to simultaneous vascularized lymph node transfer with microvascular breast reconstruction. *Ann Surg Oncol*. 2015 Sep;22(9):2919-24.
126. Nilsen TJ, Dasgupta A, Huang YC, Wilson H, Chnari E. Do Processing Methods Make a Difference in Acellular Dermal Matrix Properties? *Aesthet Surg J*. 2016 Nov;36(suppl 2):S7-S22.
127. Ohkuma R, Buretta KJ, Mohan R, Rosson GD, Rad AN. Initial experience with the use of foetal/neonatal bovine acellular dermal collagen matrix (SurgiMend™) for tissue-expander breast reconstruction. *J Plast Reconstr Aesthet Surg*. 2013 Sep;66(9):1195-201.
128. Paprottka FJ, Krezdorn N, Sorg H, Könniker S, Bontikous S, Robertson I, et al. Evaluation of Complication Rates after Breast Surgery Using Acellular Dermal Matrix: Median Follow-Up of Three Years. *Plast Surg Int*. 2017;2017:1283735.
129. Parikh RP, Doren EL, Mooney B, Sun WV, Laronga C, Smith PD. Differentiating fat necrosis from recurrent malignancy in fat-grafted breasts: an imaging classification system to guide management. *Plast Reconstr Surg*. 2012 Oct;130(4):761-72.
130. Parikh RP, Tenenbaum MM, Yan Y, Myckatyn TM. Cortiva Versus AlloDerm Ready-to-use in Prepectoral and Submuscular Breast Reconstruction: Prospective Randomized Clinical Trial Study Design and Early Findings. *Plast Reconstr Surg Glob Open*. 2018 Nov 13;6(11):e2013.
131. Petit JY, Lohsiriwat V, Clough KB, Sarfati I, Ihrari T, Rietjens M, et al. The oncologic outcome and immediate surgical complications of lipofilling in breast cancer patients: a multicenter study-milan-paris-lyon experience of 646 lipofilling procedures. *Plast Reconstr Surg*. 2011a Aug;128(2):341-6
132. Petit JY, Botteri E, Lohsiriwat V, Rietjens M, De Lorenzi F, Garusi C, et al. Locoregional recurrence risk after lipofilling in breast cancer patients. *Ann Oncol*. 2011b May 24.
133. Perez-Cano R, Vranckx JJ, Lasso JM et al. Prospective trial of adipose-derived regenerative cell (ADRC)-enriched fat grafting for partial mastectomy defects: the RESTORE-2 trial. *Eur J Surg Oncol* 2012; 38(5):382-9.

134. Pittman TA, Fan KL, Knapp A, Frantz S, Spear SL. Comparison of Different Acellular Dermal Matrix (ADM) in Breast Reconstruction: The 50/50 Study. *Plast Reconstr Surg.* 2016 Nov 21. [Epub ahead of print]
135. Pulagam SR, Poulton T, Mamounas EP. Long-term clinical and radiologic results with autologous fat transplantation for breast augmentation: case reports and review of the literature. *Breast J.* 2006 Jan-Feb;12(1):63-5.
136. Ramsden AJ, Allen V, O'Donoghue JM. Anterior rectus sheath repair with porcine collagen (Permacol) in patients undergoing breast reconstruction with free abdominal flaps. *J Plast Reconstr Aesthet Surg.* 2009 Jun;62(6):e170-1.
137. Rawlani V, Buck DW 2nd, Johnson SA, Heyer KS, Kim JY. Tissue expander breast reconstruction using prehydrated human acellular dermis. *Ann Plast Surg.* 2011 Jun;66(6):593-7.
138. Rocco N, Rispoli C, Moja L, Amato B, Iannone L, Testa S, et al. Different types of implants for reconstructive breast surgery. *Cochrane Database Syst Rev.* 2016 May 16;(5):CD010895.
139. Roehl KR, Wilhelmi BJ, Phillips LG. Breast Reconstruction. In: Townsend CM, Beauchamp RD, Evers BM, Mattox KL, editors. *Sabiston Textbook of Surgery. The Biological Basis of Modern Surgical Practice.* 19th ed. St. Louis, MO: W.B. Saunders Company; 2012. Ch.37.
140. Roostaeian J, Sanchez I, Vardanian A, Herrera F, Galanis C, Da Lio A, et al. Comparison of immediate implant placement versus the staged tissue expander technique in breast reconstruction. *Plast Reconstr Surg.* 2012 Jun;129(6):909e-918e.
141. Rosenberg JJ, Fornage BD, Chevray PM. Monitoring buried free flaps: limitations of the implantable Doppler and use of color duplex sonography as a confirmatory test. *Plast Reconstr Surg.* 2006 Jul;118(1):109–13; discussion 114–5.
142. Rosing JH, Wong G, Wong MS, Sahar D, Stevenson TR, Pu LL. Autologous fat grafting for primary breast augmentation: a systematic review. *Aesthetic Plast Surg.* 2011 Oct;35(5):882-90.
143. Rowland JH, Desmond KA, Meyerowitz BE, Belin TR, Wyatt GE, Ganz PA. Role of breast reconstructive surgery in physical and emotional outcomes among breast cancer survivors. *Journal of National Cancer Institute.* 2000 Sep;92(17):1422–9.
144. RTI Surgical™. Cortiva™. 2021. Accessed Jan 25, 2021. Available at URL address: http://www.rtix.com/en_us/products/product-implant/cortiva-cortiva-1mm-and-cortiva1mm-tailored-allograft-dermis
145. RTI Surgical™. Cortiva™ and Cortiva™ 1mm allograft dermis implant overview. 2014. Accessed Jan 25, 2021. Available at URL address: https://www.lifehealthcare.com.au/wp-content/uploads/2017/07/Cortiva_Comprehensive_Implant_Overview.pdf
146. Saint-Cyr M, Rojas K, Colohan S, Brown S. The role of fat grafting in reconstructive and cosmetic breast surgery: a review of the literature. *J Reconstr Microsurg.* 2012 Feb;28(2):99-110.
147. Salzberg CA. Noexpansive immediate breast reconstruction using human acellular tissue matrix graft (AlloDerm). *Ann Plast Surg.* 2006 Jul;57(1):1-5.
148. Salzberg CA, Ashikari AY, Koch RM, Chabner-Thompson E. An 8-year experience of direct-to-implant immediate breast reconstruction using human acellular dermal matrix (AlloDerm). *Plast Reconstr Surg.* 2011 Feb;127(2):514-24.
149. Sbitany H, Serletti JM. Acellular dermis-assisted prosthetic breast reconstruction: a systematic and critical review of efficacy and associated morbidity. *Plast Reconstr Surg.* 2011 Dec;128(6):1162-9.

150. Schnarrs RH, Carman CM, Tobin C, Chase SA, Rossmeier KA. Complication Rates With Human Acellular Dermal Matrices: Retrospective Review of 211 Consecutive Breast Reconstructions. *Plast Reconstr Surg Glob Open*. 2016 Nov 21;4(11):e1118.
151. Seth AK, Persing S, Connor CM, Davila A, Hirsch E, Fine NA, Kim JY. A comparative analysis of cryopreserved versus prehydrated human acellular dermal matrices in tissue expander breast reconstruction. *Ann Plast Surg*. 2013 Jun;70(6):632-5.
152. Seth AK, Hirsch EM, Fine NA, Kim JY. Utility of acellular dermis-assisted breast reconstruction in the setting of radiation: a comparative analysis. *Plast Reconstr Surg*. 2012; 130(4):750-758.
153. Sisti A, Grimaldi L, Tassinari J, Cuomo R, Fortezza L, Bocchiotti MA, et al. Nipple-areola complex reconstruction techniques: A literature review. *Eur J Surg Oncol*. 2016 Apr;42(4):441-65.
154. Skovsted Yde S, Brunbjerg ME, Damsgaard TE. Acellular dermal matrices in breast reconstructions - a literature review. *J Plast Surg Hand Surg*. 2016 Aug;50(4):187-96.
155. Sofregen Medical Inc., SERI™ Surgical Scaffold. Accessed Jan 27, 2021. Available at URL address: <https://www.sofregen.com/seri-surgical-scaffold/>
156. Spear SL, Wilson HB, Lockwood MD. Fat injection to correct contour deformities in the reconstructed breast. *Plast Reconstr Surg*. 2005 Oct;116(5):1300-5.
157. Spear SL, Ducic I, Cuoco F, Taylor N. Effect of obesity on flap and donor-site complications in pedicled TRAM flap breast reconstruction. *Plast Reconstr Surg*. 2007 Mar;119(3):788-95.
158. Spear SL, Ducic I, Low M, Cuoco F. The effect of radiation on pedicled TRAM flap breast reconstruction: outcomes and implications. *Plast Reconstr Surg*. 2005 Jan;115(1):84–95.
159. Spear SL, Parikh PM, Reisin E, Menon NG. Acellular dermis-assisted breast reconstruction. *Aesthetic Plast Surg*. 2008 May;32(3):418-25.
160. Synthes® North America. DermaMatrix Acellular Dermis. 2006. Accessed Jan 27, 2021. Available at URL address: <http://synthes.vo.llnwd.net/o16/LLNWMB8/US%20Mobile/Synthes%20North%20America/Product%20Support%20Materials/Brochures/MXBRODermaMatrixAcellularJ7237F.pdf>
161. Topol BM, Dalton EF, Ponn T, Campbell CJ. Immediate single-stage breast reconstruction using implants and human acellular dermal tissue matrix with adjustment of the lower pole of the breast to reduce unwanted lift. *Ann Plast Surg*. 2008 Nov;61(5):494-9.
162. U.S. Food and Drug Administration. 510(k) data base. Accessed Jan 25, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>
163. U.S. Food and Drug Administration (FDA). Products Classification. External aesthetic restoration prostheses. Revised April 1, 2020. Accessed Jan 27, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcr/CFRSearch.cfm?fr=878.3800>
164. U.S. Food and Drug Administration (FDA). Breast implants home page. Updated Sept 28, 2020a. Accessed Jan 25, 2021. Available at URL address: <http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/ImplantsandProsthetics/BreastImplants/default.htm>
165. U.S. Food and Drug Administration (FDA). Labeling for Approved Breast Implants. Updated Oct 23, 2019a. Accessed Jan 27, 2021. Available at URL address:

<http://www.fda.gov/medicaldevices/productsandmedicalprocedures/implantsandprosthetics/breastimplants/ucm063743.htm>

166. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Mentor MemoryGel™ Silicone Gel-Filled Breast Implants (P030053). November 17, 2006a. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
167. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Sientra Silicone Gel-Filled Breast Implants (P070004). March 9, 2012. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
168. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Inamed® Silicone-Filled Breast Implants (P020056). November 17, 2006b. Accessed Jan 27, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
169. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Permacol® (K992556). February 17, 2000. Accessed Jan 27, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN>
170. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Radiesse® (P050037). December 22, 2006c. Accessed Jan 25, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
171. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Juvederm® Voluma XC (P110033). October 22, 2013. Accessed Jan 25, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P110033>
172. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Phasix™ Mesh (K142818). March 31, 2015. Accessed Jan 27, 2021. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf14/K142818.pdf
173. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Biodesign Nipple Reconstruction Cylinder. (K110402). June 20, 2011. Accessed Jan 25, 2021. Available at URL address: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN>
174. U.S. Food and Drug Administration (FDA). SurgiMend Collagen Matrix. (K083898). Feb 4, 2009. Accessed Jan 27, 2021. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf8/K083898.pdf
175. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Ideal Implant® Saline-filled Breast Implant (P120011). Issued November 14, 2014. Updated November 25, 2014. Accessed Jan 27, 2021. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf12/P120011A.pdf
176. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. Mentor MemoryShape Silicone Gel-Filled Breast Implant (P060028). Issued June 14, 2013. Updated June 19, 2013. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>
177. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. SPY® Fluorescent Imaging System. (K073088). Jan 10, 2008. Accessed Jan 25, 2021. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf7/K073088.pdf
178. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. SERI™ Surgical Scaffold (K123128). April 25, 2013. Accessed Jan 27, 2021. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf12/K123128.pdf

179. U.S. Food and Drug Administration (FDA). FDA news release: FDA takes action to protect patients from risk of certain textured breast implants; requests Allergan voluntarily recall certain breast implants and tissue expanders from market. Content current as of July 24, 2019b. Accessed Jan 25, 2021. Available at URL address: <https://www.fda.gov/news-events/press-announcements/fda-takes-action-protect-patients-risk-certain-textured-breast-implants-requests-allergan>
180. U.S. Food and Drug Administration (FDA). Questions and answers about breast implant-associated anaplastic large cell lymphoma (BIA-ALCL). Page last updated Oct 23, 2019c. Accessed Jan 25, 2021. Available at URL address: <https://www.fda.gov/medical-devices/breast-implants/questions-and-answers-about-breast-implant-associated-anaplastic-large-cell-lymphoma-bia-alcl>
181. U.S. Food and Drug Administration (FDA). Inspections, Compliance, Enforcement, and Criminal Investigations. LifeCell Corporation 6/1/15. Accessed Jan 27, 2021. Available at URL address; <http://www.fda.gov/ICECI/EnforcementActions/WarningLetters/2015/ucm449737.htm>
182. U.S. Food and Drug Administration (FDA). Inspections, Compliance, Enforcement, and Criminal Investigations. Allergan Medical 5/29/15. Accessed Jan 27, 2021. Available at URL address; <http://www.fda.gov/iceci/enforcementactions/warningletters/2015/ucm449569.htm>
183. U.S. Food and Drug Administration (FDA). Inspections, Compliance, Enforcement, and Criminal Investigations. TEI Biosciences 5/29/2015. Accessed Jan 27, 2021. Available at URL address; <http://www.fda.gov/iceci/enforcementactions/warningletters/2015/ucm449557.htm>
184. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. ARTIA Reconstructive Tissue Matrix Perforated (K162752). Feb 24, 2017. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?ID=K162752>
185. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. GalaFORM 3D Scaffold (K162922). Feb 23, 2017. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?ID=K162922>
186. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. GalaFLEX™ Mesh (K140533). May 21, 2014. Accessed Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?ID=K140533>
187. U.S. Food and Drug Administration (FDA). Center for Devices and Radiological Health. GalaSHAPE 3D (K161092). Aug 23, 2016. Accessed on Jan 27, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?ID=K161092>
188. U.S. Food and Drug Administration (FDA). Medical Devices. Update on the safety of silicone gel-filled breast implants (2011) - Executive Summary. Updated Jul 16, 2018. Accessed Jan 25, 2021. Available at URL address: <https://www.fda.gov/medical-devices/breast-implants/update-safety-silicone-gel-filled-breast-implants-2011-executive-summary>
189. U.S. Food and Drug Administration (FDA). Tissue & Tissue Products. Jan 11, 2021. Available at URL address: <https://www.fda.gov/vaccines-blood-biologics/tissue-tissue-products>
190. Vardanian AJ, Clayton JL, Roostaeian J, Shirvanian V, Da Lio A, Lipa JE, et al. Comparison of implant-based immediate breast reconstruction with and without acellular dermal matrix. *Plast Reconstr Surg*. 2011 Nov;128(5):403e-410e.
191. Vashi C. Clinical Outcomes for Breast Cancer Patients Undergoing Mastectomy and Reconstruction with Use of DermACELL, a Sterile, Room Temperature Acellular Dermal Matrix. *Plast Surg Int*. 2014;2014:704323.

192. Venturi ML, Mesbahi AN, Boehmler JH 4th, Marrogi AJ. Evaluating sterile human acellular dermal matrix in immediate expander-based breast reconstruction: a multicenter, prospective, cohort study. *Plast Reconstr Surg.* 2013 Jan;131(1):9e-18e. doi: 10.1097/PRS.0b013e3182729d4f. Erratum in: *Plast Reconstr Surg.* 2013 Mar;131(3):669.
193. Wechselberger G, Schoeller T. The transverse myocutaneous gracilis free flap: a valuable tissue source in autologous breast reconstruction. *Plast Reconstr Surg.* 2004 Jul;114(1):69-73.
194. Williams SF, Martin DP, Moses AC. The History of GalaFLEX P4HB Scaffold. *Aesthet Surg J.* 2016 Nov;36(suppl 2):S33-S42.
195. Yoshimura K, Sato K, Aoi N, Kurita M, Hirohi T, Harii K. Cell-assisted lipotransfer for cosmetic breast augmentation: supportive use of adipose-derived stem/stromal cells. *Aesthetic Plast Surg.* 2008 Jan;32(1):48-55; discussion 56-7.
196. Zenn MR. Bipedicled TRAM breast reconstruction. Updated Jan 07, 2020. Accessed Jan 27, 2021. Available at URL address: <https://emedicine.medscape.com/article/1273752-overview>
197. Zenn MR. Free TRAM breast reconstruction. Updated Feb 26, 2019. Accessed Jan 27, 2021. Available at URL address: <https://emedicine.medscape.com/article/1273997-overview>
198. Zenn MR. Unipedicled TRAM breast reconstruction. Updated Feb 14, 2019. Accessed Jan 27, 2021. Available at URL address: <https://emedicine.medscape.com/article/1274334-overview>
199. Zenn MR, Salzberg CA. A Direct Comparison of Alloderm-Ready to Use (RTU) and DermACELL in Immediate Breast Implant Reconstruction. *Eplasty.* 2016 Aug 11;16:e23.
200. Zienowicz RJ, Karacaoglu E. Implant-based breast reconstruction with allograft. *Plast Reconstr Surg.* 2007 Aug;120(2):373-81.
201. Zion SM, Slezak JM, Sellers TA, Woods JE, Arnold PG, Petty PM, et al. Reoperations after prophylactic mastectomy with or without implant reconstruction. *Cancer.* 2003;98(10):2152-60.

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.