



Medical Coverage Policy

Effective Date..... 4/15/2021
Next Review Date..... 4/15/2022
Coverage Policy Number 0309

Surgical Treatment of Chest Wall Deformities

Table of Contents

Overview	1
Coverage Policy.....	1
General Background.....	2
Medicare Coverage Determinations	6
Coding/Billing Information.....	6
References	7

Related Coverage Resources

- [Breast Reconstruction Following Mastectomy or Lumpectomy](#)
- [eviCore Adult and Pediatric Chest Imaging Guidelines \(Pectus Excavatum and Pectus Carinatum; Pectus Deformities\)](#)
- [Transthoracic Echocardiography in Adults](#)
- [Transthoracic Echocardiography in Children](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgement and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses surgical repair of chest wall deformities for pectus excavatum, pectus carinatum and Poland syndrome.

Coverage Policy

Coverage for surgical repair of a chest wall deformity varies across plans, may be subject to the provisions of a cosmetic and/or reconstructive surgery benefit and may be governed by state and/or federal mandates. Please refer to the customer's benefit plan document for coverage details.

If coverage for surgical repair of a chest wall deformity is available, the following conditions of coverage apply.

Surgical repair of pectus excavatum is considered medically necessary when imaging studies (e.g., computerized tomography [CT] scans, radiographs, magnetic resonance imaging [MRI]) confirm a pectus index (i.e., Haller index) greater than 3.25 and EITHER of the following criteria is met:

- Pulmonary function studies demonstrate restrictive or obstructive lung disease
- Cardiac imaging (e.g., echocardiography, stress echocardiography, MRI) demonstrates findings consistent with external cardiac compression

Surgical repair of pectus carinatum is considered medically necessary when there is documented evidence of significant physical functional impairment (e.g., cardiac or respiratory insufficiency), and the procedure is expected to correct the impairment.

Surgical treatment of a chest wall deformity associated with Poland syndrome, including breast reconstruction, is considered medically necessary. Coverage of breast reconstruction is limited to a one- or two-stage approach to correct breast asymmetry.

Note: Please reference the Cigna Medical Coverage policy “Breast Reconstruction Following Mastectomy or Lumpectomy” for conditions of coverage related specifically to breast reconstruction procedures.

Not Covered

When performed for indications other than those listed above, surgical repair of a chest wall deformity is considered not medically necessary.

Magnetic Mini Mover Procedure (3MP) for the treatment of pectus excavatum is considered experimental, investigational or unproven.

General Background

The thorax (i.e. chest cavity) is a rigid structure that protects the thoracic organs and supports the upper extremities. Commonly reported chest wall deformities include pectus excavatum (PE), pectus carinatum (PC) and Poland syndrome. In many cases, primarily cosmetic complaints without functional impairment are associated with the abnormality. However, abnormalities of the chest wall can lead to restrictive pulmonary disease, impaired respiratory muscle strength, and decreased ventilatory performance in response to physical stress (Boas, 2020). Cardiac or respiratory impairment may result in functional limitations, such as activity intolerance. Other symptoms may include mild to moderate exercise limitation, respiratory infections, and asthmatic symptoms, as well as decreased stamina and endurance.

Pectus Excavatum

Pectus excavatum (PE), also referred to as a sunken chest or funnel chest, is the most common congenital chest wall deformity. The deformity may be deeper on the right side than on the left side and result in a rotation of the sternum. It is usually diagnosed within the first year of life, with wide variations in the degree of sternal depression. Although most patients are asymptomatic, during periods of rapid bone growth (e.g., puberty), the appearance of the chest may worsen and symptoms may develop. Moderate to severe deformities may displace the heart into the left chest, decreasing stroke volume and cardiac output. Chest deformities may also depress the sternal volume, adversely affecting the flow of air in and out of the lungs. Scoliosis, congenital heart disease and functional heart murmurs can also be associated with PE. Symptoms may include fatigue, dyspnea, chest discomfort and palpitations with mild exercise. The body generally compensates by increasing the heart rate with activity to overcome the decreased cardiac output and by more rapid, shallow breathing to compensate for the respiratory deficit. Pulmonary effects associated with PE generally include restrictive lung disease, atelectasis, and paradoxical respiration. Restricted lung disease is indicative when the total lung capacity (TLC) (forced vital capacity plus residual capacity) is less than 80% of the predicted value for an individual (Rakel, 2011; Boas, 2020).

Pectus Carinatum

Pectus carinatum (PC) (i.e., pigeon breast or chicken breast) is a congenital chest deformity characterized by an anterior protrusion deformity of the sternum and costal cartilages. PC is typically not confirmed until after the growth spurts of early adolescence. This deformity produces a rigid chest and, while symptoms are uncommon,

it may result in inefficient respiration as a result of the restrictive chest formation. Three types of PC-related defects have been identified in the literature:

- anterior displacement of the body of the sternum and symmetrical concavity of the costal cartilages
- lateral depression of the ribs on one or both sides of the sternum
- pouter pigeon breast (the least common of the three): a defect that consists of an upper or chondromalacial prominence with protrusion of the manubrium and depression of the sternal body

The degree of physiological impairment is related to the degree of chest deformity. Patients with PC may develop symptoms as a result of restricted air exchange; complete expiration of air from the lungs may not occur. In addition, pain may result from the secondary pressures that develop from the overgrowth of cartilage. Other conditions that may be associated with PC include frequent respiratory infections, asthma, rickets, mitral valve disease, Marfan's syndrome, scoliosis and other cardiac changes.

Non-surgical treatment may include orthotic bracing. Bracing may be effective for correction of the chest wall deformity associated with PC, however this type of treatment is often aimed at improving cosmesis. Compression orthotic braces exert pressure on the anteroposterior direction and may be recommended for skeletally immature children (e.g., < age 18 years) with mild to moderate chest deformities. Prolonged use of the orthotic device is often required for total correction and compliance is an important factor for successful remodeling. As a result candidates must be motivated to wear the brace (Goretsky, et al., 2004; Obermeyer, Goretsky, 2012).

Poland Syndrome

Poland syndrome (i.e., Poland's anomaly, Poland's syndactyly), is a rare congenital disorder associated with lateral depression of the ribs on one or both sides of the sternum. The right side of the body is affected twice as often as the left. When the anomaly occurs on the left side of the body, the heart and lungs are vulnerable, because they may be covered only by skin, fascia and pleura. Although the anomaly is associated with a wide range of malformations, the condition is characterized by absence or hypoplasia of the pectoralis major muscle, absence or hypoplasia of the pectoralis minor muscle, absence of costal cartilages, hypoplasia of the breast and subcutaneous tissue, and a variety of hand and upper-extremity anomalies. In cases of severe cartilage deficiency, patients may develop lung hernia and paradoxical respiratory motion. In less severe cases, patients may develop a simple flattening of the anterior chest wall.

Diagnosis and Evaluation

The severity of the chest wall abnormality is dependent upon the depth, symmetry and width of the deformity. Chest radiographs are commonly used to determine the degree of chest wall deformity. Plain anteroposterior and lateral radiographs are used to determine the Haller index (a measurement of chest diameter). Cross-sectional imaging such as computerized tomography (CT) scans and magnetic resonance imaging (MRI) may be used to evaluate the degree of cardiac compression (ventricular compression), pulmonary compression, and cardiac displacement. CT scan ratios that reveal transverse to AP diameter of greater than 3.25 are considered significant for pectus excavatum. A normal chest has an index of 2.5 (Malek, et al., 2003; Fonkalsrud, 2004).

Echocardiography and/or electrocardiography may also be used to evaluate cardiac status. Respiratory status can be determined with the use of pulmonary function studies. In some cases, pulmonary function studies may reveal a restrictive pattern (incomplete lung expansion) and a subsequent decrease in pulmonary volume and reserve. The forced expiratory volume (in one second) (FEV₁), forced vital capacity (FVC), and total lung capacity (TLC) are reduced while the ratio of FEV₁/FVC may be normal or increased in the presence of restrictive airway disease. TLC <80% predicted value signifies restrictive pulmonary disease.

The diagnosis of Poland syndrome is usually obtained by clinical examination. Chest wall abnormalities and determining the presence of latissimus dorsi muscles may require CT scans while chest radiographs may be utilized to evaluate rib formation.

Surgical Treatment

Indications for surgical correction are controversial and vary widely. Surgical repair is offered primarily as a method of improving cosmesis and psychological factors but may be necessary to improve cardiopulmonary function in some patients, as the disfigurement may be accompanied by physiologic impairment.

Pectus Excavatum/Pectus Carinatum: While the optimal age for surgical repair is generally between the ages of 11 and 18 years, and may be performed in adults, each case must be reviewed individually for the presence of impaired cardiopulmonary symptoms. Criteria that may be used to demonstrate severe PE and the need for surgical repair include two or more of the following (Goretsky, et al., 2004):

- a Haller CT index greater than 3.25
- pulmonary function studies that indicate restrictive or obstructive airway disease
- a cardiology evaluation, where the compression is causing murmurs, mitral valve prolapse, cardiac displacement, or conduction abnormalities on the echocardiogram or EKG
- documentation of progression of the deformity with associated physical symptoms other than isolated concerns of body image
- a failed Ravitch procedure
- a failed minimally invasive procedure

Surgery for PE may be performed using any one of several techniques, including a sternal osteotomy (i.e., a modified osteotomy that involves supporting, removing and repositioning the sternum) or implantation of a Silastic mold in the subcutaneous space to fill the defect without altering the thoracic cage. Surgical correction often employs a metal bar behind the sternum; the bar may be removed in one to two years, after remodeling has occurred. The standard surgical procedure is the open Ravitch procedure, which involves extensive dissection, cartilage resection and sternal osteotomy. More recently, minimally invasive techniques, such as the Nuss procedure (i.e. a minimally invasive repair of pectus excavatum [MIRPE]), have been utilized. These procedures involve the insertion of a convex steel bar beneath the sternum through small thoracic incisions and do not require cartilage resection or osteotomy.

Surgical repair may also be recommended for the treatment of PC. Surgical repair of PC may involve removing the affected cartilages and mobilizing the skin and pectoralis muscle flaps. To straighten the sternum, any one of the following surgeries may be performed:

- an osteotomy
- a subperichondrial resection of the involved costal cartilages
- a wedge-shaped osteotomy in the anterior sternal plate

Poland Syndrome: Patients with Poland syndrome typically present for surgical reconstruction to correct deformity or defect of the pectoralis muscles and/or ribs, and to correct breast asymmetry. In cases of rib aplasia, operative reconstruction may eliminate paradoxical motion, improving respiratory impairment. For more severe conditions, reconstructive surgery also provides protection of the underlying heart and lung structures. Surgical repair of the chest wall includes the reconstruction of the pectoral muscles and resection of deformed cartilages. This repair typically involves muscle transfers and/or flaps to match normal development of the unaffected side, reconstruction of the axillary line, and correction of infraclavicular flattening. Techniques for repair of large defects include the use of rib grafts, other bony allo- or autografts, mesh, or a combination of these techniques (Fokin, 2002).

Surgical treatment of Poland syndrome often consists of reconstruction of the breast and nipple on the affected side by a plastic surgeon, in addition to surgical repair of the chest wall muscles and hypoplastic bone. Surgery may be performed early (approximately age 13) in males, however, in females, reconstructive surgery is often deferred until breast development is complete. If there are rib abnormalities and paradoxical motion, the rib grafts or other chest wall stabilization may occur before breast development is complete. Generally, reconstruction of the breast involves tissue expansion, placement of permanent breast implants and may involve myocutaneous or latissimus dorsi flaps if there is an associated anomaly of the pectoral muscle. Nipple-areolar reconstruction is generally performed at a later stage. Consequently, for patients with Poland syndrome,

treatment provided before complete breast development may involve the use of tissue expanders in the affected side which can be inflated periodically to match development of the unaffected breast. Expanders allow for tissue expansion and accommodation of a permanent implant and latissimus muscle upon completion of breast development. Once breast development is complete, the expander is removed and a permanent prosthesis is inserted and breast reconstruction is performed.

U.S. Food and Drug Administration (FDA)

Reconstructive repair of chest wall deformities may involve the placement of metal bars, struts, and stabilizers to correct the defect. These appliances are categorized as Class II medical devices by the U.S. Food and Drug Administration (FDA), and numerous manufacturers have received 510(k) FDA approval for bone fixation devices for repair of PE and other anterior chest wall deformities. Examples include Park's Pectus System (TDM Co. Ltd., Houston, TX) and the Lorenz Pectus Support Bar System (W. Lorenz Surgical, Inc., Jacksonville, FL).

Literature Review

Published evidence evaluating surgical repair of chest wall deformities consists of meta-analyses, case series, cross-comparison studies and prospective trials. The reported clinical outcomes are mixed; differences among outcomes may be related to patient selection criteria, the degree of severity of the deformity, the surgical technique utilized, and growth effects. Many authors evaluated and reported on the methods of surgical repair and improved cosmetic outcomes while some evaluated the impact of PE or PC on cardiopulmonary function. There is little consensus regarding the degree of cardiopulmonary impairment, if any, that is associated with these anomalies. The effects of surgery on exercise tolerance are not clearly established, however, data suggesting improvement in cardiovascular and/or pulmonary function and activity tolerance after surgical repair have been reported (Kelly, et al., 2013; Jaroszewski, Fonkalsrud, 2007; Kubiak, et al., 2007; Lawson, et al., 2005; Bawazir, et al., 2005; Fonkalsrud and Anselmo, 2004; Haller and Loughlin, 2000; Fonkalsrud, et al., 1994). Outcome measures of these studies generally included total lung capacity (TLC), functional residual capacity (FRC), vital capacity, (VC), expiratory flow rate (EFR), and maximum expiratory flow rate (MEFR), exercise tolerance and endurance typically measured prior to surgery, immediately following surgery and three to six months postoperatively. Improvement was generally seen only with increased periods of exercise and not with routine pulmonary function testing at rest. The results of some meta-analyses and other published clinical studies in the medical literature are also mixed, some results tend to support improvement in cardiopulmonary function following surgery (Johnson, et al., 2008; Malek, et al., 2006a, Malek, et al., 2006b) while others do not (Zganjer and Zganjer, 2010; Castellani, et al, 2010; Guntheroth and Spiers, 2007).

Magnetic Mini-Mover Procedure (3MP)

A procedure that is currently under investigation for the treatment of PE is the Magnetic Mini-Mover Procedure (3MP) which utilizes a magnetically coupled implant to pull the sternum forward and remodel the deformed costal cartilage. The internal magnetic implant is surgically placed on the sternum and an external magnet is applied using a custom-fitted chest wall orthotic device. The Magnetic Mini-Mover device is currently under investigation in a United States Food and Drug Administration (FDA) sponsored Phase III Investigational Device Exemption (IDE) clinical trial (NCT01327274); IDE approval allows the device to be used in a clinical study for the collection of safety and effectiveness data. Current evidence evaluating the magnetic mini mover procedure in the published peer-reviewed scientific literature is primarily in the form of pilot studies and small case series (n=15) (Graves, et al., 2017; Harrison, et al 2012; Harrison, et al., 2007).

Professional Societies/Organizations

The American Pediatric Surgical Association (APSA) published its Pectus Carinatum Guideline in 2012 (APSA, 2012). The guideline recommendations were based upon a systematic review of medical literature and peer-reviewed journals in or after 1973. Recommendations pertaining to surgical intervention included:

- Operative therapy plans should be individualized and age-appropriate, with surgery typically being deferred until pubertal growth is nearly complete
- Open surgical techniques are acceptable for repair when performed by experienced pediatric surgeons
- Thoracoscopic and other minimally invasive techniques are acceptable in some children, when performed by pediatric surgeons with advanced skills and experience in the technique
- Surgery in children ages 5 years and younger is discouraged due to the risk of disruption of normal chest wall growth and chest wall restriction

- The pediatric surgeon should be involved in extended post-operative follow up

Use Outside of the US: No relevant information.

Medicare Coverage Determinations

	Contractor	Policy Name/Number	Revision Effective Date
NCD		No National Coverage Determination found	
LCD		No Local Coverage Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.

Coding/Billing Information

Note: 1) This list of codes may not be all-inclusive.

- 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Pectus Carinatum/Pectus Excavatum

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
21740	Reconstructive repair of pectus excavatum or carinatum; open
21742	Reconstructive repair of pectus excavatum or carinatum; minimally invasive approach (Nuss procedure), without thoracoscopy
21743	Reconstructive repair of pectus excavatum or carinatum; minimally invasive approach (Nuss procedure), with thoracoscopy

Poland Syndrome

Considered medically necessary when used to report surgical repair of a chest deformity associated with Poland syndrome:

CPT®* Codes	Description
15734	Muscle, myocutaneous, or fasciocutaneous flap; trunk
15756	Free muscle or myocutaneous flap with microvascular anastomosis
20900	Bone graft, any donor area; minor or small (eg, dowel or button)
20902	Bone graft, any donor area; major or large

Considered Experimental, Investigational, Unproven when used to report Magnetic Mini Mover (3MP) procedure:

CPT®* Codes	Description
21743	Reconstructive repair of pectus excavatum or carinatum; minimally invasive approach (Nuss procedure), with thoracoscopy

*Current Procedural Terminology (CPT®) ©2020 American Medical Association: Chicago, IL.

References

1. American Pediatric Surgical Association (APSA). Pectus Carinatum Guideline. 2012. Accessed February 19, 2021. Available at URL address: <https://apsaped surg.org/resources/resources/practice-guidelines/apsa-guidelines/>
2. Aronson DC, Bosgraaf RP, Merz EM, van Steenwijk RP, van Aalderen WM, van Baren R. Lung function after the minimal invasive pectus excavatum repair (Nuss procedure). *World J Surg.* 2007 Jul;31(7):1518-22.
3. Ateş O, Karakuş OZ, Hakgüder G, Olguner M, Akgür FM. Pectus carinatum: the effects of orthotic bracing on pulmonary function and gradual compression on patient compliance. *Eur J Cardiothorac Surg.* 2013 Sep;44(3):e228-32.
4. Bawazir OA, Montgomery M, Harder J, Sigalet DL. Midterm evaluation of cardiopulmonary effects of closed repair of pectus excavatum. *J Pediatr Surg.* 2005 May;40(5):863-7.
5. Boas S. Pectus Excavatum. Skeletal Diseases Influencing Pulmonary Function. In: Behrman RE, Kliegman RM, Jenson HB, editors; *Nelson Textbook of Pediatrics*, 21st ed. Copyright © 2020. Chapter 445, 2326-2329.e3
6. Bohosiewicz J, Kudela G, Koszutski T. Results of Nuss procedure for the correction of pectus excavatum. *Eur J Pediatr Surg.* 2005 Feb;15(1):6-10.
7. Brigato RR, Campos JR, Jatene FB, Moreira LF, Rebeis EB. Pectus excavatum: evaluation of Nuss technique by objective methods. *Interact Cardiovasc Thorac Surg.* 2008 Dec;7(6):1084-8.
8. Castellani C, Windhaber J, Schober PH, Hoellwarth ME. Exercise performance testing in patients with pectus excavatum before and after Nuss procedure. *Pediatr Surg Int.* 2010 Jul;26(7):659-63.
9. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed February 18, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx>
10. Centers for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed February 18, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/ncd-alphabetical-index.aspx>
11. Cheng YL, Lee SC, Huang TW, Wu CT. Efficacy and safety of modified bilateral thoracoscopy-assisted Nuss procedure in adult patients with pectus excavatum. *Eur J Cardiothorac Surg.* 2008 Nov;34(5):1057-61.
12. Cohee AS, Lin JR, Frantz FW, Kelly RE Jr. Staged management of pectus carinatum. *J Pediatr Surg.* 2013 Feb;48(2):315-20.
13. Coln E, Carrasco J, Coln D. Demonstrating relief of cardiac compression with the Nuss minimally invasive repair for pectus excavatum. *J Pediatr Surg.* 2006 Apr;41(4):683-6; discussion 683-6.
14. de Oliveira Carvalho PE, da Silva MV, Rodrigues OR, Cataneo AJ. Surgical interventions for treating pectus excavatum. *Cochrane Database Syst Rev.* 2014 Oct 29;10:CD008889. doi: 10.1002/14651858.CD008889.pub2.
15. Desmarais TJ, Keller MS. Pectus carinatum. *Curr Opin Pediatr.* 2013 Jun;25(3):375-81.
16. Fokin AA, Robicsek F. Poland's syndrome revisited. *Ann Thorac Surg.* 2002 Dec;74(6):2218-25.

17. Fonkalsrud EW, Anselmo DM. Less extensive techniques for repair of pectus carinatum: the undertreated chest deformity. *J Am Coll Surg*. 2004 Jun;198(6):898-905.
18. Fonkalsrud EW, Beanes S. Surgical management of pectus carinatum: 30 years' experience. *World J Surg*. 2001 Jul;25(7):898-903.
19. Fonkalsrud EW. Current management of pectus excavatum. *World J Surg*. 2003 May;27(5):502-8.
20. Fonkalsrud EW, DeUgarte D, Choi E. Repair of pectus excavatum and carinatum deformities in 116 adults. *Ann Surg*. 2002 Sep;236(3):304-12; discussion 312-4.
21. Fonkalsrud EW. Management of pectus deformities in female patients. *Am J Surg*. 2004 Feb;187(2):192-7.
22. Fonkalsrud EW, Mendoza J. Open repair of pectus excavatum and carinatum deformities with minimal cartilage resection. *Am J Surg*. 2006 Jun;191(6):779-84.
23. Fonkalsrud EW, Salman T, Guo W, Gregg JP. Repair of pectus deformities with sternal support. *J Thorac Cardiovasc Surg*. 1994;107:37-42.
24. Frantz FW. Indications and guidelines for pectus excavatum repair. *Curr Opin Pediatr*. 2011 Aug;23(4):486-91.
25. Frey AS, Garcia VF, Brown RL, Inge TH, Ryckman FC, Cohen AP, et al. Nonoperative management of pectus carinatum. *J Pediatric Surg*. 2006 Jan;41(1):40-5;discussion 40-5.
26. Goretsky MJ, Kelly RE, Croitoru D, Nuss D. Chest wall anomalies: pectus excavatum and pectus carinatum. *Adolesc Med Clin*. 2004 Oct;15(3):455-71.
27. Graves CE, Hirose S, Raff GW, Iqbal CW, Imamura-Ching J, Christensen D, Fechter R, Kwiat D, Harrison MR. Magnetic Mini-Mover Procedure for pectus excavatum IV: FDA sponsored multicenter trial. *J Pediatr Surg*. 2017 Jun;52(6):913-919.
28. Guntheroth WG; Spiers PS. Cardiac function before and after surgery for pectus excavatum. *Am J Cardiol*. 2007 Jun;99(12):1762-4.
29. Haller JA Jr., Loughlin GM. Cardiorespiratory function is significantly improved following corrective surgery for severe pectus excavatum: proposed treatment guidelines. *J Cardiovasc Surg (Torino)*. 2000 Feb;41(1):125-30.
30. Harrison MR, Estefan-Ventura D, Fechter R, Moran AM Jr, Christensen D. Magnetic Mini-Mover Procedure for pectus excavatum: I. Development, design, and simulations for feasibility and safety. *J Pediatr Surg*. 2007 Jan;42(1):81-5; discussion 85-6.
31. Harrison MR, Gonzales KD, Bratton BJ, Christensen D, Curran PF, Fechter R, Hirose S. Magnetic mini-mover procedure for pectus excavatum III: safety and efficacy in a Food and Drug Administration-sponsored clinical trial. *J Pediatr Surg*. 2012 Jan;47(1):154-9.
32. Jaroszewski DE, Fonkalsrud EW. Repair of pectus chest deformities in 320 adult patients: 21 year experience. *Ann Thorac Surg*. 2007 Aug;84(2):429-33.
33. Jayaramakrishnan K, Wotton R, Bradley A, Naidu B. Does repair of pectus excavatum improve cardiopulmonary function? *Interact Cardiovasc Thorac Surg*. 2013 Jun;16(6):865-70.

34. Johnson JN, Hartman TK, Pianosi PT, Driscoll DJ. Cardiorespiratory function after operation for pectus excavatum. *J Pediatr*. 2008 Sep;153(3):359-64.
35. Kelly RE Jr, Cash TF, Shamberger RC, Mitchell KK, Mellins RB, Lawson ML, et al. Surgical repair of pectus excavatum markedly improves body image and perceived ability for physical activity: multicenter study. *Pediatrics*. 2008 Dec;122(6):1218-22.
36. Kelly RE, Goretsky MJ, Obermeyer R, Kuhn MA, Redlinger R, Haney TS, Moskowitz A, Nuss D. Twenty-one years of experience with minimally invasive repair of pectus excavatum by the Nuss procedure in 1215 patients. *Ann Surg*. 2010 Dec;252(6):1072-81.
37. Kelly RE Jr, Mellins RB, Shamberger RC, Mitchell KK, Lawson ML, Oldham KT, Azizkhan RG, Hebra AV, Nuss D, Goretsky MJ, Sharp RJ, Holcomb GW 3rd, Shim WK, Megison SM, Moss RL, Fecteau AH, Colombani PM, Cooper D, Bagley T, Quinn A, Moskowitz AB, Paulson JF. Multicenter study of pectus excavatum, final report: complications, static/exercise pulmonary function, and anatomic outcomes. *J Am Coll Surg*. 2013 Dec;217(6):1080-9.
38. Kelly RE Jr; Shamberger RC; Mellins RB; Mitchell KK; Lawson ML; Oldham K; et al. Prospective multicenter study of surgical correction of pectus excavatum: design, perioperative complications, pain, and baseline pulmonary function facilitated by internet-based data collection. *J Am Coll Surg*. 2007 Aug;205(2):205-16.
39. Kubiak R, Habelt S, Hammer J, Häcker FM, Mayr J, Bielek J. Pulmonary function following completion of Minimally Invasive Repair for Pectus Excavatum (MIRPE). *Eur J Pediatr Surg*. 2007 Aug;17(4):255-60.
40. Lawson ML, Mellins RB, Paulson JF, Shamberger RC, Oldham K, Azizkhan RG, et al. Increasing severity of pectus excavatum is associated with reduced pulmonary function. *J Pediatr*. 01-AUG-2011; 159(2): 256-61.e2.
41. Lawson ML, Mellins RB, Tabangin M, Kelly Jr RE, Croitoru DP, Goretsky MJ, Nuss D. Impact of pectus excavatum on pulmonary function before and after repair with the Nuss procedure. *J Pediatr Surg*. 2005 Jan;40(1):174-80;discussion 180.
42. Lee RT, Moorman S, Schneider M, Sigalet DL. Bracing is an effective therapy for pectus carinatum: interim results. *J Pediatr Surg*. 2013 Jan;48(1):184-90.
43. Lopez M, Patoir A, Varlet F, Perez-Etchepare E, Tiffet T, Villard A, Tiffet O. Preliminary study of efficacy of dynamic compression system in the correction of typical pectus carinatum. *Eur J Cardiothorac Surg*. 2013 Nov;44(5):e316-9.
44. Maagaard M, Tang M, Ringgaard S, Nielsen HH, Frøkiær J, Haubuf M, Pilegaard HK, Hjortdal VE. Normalized cardiopulmonary exercise function in patients with pectus excavatum three years after operation. *Ann Thorac Surg*. 2013 Jul;96(1):272-8.
45. Malek MH, Berger DE, Housh TJ, Marelich WD, Coburn JW, Beck TW. Cardiovascular function following surgical repair of pectus excavatum: a metaanalysis. *Chest*. 2006a Aug;130(2):506-16.
46. Malek MH, Berger DE, Marelich WD, Coburn JW, Beck TW, Housh TJ. Pulmonary function following surgical repair of pectus excavatum: a meta-analysis. *Eur J Cardiothorac Surg*. 2006b Oct;30(4):637-43.
47. Malek MH, Fonkalsrud EW, Cooper CB. Ventilatory and cardiovascular responses to exercise in patients with pectus excavatum. *Chest*. 2003 Sep;124(3):870-82.
48. Mansour KA, Thourani VH, Odessey FA, Durham MM, Miller JI Jr, Miller DI. Thirty-year experience with repair of pectus deformities in adults. *Ann Thorac Surg*. 2003 Aug;76(2):391-5;discussion 395.

49. Mayer, OH. Pectus excavatum: etiology and evaluation. In: UpToDate, Redding, G (ed.). Jan 21, 2020. UpToDate, Waltham, MA. Accessed February 18, 2021.
50. Mayer, OH. Pectus excavatum: treatment. In: UpToDate, Redding, G (ed.). Jun 15, 2020. UpToDate, Waltham, MA. Accessed February 18, 2021.
51. Nuchtern, JG and Mayer, OH. Pectus carinatum. In: UpToDate, Redding, G (ed.). Dec 1, 2020. UpToDate, Waltham, MA. Accessed February 19, 2021.
52. Obermeyer RJ, Goretsky MJ. Chest wall deformities in pediatric surgery. *Surg Clin North Am.* 2012 Jun; 92(3): 669-84.
53. Oncel M, Tezcan B, Akyol KG, Dereli Y, Sunam GS. Clinical experience of repair of pectus excavatum and carinatum deformities. *Cardiovasc J Afr.* 2013 Sep;24(8):318-21.
54. Petersen C, Leonhardt J, Duderstadt M, Karck M, Ure BM. Minimally invasive repair of pectus excavatum - shifting the paradigm? *Eur J Pediatr Surg.* 2006 Apr;16(2):75-8.
55. Poston PM, McHugh MA, Rossi NO, Patel SS, Rajput M, Turek JW. The case for using the correction index obtained from chest radiography for evaluation of pectus excavatum. *J Pediatr Surg.* 2015 Nov;50(11):1940-4.
56. Poston PM, Patel SS, Rajput M, Rossi NO, Ghanamah MS, Davis JE, Turek JW. The correction index: setting the standard for recommending operative repair of pectus excavatum. *Ann Thorac Surg.* 2014 Apr;97(4):1176-9; discussion 1179-80.
57. Rakel: Textbook of Family Medicine, 9th ed. Ch 16. Pulmonary Medicine. Pulmonary function testing. Copyright ©2015 Saunders.
58. Robicsek F, Watts LT. Pectus carinatum. *Thorac Surg Clin.* 2010 Nov;20(4):563-74.
59. Sarwar ZU, DeFlorio R, O'Connor SC. Pectus excavatum: current imaging techniques and opportunities for dose reduction. *Semin Ultrasound CT MR.* 2014 Aug;35(4):374-81.
60. Sigalet DL, Montgomery M, Harder J. Cardiopulmonary effects of closed repair of pectus excavatum. *J Pediatr Surg.* 2003 Mar;38(3):380-5; discussion 380-5.
61. St Peter SD, Juang D, Garey CL, Laituri CA, Ostlie DJ, Sharp RJ, Snyder CL. A novel measure for pectus excavatum: the correction index. *J Pediatr Surg.* 2011 Dec;46(12):2270-3. doi: 10.1016/j.jpedsurg.2011.09.009.
62. Townsend. Chest wall. In: Sabiston Textbook of Surgery, 17th ed. Elsevier; 2004. p. 1711-4.
63. Warner BW. What's new in pediatric surgery. *J Am Coll Surg.* 2004 Aug;199(2):273-85.
64. Zhao L, Feinberg MS, Gaides M, Ben-Dov I. Why is exercise capacity reduced in subjects with pectus excavatum? *J Pediatr.* 2000 Feb;136(2):163-7.
65. Zganjer M, Zganjer V. Surgical correction of the funnel chest deformity in children. *Int Orthop.* 2010 Dec

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.