



Medical Coverage Policy

Effective Date..... 4/15/2021
Next Review Date..... 4/15/2022
Coverage Policy Number 0335

Otoplasty and External Ear Reconstruction

Table of Contents

Overview	1
Coverage Policy.....	1
General Background.....	2
Medicare Coverage Determinations	5
Coding/Billing Information.....	5
References	6

Related Coverage Resources

- [Cochlear and Auditory Brainstem Implants](#)
- [Prosthetic Devices](#)
- [Hearing Aids](#)
- [Scar Revision](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgement and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses otoplasty and external ear reconstruction. Otoplasty, a procedure to correct protruding ears, is performed to improve the appearance of ears. External ear reconstruction is a surgical procedure that attempts to reconstruct the external ear to normal anatomical shape.

Coverage Policy

Coverage for otoplasty and/or external ear reconstruction is dependent on benefit plan language and may be subject to the provisions of a cosmetic and/or reconstructive surgery benefit. In addition, this service may be governed by state mandates. Please refer to the applicable benefit plan documents and schedule of copayments to determine benefit availability and the terms, conditions and limitations of coverage.

External ear reconstruction for the treatment of an external ear deformity or the absence of an external ear is considered medically necessary when ANY of the following criteria is met:

- hearing is expected to improve

- reconstruction is necessary to allow the use of a conventional air conduction hearing aid
- photographs demonstrate that the external ear deformity is preventing the functional ability to use eyewear for the correction of visual loss

Non-surgical external ear molding is considered medically necessary for a congenital external ear malformation with a functional impairment of hearing.

Each of the following are considered cosmetic in nature and not medically necessary when performed solely to improve physical appearance:

- external ear reconstruction
- ear molding

Otoplasty (CPT® code 69300) is considered cosmetic in nature and not medically necessary for ANY indication, including ALL of the following:

- prominent/protruding ears
- lop ears
- cupped ears
- constricted ears

General Background

Abnormal ear development may result from trauma or disease although most often the deformity is congenital. While some abnormalities require no intervention and will self-correct (e.g., caused by abnormal positioning in utero), some may be corrected non-surgically (e.g., ear molds), and others may require surgical correction.

Nonsurgical treatment (e.g., ear molds) is generally employed as treatment shortly after birth when infant ear cartilage is soft and moldable. Ear molds used to improve the appearance of the external ear in the absence of a functional deficit is considered cosmetic.

External ear reconstruction involves various degrees of surgical repair and may be performed to correct the congenital absence of an external ear for conditions such as microtia and anotia or to correct an external ear that has been altered as a result of trauma or surgery. External ear deformities usually do not result in a functional deficit, such as hearing impairment (i.e., inability to hear normal conversation). In the absence of hearing impairment or other functional deficit external ear deformities generally do not require any intervention; treatment is considered cosmetic.

Otoplasty is a procedure performed solely for cosmetic purposes.

Congenital Abnormalities

Prominent/Protruding Ears: Prominent ears are a congenital abnormality in which the ears tend to project excessively from the skull without causing a functional deficit (Liaw, et al., 2017). This condition may occur as a result of an inadequately formed antihelix (i.e., the outer frame of the auricle), an overdeveloped or excessively deep concha (i.e., hollow portion of the outer ear), or a combination of these conditions (American Society of Plastic Surgeons [ASPS], 2005; Reaffirmed June 2015). Normal prominence is defined as 1.2–2.0 cm from the post-auricular scalp to the lateral aspect of the superior helix. Ear prominence is typically defined as a protrusion of the helix 2 cm or more from the postauricular scalp. Otoplasty performed to correct prominent ears involves recreating an antihelical fold and possibly in setting or resecting the concha to decrease the prominence. The primary goal of surgical correction for prominent/protruding ears is improvement of physical appearance (i.e., cosmesis). In comparison to conventional methods which usually involve a post-auricular incision, incisionless otoplasty is a noninvasive method of correction under investigation that involves use of percutaneous sutures and scoring of the cartilage. It has been suggested incisionless otoplasty may result in fewer complications when compared to conventional methods.

Microtia: Microtia describes an incompletely formed ear and is commonly associated with congenital aural atresia (Murakami, et al., 2010; Kelley and Scholes, 2007). It may occur as a single disorder, as a feature of hemifacial microsomia complex (i.e., one side of the face does not grow in proportion to the other side), or as part of a congenital syndrome, such as Treacher Collin's syndrome. While there is no universally accepted classification system for microtia, a system that assigns grades based on the severity of the deformity has been adopted (Murakami, et al., 2010; Zim, 2003). Microtia may be divided into the following categories:

- Type I** A mildly deformed ear that has a slightly dysmorphic helix and antihelix. The external auditory meatus is usually present.
- Type II** Ears that have all major structures present to some degree, but with an absolute deficiency of tissue; surgical correction requires the addition of cartilage and skin; the external auditory meatus is present but may demonstrate some degree of deformity. The auricle is usually hook-, S- or question-mark shaped in appearance.
- Type III** Few or no recognizable landmarks of the auricle or canal although the lobule is usually present and positioned anteriorly.

Microtia may result in subtle abnormalities of the size, shape and location of the pinna and ear canal, or it may occur as a major deformity, with small remnants of skin and cartilage, as well as absence of the ear canal opening. Mild ear deformities are associated with altered physical appearance and are usually not associated with a functional deficit. Deformities that may be considered Type I deformities include mildly constricted ears, lop-ear deformities (characterized by an absence of the antihelical fold causing the ear to fall forward) and cupped-ear deformities (excessive cartilage of the ear canal causing the ear to project outward). With these deformities, all major structures are present to some degree.

Type II deformities may include mini-ear and severe cup deformities. The external auditory meatus is present, although it may demonstrate some degree of stenosis.

Anotia is the complete absence of the external ear and auditory canal and may be considered Type III microtia, although a few sources consider this a fourth degree of severity.

The inner ear function of the affected ear usually remains adequate, resulting in some ability to hear on the affected side (Bonilla, 2021) and the contralateral ear is usually normal, allowing for normal development of speech. However, sensorineural, conductive, or mixed hearing loss may be present in the microtia patient (Kelley and Scholes, 2007; Beahm and Walton, 2002) and it has been reported that hearing impairment may be reduced by approximately 40% on the affected side. Congenital deformities of the ear may be coupled with abnormalities involving the external ear canal (EAC) and tympanic membrane; consequently, these abnormalities may affect sound conduction. Microtia has also been associated with middle ear abnormalities; patients with complete or partial stenosis of the EAC commonly have severe ossicular malformations (Kim, et al., 2002). Additionally, malformations of the external ear can interfere with the ability to wear hearing aids or glasses, which is of particular importance given the high incidence of craniofacial or ocular abnormalities associated with microtia (Cuccolo, et al., 2019).

Ear reconstruction may be performed to improve physical appearance for patients with microtia, although, when considering surgery, emphasis is also placed on restoring sufficient hearing to allow normal speech development. The aim of reconstructive surgery is to improve hearing to a point where the child may not need a hearing aid or to provide an ear canal and pinna so that the child can use an air-conduction hearing aid (Haddad and Dodhia, 2020). Other operations, such as canal or middle ear reconstruction, may be performed to improve patient outcomes. Surgery performed to improve hearing is recommended if there are bilateral deformities resulting in conductive hearing loss (Haddad, 2011) or for unilateral microtia with impaired hearing of the normal ear (Medicare Services Advisory Committee [MSAC], 2000). In patients with bilateral microtia, bone conduction hearing aids are often recommended (Murakami, et al., 2010; Kelley and Scholes, 2007). Hearing amplification is not usually required for unilateral atresia, although binaural hearing is superior to monaural in terms of sound localization and speech perception.

Although it may be performed on adults, it is generally recommended that external ear reconstruction for treatment of ear deformities, more specifically microtia, be performed when the patient is between ages six and

eight. By this age, the ear has reached 85–90% of its adult size. In addition, at this time, the patient's rib size is sufficient to allow a rib graft. Early surgery may also result in the avoidance of social problems for the child. In cases of bilateral microtia, reconstructions may begin as early as age four.

Trauma/Neoplasm

Trauma to the ear may result from burn injuries, human or animal bites, falls or motor vehicle accidents. The unavoidable exposure to sun of the helical rim of the ear contributes to the development of skin neoplasm and removal with precise margin control is recommended. Despite efforts to preserve healthy tissue in the presence of tissue injury or neoplasm, reconstruction is often necessary to improve physical appearance and function. For some patients, auricular prostheses may be considered an alternative to ear reconstruction. Nonetheless, reconstruction to improve physical appearance in the absence of improving function is considered cosmetic.

Cochlear Implant

Sensorineural hearing loss may occur as a result of congenital defects, disease or trauma of the inner ear, and can cause significant hearing impairment. When the hearing loss becomes profound and a hearing aid is ineffective, a cochlear implant may maximize hearing ability for patients.

Cochlear implants have two integral components:

- The internal component consists of a receiver-stimulator connected to an intracochlear electrode array made up of electrode rings that are integrated into a silicone carrier. The stimulator is implanted in the skull near the cochlea and is connected to the electrode array via a lead wire.
- The external component consists of a microphone worn on the external ear, a speech processor worn at ear-level or on the body, and a transmitter worn behind the ear.

Although considered rare, sensorineural hearing loss may occur with congenital ear anomalies such as aural atresia and microtia. Aural atresia is a congenital defect characterized by malformations of the external and middle ear structures. Consequently, otoplasty may be considered as part of the reconstructive process and implantation of a cochlear device. Cochlear microphone placement may be difficult in some cases and external ear reconstruction may be required to facilitate use of the device.

Treatment

Non-surgical treatment of congenital external ear malformation includes splinting and molding. The ideal treatment is initiated within the first two weeks of life and may continue until the infant is three months old. Splints should be light, inexpensive, quick to fabricate, adjustable to complicated cartilage contours, simple to apply and remove, and easy to reform. The molding process involves the application of a rigid cradle that surrounds the patient's auricle and adheres firmly to the skin (Schultz, et al., 2017). Nonsurgical treatment of microtia, involving a prosthetic device, is an alternative to surgical correction. Bone-anchored hearing aid devices are often used to improve conductive hearing loss for cases of bilateral microtia involving hearing impairment.

Surgical repair is generally performed for cosmetic purposes and in some rare situations, functional reasons. The overall goal is to reconstruct an ear that is normal in appearance and function. For some cases an incision is made behind the ear to reduce one or more components, for other more extensive cases reconstruction may involve cartilage reshaping and sculpturing. The reconstruction surgery for severe cases typically involves multiple stages that are performed at least three to six months apart. The initial stages involve the removal of scarred, deformed tissue and the implantation of costal cartilage (e.g., rib cartilage grafting); additional stages are performed for lobule transfer, postauricular skin grafting and tragus reconstruction (Murakami, et al., 2010). Although numerous implants are available for surgical reconstruction of the ear, the gold standard of therapy for treating microtia deformities is autologous rib cartilage grafting. In cases where there is associated aural atresia or decreased hearing in the contralateral normal ear, a separate surgery is indicated to restore hearing function. Medpor ear reconstruction (i.e., porous polyethylene framework and temporoparietal fascial flap) is a well-established alternative choice of treatment for microtia and atresia surgery. Medpor ear reconstruction involves a single-stage reconstruction with a Medpor implant versus multiple-staged approach using rib graft reconstruction. There are several types of porous polyethylene implants which include, but are not limited to Medpor, Omnipor and Supor.

Complications associated with otoplasty and/or external ear reconstructive procedures include bleeding, infection and possibly pneumothorax if a rib graft is used. Complications associated with middle ear surgery for improvement of hearing include restenosis of the external auditory canal and damage to the facial nerve (Bonilla, 2021).

Professional Societies/Organizations

American Academy of Pediatrics (AAP): Guidelines and/or position statements from the AAP do not comment on the performance of otoplasty for treatment of external ear deformities.

American Society of Plastic Surgeons (ASPS): According to the ASPS, otoplasty is considered a reconstructive surgery that may be performed in children or adults, although the procedure is more common in children (ASPS, 2005; Reaffirmed June 2015).

Use Outside of the US: The National Institute for Health and Care Excellence (NICE) issued a procedural guidance for incisionless otoplasty (NICE, 2012) using minimal percutaneous access for the treatment of protruding or prominent ears. Interventional procedure guidance documents were not found for conventional otoplasty or external ear reconstruction. According to the guidance, incisionless otoplasty should only be used with special arrangements for clinical governance, consent and audit, or research due to inadequate evidence.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No National Coverage Determination found	
LCD		No Local Coverage Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.

Coding/Billing Information

Note: 1) This list of codes may not be all-inclusive.

2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

External Ear Reconstruction

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
69310	Reconstruction of external auditory canal (meatoplasty) (eg, for stenosis due to injury, infection) (separate procedure)
69320	Reconstruction external auditory canal for congenital atresia, single stage
69399 [†]	Unlisted procedure, external ear

†Note: Considered Medically Necessary when used to report ear molding and the above criteria in the policy statement is met.

Otoplasty

Considered Cosmetic/Not Medically Necessary

CPT®* Codes	Description
69300	Otoplasty, protruding ear, with or without size reduction

References

1. Adamson PA, Doud Galli SK, Chen T. Otoplasty. Cummings CW, Flint PW, Haughey BH, Robbins KT, Thomas JR, Harker LA, et al. editors. In: Otolaryngology: Head and Neck Surgery, 5th ed., Ch 33, Otoplasty. Copyright ©2010 Mosby, Inc.
2. Adamson PA, Litner JA. Otoplasty technique. *Otolaryngol Clin North Am.* 2007 Apr;40(2):305-18.
3. American Society of Plastic Surgeons (ASPS). Ear deformity: prominent ears: recommended criteria for third-party payer coverage [position paper]. Socioeconomic Subcommittee. Approved by ASPS Board of December 2005. Reaffirmed June 2015. Accessed February 11, 2021. Available at URL address: <http://www.plasticsurgery.org/for-medical-professionals/legislation-and-advocacy/health-policy-resources/recommended-insurance-coverage-criteria.html>
4. Beahm EK, Walton RL. Auricular reconstruction for microtia: Part I: anatomy, embryology, and clinical evaluation. *Plast Reconstr Surg.* 2002 Jun;109(7):2473-82.
5. Bonilla JA. Microtia. In: eMedicine specialties > pediatrics > otolaryngology. Updated January 4, 2021. Accessed February 11, 2021. Available at URL address: <http://www.emedicine.com/ped/topic3003.htm>
6. Brent B. Technical advances in ear reconstruction with autogenous rib cartilage grafts: personal experience with 1200 cases. *Plast Reconstr Surg.* 1999 Aug;104(2):319-34.
7. Brent B. Reconstructive ear surgery. Microtia, Auricular Reconstruction. Copyright © 1998-2011. Accessed February 11, 2021. Available at URL address: <http://www.microtia.us.com/>
8. Brodland DG. Auricular reconstruction. *Dermatol Clin.* 2005 Jan;23(1):23-14, v.
9. Burns JL, Blackwell SJ. Anotia. In: Townsend CM Jr, Beauchamp RD, Evers BM, Mattox KL, editors. Sabiston textbook of surgery. 19th ed. Philadelphia, PA: Saunders; 2012. Chapter 69: plastic surgery. p. 2189.
10. Cuccolo NG, Zwierstra MJ, Ibrahim AMS, Peymani A, Afshar S, Lin SJ. Reconstruction of Congenital Microtia and Anotia: Analysis of Practitioner Epidemiology and Postoperative Outcomes. *Plast Reconstr Surg Glob Open.* 2019 Jun 19;7(6):e2318.
11. Daniali LN, Rezzadeh K, Shell C, Trovato M, Ha R, Byrd HS. Classification of Newborn Ear Malformations and their Treatment with the EarWell Infant Ear Correction System. *Plast Reconstr Surg.* 2017 Mar;139(3):681-691.
12. Ear Plastic Surgery. American Academy of Otolaryngology—Head and Neck Surgery (AAO-HNS). Patient Health Information. Ear Plastic Surgery. © Copyright 2021. American Academy of Otolaryngology — Head and Neck Surgery. Accessed February 11, 2021. Available at URL address: https://www.enthealth.org/be_ent_smart/ear-plastic-surgery/
13. FACES: The National Craniofacial Association. Microtia. © FACES: The National Craniofacial Association. Not dated. Accessed February 11, 2021. Available at URL address: <https://www.faces-cranio.org/microtia>
14. Fioramonti P, Serratore F, Tarallo M, Ruggieri M, Ribuffo D. Otoplasty for prominent ears deformity. *Eur Rev Med Pharmacol Sci.* 2014;18(21):3156-65.

15. Gosain AK, Kumar A, Huang G. Prominent ears in children younger than 4 years of age: what is the appropriate timing for otoplasty? *Plast Reconstr Surg*. 2004 Oct;114(5):1042-54.
16. Haddad J, Dodhia SM. Congenital Malformations of the Ear. In: Kliegman RM, St Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM. *Nelson Textbook of Pediatrics*. 21st edition. Philadelphia PA: Elsevier Inc; 2020. Ch 66. Pgs 3411-3414.e1.
17. Haddad J. Congenital malformations. In: Kliegman RM, editor. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia, PA: Saunders; 2011. Chapter 630.
18. Haytoglu S, Haytoglu TG, Bayar Muluk N, Kuran G, Arikan OK. Comparison of two incisionless otoplasty techniques for prominent ears in children. *Int J Pediatr Otorhinolaryngol*. 2015 Apr;79(4):504-10.
19. Kelley P, Hollier L, Stal S. Otoplasty: evaluation, technique, and review. *J Craniofac Surg*. 2003 Sep;14(5):643-53.
20. Kelley PE, Scholes MA. Microtia and congenital aural atresia. *Otolaryngol Clin North Am*. 2007 Feb;40(1):61-80, vi.
21. Kim SY, Bothwell NE, Backous DD. The expanding role of the otolaryngologist in managing infants and children with hearing loss. *Otolaryngol Clin North Am*. 2002 Aug;35(4):699-710.
22. Leach, LL Jr, Biavati MJ. Ear reconstruction. In: *eMedicine specialties > otolaryngology and facial plastic surgery > reconstructive surgery*. Updated April 12, 2019. Accessed February 11, 2021. Available at URL address: <http://www.emedicine.com/ent/topic79.htm>
23. Lennon C, Chinnadurai S. Nonsurgical Management of Congenital Auricular Anomalies. *Facial Plast Surg Clin North Am*. 2018 Feb;26(1):1-8.
24. Liaw J, Patel VA, Carr MM. Congenital anomalies of the external ear. *Operative Techniques in Otolaryngology*. 2017 June; 28(2): 72-76.
25. Lin K, Marrinan MS, Shapiro WH, Kenna MA, Cohen NL. Combined microtia and aural atresia: issues in cochlear implantation. *Laryngoscope*. 2005 Jan;115(1):39-43.
26. McKinnon BJ, Jahrsdoerfer RA. Congenital auricular atresia: update on options for intervention and timing of repair. *Otolaryngol Clin North Am*. 2002 Aug;35(4):877-90.
27. Medicare Services Advisory Committee (MSAC); Doust J, Murray A-M, Vandervord J. Total ear reconstruction: MSAC application 1024 [assessment report]. Canberra, Australia: Commonwealth of Australia; 2000 Mar. Endorsed 2000 March 6 by the Commonwealth Minister for Health and Aged Care. Accessed February 11, 2021. Available at URL address: <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/1024-public>
28. Murakami CS, Quatela VC, Si K, Shvidler J. Reconstruction surgery of the ear: Microtia Reconstruction. In: Cummings CW, Flint PW, Haughey BH, Robbins KT, Thomas TR, Harker LA, et al., editors. *Otolaryngology: Head and Neck Surgery*. 5th ed. Copyright ©2010. Chapter 192.
29. National Institute for Health and Care Excellence. Incisionless Otoplasty (IPG422) March 28, 2012. Accessed February 11, 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg422>
30. Owsley TG. Otoplastic surgery for the protruding ear. *Atlas Oral Maxillofac Surg Clin North Am*. 2004 Mar;12(1):131-9.
31. Renner G, Lane RV. Auricular reconstruction: an update. *Curr Opin Otolaryngol Head Neck Surg*. 2004 Aug;12(4):277-80.

32. Schultz K, Guillen D, Maricevich RS. Newborn Ear Deformities: Early Recognition and Novel Nonoperative Techniques. *Semin Plast Surg.* 2017 Aug;31(3):141-145.
33. Strychowsky JE1, Moitri M, Gupta MK, Sommer DD. Incisionless otoplasty: a retrospective review and outcomes analysis. *Int J Pediatr Otorhinolaryngol.* 2013 Jul;77(7):1123-7.
34. Types of hearing loss. American Speech-Language-Hearing Association. Not dated. Accessed February 11, 2021. Available at URL address: <http://www.asha.org/public/hearing/Types-of-Hearing-Loss/>
35. Zim SA. Microtia reconstruction: an update [review]. *Curr Opin Otolaryngol Head Neck Surg.* 2003 Aug;11(4):275-81.

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.