



# Medical Coverage Policy

Effective Date..... 8/15/2021  
Next Review Date..... 8/15/2022  
Coverage Policy Number ..... 0403

## Male Sexual Dysfunction Treatment: Non-pharmacologic

### Table of Contents

Overview ..... 1  
 Coverage Policy..... 1  
 General Background.....2  
 Medicare Coverage Determinations ..... 13  
 Coding/Billing Information..... 13  
 References ..... 15

### Related Coverage Resources

- [Collagenase clostridium histolyticum](#)
- [Sildenafil \(Viagra®\)](#)
- [Vardenafil](#)
- [Avanafil](#)

#### **INSTRUCTIONS FOR USE**

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

### Overview

This Coverage Policy addresses devices and procedures used in the treatment of erectile dysfunction.

### Coverage Policy

**Coverage for male sexual dysfunction varies across plans. Refer to the customer's benefit plan document for coverage details.**

**If coverage is available for the treatment of male sexual dysfunction, the following conditions of coverage apply.**

#### **Vacuum Erection Device**

**A vacuum erection device is considered medically necessary for the treatment of erectile dysfunction.**

## **Penile Prosthesis**

**Surgical implantation of a penile prosthesis is considered medically necessary for the treatment of erectile dysfunction when ALL of the following criteria are met:**

- erectile dysfunction has persisted for at least six months
- a comprehensive history and physical exam, including appropriate laboratory testing, has been completed
- there is failure, contraindication or intolerance to conservative medical management including FDA pharmacological therapy (e.g., oral PDE5 inhibitors, intracavernosal injection, intraurethral medication) and/or a vacuum erection device

**Surgical reimplantation of a medically necessary penile prosthetic device, following the medically necessary removal of a penile prosthesis, is considered medically necessary when benefit coverage is available.**

**Removal of a penile prosthesis is considered medically necessary for ANY of the following indications:**

- infection
- mechanical failure
- urinary obstruction
- intractable pain

**A penile prosthesis for ANY other indication is considered not medically necessary.**

## **Not Covered Procedures**

**Each of the following procedures for the treatment of erectile dysfunction is considered experimental, investigational or unproven:**

- venous occlusive surgery (e.g., venous ligation)
- sural nerve grafting during radical retroperitoneal prostatectomy
- extracorporeal shock wave therapy (ESWT)

## **General Background**

Erectile dysfunction (ED) (i.e., impotence) is defined as the inability to achieve or maintain an erection sufficient for satisfactory sexual performance. ED usually has a physical cause in older men and is treatable at all ages. An age correlation exists for the prevalence of ED, however it is not an inevitable part of the aging process. Burnett and Ramasamy report the worldwide prevalence is 1%–10% for men younger than the age 40, up to 15% for men age 40–49, up to 30% for men age 50–59, up to 40% for men age 60–69, and 50%–100% for men between age 70–90. Additionally, it is estimated that in 1995, there were more than 152 million men worldwide who experienced ED. The 2025 prevalence projection was that approximately 322 million men would have ED. They report that the trend is maintained irrespective of racial/ethnic background or geographic region.

It was estimated that there were more than 152 million men worldwide who experienced ED in 1995, with a projection of the prevalence reaching approximately 322 million men having ED by 2025 (Aytac et al., 1999). This trend is maintained irrespective of racial/ethnic background or geographic region. (Santucci and Bandukwala, 2016; Rosen, et al., 2005; Fazio, et al., 2004; Morales, 2003).

There are multiple causes of organic ED including disease processes, trauma, drug and alcohol use/abuse, as well as smoking. ED may occur as a result of an underlying medical condition, such as diabetes, kidney disease, hormonal imbalance, multiple sclerosis, atherosclerosis, vascular disease or neurological disease. Injury to the penis, spinal cord, prostate, bladder, and pelvis may also cause ED due to damage to nerves smooth muscles,

arteries or fibrous tissue of the corpora cavernosa. Surgery, especially radical prostate or bladder surgery can injure the nerves and arteries near the penis resulting in ED. One of the side effects of medications, such as antihypertensive drugs, antihistamines, antidepressants, tranquilizers, histamine-receptor antagonists for treatment of gastric ulcers, opiates, and appetite suppressants is ED. Peyronie's disease, which causes scarring of the fibrous tissue of the penis, and priapism (i.e., persistent, abnormal erection of the penis) are also associated with ED. Other possible contributing factors of ED include smoking, which affects blood flow, and hormonal abnormalities. Psychological factors (e.g., stress, anxiety, depression, and low self-esteem) cause 10–20% of ED cases (Santucci and Bandukwala, 2016; McVary, 2007; Rosen, et al., 2005; Fazio, et al., 2004; Morales, 2003).

Men with symptoms of ED should have a complete medical, sexual, and psychosocial history taken along with a physical examination and selective laboratory testing. Patients should be informed that ED is a risk factor for underlying cardiovascular disease and other health conditions that may warrant further evaluation and treatment. The laboratory tests that are routinely performed as a part of an ED evaluation are serum testosterone, glucose/hemoglobin A1c, and in some cases serum lipids. No routine serum study is likely to alter ED management. However, serum studies may provide information on the etiology of ED and reveal the presence of additional conditions that require treatment. In addition, a prostate specific antigen (PSA) may be appropriate in some men with ED (American Urological Association [AUA], 2018). Vascular flow to the corpora cavernosa may be evaluated using a penile doppler examination. Color duplex ultrasonography, which measures cavernous artery diameter and pressures, may also be used to assess venous leakage. According to the American Association of Clinical Endocrinologists (AACE), "occasionally, the measurement of nocturnal penile tumescence and rigidity is useful, especially to distinguish between psychologic and organic erectile dysfunction" (AACE, 2003).

The method of treatment for ED is dependent upon the etiology of the condition. Psychologically-based ED, without organic cause (e.g., secondary to depression, anxiety, stress) may dissipate with psychotherapy and/or behavioral therapy. Organic ED can occur as a secondary condition to several diseases and/or their treatment. Treatment of underlying diseases such as diabetes mellitus, hypertension, heart disease and endocrine conditions (e.g., hypogonadism, hyperprolactinemia, and thyroid disorders), and cessation or modification of prescription medications (e.g., antihypertensives) may be indicated. Discontinuing alcohol consumption and illicit drug use, and/or making lifestyle modifications (e.g., avoiding smoking, maintaining ideal body weight and engaging in regular exercise) may reverse ED. Treatment of Peyronie's disease resulting in severe curvature may involve the concomitant use of incision/grafting and prosthesis insertion due to the significant incidence of erectile dysfunction following surgery on the penis for Peyronie's plaques (Taylor and Levine, 2007). There is some controversy regarding testosterone replacement therapy, which includes oral preparations, intramuscular injections, topical gels, and transdermal preparations. Topical gels are the most commonly prescribed forms of testosterone replacement (Brant, et al., 2007; McVary, 2007; Seftel, et al., 2004; Morales, 2003).

Treatment algorithms for erectile dysfunction (ED) involve using medical therapies such as phosphodiesterase type 5 (PDE5) inhibitors, intracavernosal injection therapy of vasoactive agents, as well as vacuum erection devices (Chung, 2019). Current practice guidelines suggest that the choice of PDE5 inhibitor should be based upon on the patient's preferences, including cost, ease of use, and adverse effects (AUA, 2018). Oral agents (e.g., PDE-5 inhibitors) are successful in 70–80% of men. With the availability of oral agents and minimally invasive options, surgical implantation typically occurs when these less invasive options are unavailable, unsuccessful or provide inadequate erectile function (Chung, 2019; Brant, et al., 2007; McVary, 2007; Sadeghi-Nejad, 2007; Jain and Terry, 2006; Carson, 2005; Fazio, et al., 2004; Morales, 2003).

In addition to the surgically implanted prostheses, other procedures may be recommended for ED that is refractory to medical therapy. Vascular surgical procedures include penile arterial bypass or revascularization and venous ligation for the treatment of vasculogenic ED. For those with ED unresponsive to nonsurgical treatments, vascular surgery may be the preferred treatment option that offers the possibility of spontaneous, unaided erections. The success rates for arterial revascularization are low but reasonable success rates may be achieved in young, nonsmoking, otherwise healthy men with recently acquired ED due to a focal arterial occlusion. Techniques to improve the veno-occlusive mechanism with ligation of the dorsal, cavernous, and crural veins have been largely abandoned in favor of medical therapies (PDE-5 inhibitors) (Khera, 2020). Sural nerve grafting has been proposed as a surgical intervention for ED that occurs in association with radical

prostatectomy. The Nesbit and Lue procedures are established for the correction of penile deformities caused by Peyronie's disease. Extracorporeal Shock Wave Therapy (ESWT) has also been proposed as a treatment for Peyronie's disease.

### **Non-pharmacologic Treatment of Erectile Dysfunction**

**Vacuum Erection Device:** An alternative treatment for erectile dysfunction is a vacuum erection device (VED). This device functions as an external aid; however, some users may find it difficult to use. The device causes an erection by creating a partial vacuum, drawing blood into the penis, engorging and expanding it. The device has three components: a plastic cylinder, in which the penis is placed; a pump that draws air out of the cylinder; and an elastic band that is placed around the base of the penis to maintain the erection when the cylinder is removed.

**Penile Prosthesis:** When nonsurgical therapies have proven ineffective, a penile prosthesis may be surgically implanted. Since surgery destroys the corpus cavernosum of the penis, this procedure precludes any future pharmacological treatment (Morales, 2003).

Complications of penile prostheses include erosion of the device, mechanical failure and the possibility of infection. Device extrusion, migration, urinary obstruction and prolonged or intractable pain are other potential risks. The average infection rate post-operatively ranges from 2–4% over a two year period, with most infections becoming evident during the first year. Some bacterial species can lie indolent for as long as two years before causing clinical signs of infection. Men with diabetes, spinal cord injuries or urinary tract infections have an increased risk of prosthesis-associated infections. If the infection cannot be successfully treated with antibiotics, it may be necessary to remove the prosthesis. Replacement with a new prosthesis should be delayed after removal of an infected prosthesis to allow adequate healing and eradication of the offending microorganism (Santucci and Bandukwala, 2016).

### **U.S. Food and Drug Administration (FDA)**

There are two types of mechanical devices for treatment of erectile dysfunction: external penile rigidity devices and penile rigidity implants. Both are regulated by the FDA. External penile rigidity devices are classified by the FDA as Class II medical devices and are exempt from the premarket notification requirements of the 510(k) process (FDA, 2004). Examples of these devices are the Rejoyn Vacuum Therapy System (Rejoyn Medical Systems, Nevada City, CA) and Osbon Erecaid™ Vacuum Therapy (Timm Medical Technologies, Dodge City, KS).

Penile rigidity implants are either noninflatable (i.e., semirigid rods) or inflatable. Noninflatable devices are classified by the FDA as Class II medical devices and consist of a pair of semi-rigid rods or cylinders that are surgically implanted in the corpora cavernosa. The purpose of the device is to provide adequate penile rigidity for intercourse. This classification includes the following designs (FDA, 2000):

- rod prosthesis: a flexible, solid cylinder of polymer material
- malleable prosthesis: a flexible polymer cylinder that incorporates an internal metal core
- single-hinged prosthesis: a highly flexible material that enables the user to position the penis downward for concealment
- multiple-hinged prosthesis: a series of hinged segments, encapsulated in a polymer sheath

The Spectra™ (AMS Men's Health/Boston Scientific) and the Genesis® Penile Prosthesis (Coloplast, Minneapolis, MN) are examples of rigid penile prostheses.

Inflatable devices are classified by the FDA as Class III medical devices and consist of paired cylinders, surgically implanted inside the penis, which can be expanded using pressurized fluid. Tubes connect the cylinders to a reservoir filled with radiopaque fluid implanted in the abdomen and a subcutaneous pump implanted in the scrotum. The user inflates the cylinders by pressing on the small pump, located under the skin in the scrotum (FDA, 2020). The AMS 700™ (Boston Scientific) and the Titan® (Coloplast, Minneapolis, MN) are examples of inflatable penile prostheses.

### **Literature Review**

Due to the nature of these devices, outcomes reported in studies evaluating their effectiveness are largely self-reported and subjective (e.g., patient satisfaction questionnaires). Objective outcome measures that have been reported in the medical literature include rate of mechanical failures and defects, and complications. Published evidence supports improved patient satisfaction with the use of penile implants when compared to sildenafil or intracavernous injections (Rajpurkar, et al., 2003); improved quality of life (Ferguson, et al., 2003); and improved erectile function (Mulhall, et al., 2003). Patient satisfaction has been reported to range from 71% to 91.2% with the use of implantable penile prostheses (Paranhos, et al., 2010; Knoll, et al., 2009; Israilov, et al., 2005; Minervini, et al., 2006; Zermann, et al., 2006; Ferguson, et al., 2003). Wilson et al. (2007) reported an estimated mechanical revision rate of 79.4% for device survival at 10 years compared to 71.2% at 15 years. The authors also noted with newer devices a 10-year mechanical survival and freedom from mechanical breakage increased to 88.6% and 97.9%, respectively. In general, the medical literature indicates these devices are safe and effective for the treatment of ED for a carefully selected subset of individuals whose condition is organic in nature and have failed more conservative treatment.

### **Vascular Surgery**

Patients who are considered for vascular surgical therapy typically have appropriate preoperative evaluation, which may include the combined injection and stimulation (CIS) test, dynamic infusion pharmacocavernosometry and cavernosography (DICC), duplex ultrasonography, and possibly arteriography. Penile arterial reconstructive surgery, also referred to as penile arterial bypass or revascularization, is one intervention that has the potential to cure patients with ED. During penile revascularization procedures, an arterial blockage is bypassed usually by anastomosing the inferior epigastric artery to the dorsal artery of the penis. Young men without other vascular risk factors (e.g., diabetes, high blood pressure, lipid disorders, cigarette smoking), who have ED due to pure artery blockage, are ideal candidates for this procedure. For posttraumatic arteriogenic ED in young patients, surgical penile revascularization has a 60–70% long-term success rate (Hatzimouratidis, et al., 2018).

### **Penile Vein Ligation**

Venous ligation is a surgical procedure used to treat veno-occlusive ED, or erectile failure caused by venous insufficiency. The procedure involves the removal or ligation of the veins leaving the corpora cavernosa. Penile vein ligation techniques range from dorsal and accessory vein ligation to complete ligation and excision of the dorsal, cavernous, and crural veins. Surgery of the penile venous system has been reported to have some efficacy in patients with venous leakage. However, the tests necessary to establish this diagnosis have been incompletely validated. Therefore, it is difficult to select patients who will have a predictably good outcome. In general, the long-term benefits of venous ligation surgery have been limited. Success rates within the first year range from 23–80% but consistently decrease on longer follow-up (Rao and Donatucci, 2001).

### **Literature Review**

Studies in the published peer-reviewed medical literature evaluating penile vein ligation for venogenic ED include case series primarily with small sample sizes. A larger series by Hsu et al. (2010) compared patients with veno-occlusive dysfunction who were treated with a venous stripping surgical method (n=178) with patients who were treated without this surgery (n=163). At an average follow-up of 7.7 years, there were no statistically significant differences in outcomes between the surgery and control groups as measured by the International Index of Erectile Function (IIEF-5) scores.

Cayan (2008) reported a series of 26 patients who underwent penile venous surgery with crural ligation for primary venous leakage. Postoperatively complete improvement in erectile function occurred in 11 men (42.3%), partial improvement occurred in eight (30.8%), and erectile function remained unchanged in seven (26.9%). Earlier case series have reported success rates of 31%–60% (Berardinucci, et al., 1996; Kim and McVary, 1995).

### **Sural Nerve Graft with Radical Prostatectomy**

Despite advances in radical prostatectomy procedures, ED remains a significant postoperative complication. When both neurovascular bundles are spared during radical retropubic prostatectomy (RRP), potency rates of up to 70% have been reported, but rates of 30–60% have been observed. For intentional resection of both neurovascular bundles, the return of erectile function is the exception (Kim, et al., 2001). Sural nerve grafting has been proposed as an intervention at the time of RRP to prevent ED associated with the procedure. In sural nerve

grafting, a portion of the sural nerve is harvested from one leg and then anastomosed to the divided ends of the cavernous nerves which are resected during a radical prostatectomy.

### **Literature Review**

There is limited data in the scientific peer-reviewed literature regarding the long-term outcomes of sural nerve grafting. An RCT by Davis et al. (2009) compared outcomes of patients who underwent unilateral nerve-sparing radical prostatectomy with a sural nerve graft (n=66) to those who had unilateral nerve-sparing radical prostatectomy alone (n=41). At 24-month follow-up, there was no significant difference in the return of erectile function for the sural nerve graft group versus the control group (p=0.962).

Nonrandomized controlled comparison studies and case series have also evaluated the safety and effectiveness of sural nerve grafting with radical prostatectomy (Siddiqui, et al., 2014; Satkunasivam, et al., 2009; Hanson, et al., 2008; Zorn, et al., 2008; Sim, et al., 2006; Porpiglia, et al., 2005; Kim, et al., 2001). Study populations have ranged from 28–40 with a follow-up range of 12–24 months. Small sample sizes have limited the generalizability of results. In addition, the results of these studies have not consistently shown a statistically significant difference in erectile function after sural nerve grafting.

### **Extracorporeal Shock Wave Therapy (ESWT)**

ESWT is a noninvasive treatment that involves delivery of low- or high-energy shock waves via a device to a specific site within the body. These pressure waves travel through fluid and soft tissue; their effects occur at sites where there is a change in impedance, such as the bone/soft-tissue interface.

### **Peyronie's Disease (PD)**

PD is a localized connective tissue disorder of unknown cause, and is characterized by the formation of inelastic fibrous plaques within the tunica albuginea or erectile tissue of the penis. For many patients, PD results in sexual problems due to the difficulty in attaining and/or maintaining erections. In a significant number of patients, the disorder improves or resolves spontaneously. Medical therapies, including antioxidants (such as vitamin E and potassium aminobenzoate) and corticosteroids injected directly into the plaque, lack adequate scientific support. Medical treatment options for PD typically include oral or intralesional drug therapy. Surgical management is considered for patients who have penile deformity compromising sexual function and whose PD has persisted for more than 12 months, and is refractory to medical therapy (Brant, et al., 2021).

ESWT has been investigated as a treatment option for PD. Various hypotheses about its mechanism of action exists, including direct damage to the plaque resulting in an inflammatory reaction with increased macrophage reaction leading to plaque lysis, improved vascularity resulting in plaque resorption, and the creation of contralateral scarring of the penis resulting in "false" straightening (Taylor and Levine, 2007). There is a lack of standardization regarding issues such as shockwave dosage, energy levels and number of sessions required for a therapeutic effect in patients with PD. Currently, the treatment of PD is not a U.S. Food and Drug Administration (FDA)-approved indication for ESWT.

### **Literature Review**

The use of ESWT for the treatment of PD has been examined in several RCTs, systematic review/meta-analysis and observational studies. Gao et al. (2016) conducted a meta-analysis of the evidence on ESWT for PD (n=6 studies/443 patients). Data was extracted from RCTS (n=3 studies/238 subjects), case-control studies (n=2 studies/111 subjects), and one cohort study (n=94 subjects). The primary outcomes were lessening of plaques and improvement of penile curvature. Secondary outcomes included relief and complete remission of pain, and improvement of sexual function. Follow-up ranged from four weeks to six months. Pooling data showed statistically significant pain relief and remission of pain (p<0.0001) after ESWT. A decrease of penile plaque was also observed (p=0.02). However, insignificant differences were found in improvement of penile curvature (p=0.06) and sexual function (p=0.18) between ESWT and placebo groups. A meta-analysis of RCTs only showed similar results for all parameters. Study limitations include the short-term follow-up in trials and the inclusion of low-quality evidence. The authors concluded that although the results of this meta-analysis suggests that ESWT may be an effective and relatively safe choice for PD patients with penile plaque and painful erection, the efficacy of ESWT could be very limited in patients with penile curvature and ED.

Chitale et al. (2010) randomized men with Peyronie's disease to receive ESWT (n=16) or sham (n=20). Inclusion requirements were stable penile deformity secondary to PD, recent onset of painless deformity of the penis on erection, and stable for > six months; pain and/or angulation of the penis on erection; difficult intercourse due to penile curvature, and partner dissatisfaction; a degree of ED (partial) associated with penile deformity; palpable plaque along the penis with penile deformity; aged > 18 years. The exclusion criteria were: congenital curvature of the penis; previous treatment for PD (surgical/medical); patients on warfarin and patients with total ED in need of therapy for ED. Primary outcome measures were the difference in the angle of deformity and the difference in IIEF score before and after treatment. Secondary outcome measures included the difference in VAS scores before and after treatment. At six months of follow-up there was no significant difference in the mean change between the control and intervention groups on any outcome measure. The study is limited by its small sample size. Results do not support the effectiveness of ESWT for PD.

Palmieri et al. (2009) conducted an RCT comparing ESWT as a treatment for PD with a duration of less than 12 months (n=50) to placebo (n=50). At 24 weeks, the mean visual analog scale (VAS) score was reported to be significantly lower in both groups compared to baseline. Mean plaque size and mean curvature degree were significantly higher in the placebo group compared to baseline and ESWT values. Study limitations include small sample size, restricted inclusion criteria, and short-term follow-up.

In general, the evidence evaluating the safety and efficacy of ESWT for PD consists of observational studies and case series with relatively small sample sizes (n=44–325) and short-term follow-up (di Mauro, et al., 2019; De Berardinis, et al., 2010; Poulakis, et al., 2006; Srirangam, et al., 2006; Strebel, et al., 2004; Hauck, et al., 2004). These studies have yielded inconsistent results.

### **Vasculogenic ED**

Low-intensity ESWT (LI-ESWT) has recently been investigated as a treatment for patients with mild to severe ED resulting from altered blood flow to the penis. Low-intensity extracorporeal shockwave therapy (Li-ESWT) is noninvasive and uses acoustic waves, which can pass through tissue and be focussed to target specific areas or organs to induce the desired effects (Young Academic Urologists Men's Health Group, 2017). LI-ESWT is thought to induce mechanical stress and cellular microtrauma which results in the development of new blood vessels in the treated tissue of the corpora cavernosa. LI-ESWT aims to restore the erectile mechanism in order to enable natural or spontaneous erections. Patients who might be offered LI-ESWT include those with vasculogenic ED amenable to microvascular surgery.

### **Literature Review**

Evidence in the published peer-reviewed medical literature evaluating ESWT for ED includes RCTs, cohort studies and systematic reviews/meta-analyses.

Vinay et al. (2021) conducted a double-blind, sham controlled, randomized controlled trial that assessed the effect of electromagnetic LI-ESWT on the erectile function of vascular phosphodiesterase type 5 inhibitor (PDE5I) refractory ED patients. Included patients were predominantly, middle-aged men with a high prevalence of multiple cardiovascular risk factors with a history of ED of more than six months refractory to PDE5I drugs. Patients (n=80) were randomized to LI-ESWT (n=40) or sham treatment (n=40). All patients in the active LI-ESWT group and 36 sham treated patients completed the study protocol (n=76). All patients in the study stopped using ED drugs during study protocol and follow-up, with a wash-out period of two months. The LI-ESWT group received 5000 shocks once a week for four weeks using the RENOVA® electromagnetic device (Direx Group, Wiesbaden, Germany). The sham group was treated with a device probe, identical to the treatment group, but without shockwaves. Outcomes measured the response to treatment at one, three and six months after the last session using the following questionnaires: index of erectile function–erectile function domain (IIEF-EF), erection hardness score (EHS) (> 2), sexual encounter profile (SEP2 and SEP3) and global assessment question (GAQ1). In each of the four LI-ESWT sessions and three post-treatment evaluations, patients were questioned about the use of ED drugs. At one month, there were not any significant differences between the groups in any of the EF questionnaire scores. At the three month follow-up, there was a significant difference in the median change of the IIEF-EF score for active and sham groups in favor of the LI-ESWT group (p=0.004). There were not any significant differences between groups regarding the percentage of men with an EHS > 2 or the number of positive answers to SEP2, SEP3 and GAQ1 during the three month evaluation. At the six month follow-up, there were significant differences between the groups in EHS > 2 and positive answers to GAQ-1 in favor of the

LI-ESWT group ( $p=0.028$  and  $p=0.011$ , respectively). SEP2 and SEP3 positive answers did not present significant differences between groups. No adverse events were observed during the study. Author noted limitations included: small patient population, penile hemodynamics were not measured to diagnose vasculogenic ED or to confirm the improvement of cavernous blood inflow or penile rigidity and the short-term follow-up. The authors concluded that the study showed that penile electromagnetic shockwave therapy may improve erectile function, to a modest extent, on certain patients that do not respond to PDE5I. However, randomized sham controlled trials with long-term follow-up is needed to compare different ED etiologies and protocol characteristics. No health disparities were identified by the investigators.

Ortac et al. (2021) reported on a prospective, single-blinded, placebo randomized controlled trial that compared outcomes in ED patients after ESWT and placebo treatment. Included patients were age 18–75 with a diagnosis of mild ED (IIEF-EF score = 17–25) which was confirmed by Penile Doppler ultrasonography (US) at least six months prior to study. The study was comprised of a four week washout phase, a four week treatment phase, and a 48-week follow-up. Patients ( $n=66$ ) were randomized to ESWT ( $n=44$ ) or placebo ( $n=22$ ). Mean age of ESWT and the placebo group was  $42.32 \pm 9.88$  and  $39.86 \pm 11.64$  ( $p=0.374$ ) respectively. The ESWT group received 3000 shock wave pulses once a week using the DUOLITH® SD1 shock wave generator. The placebo group also received treatment once a week, however no shocks were given during treatment. Outcomes measured changes from baseline in patient-reported outcomes of erectile function (International Index of Erectile Function domain scores [IIEF-EF]), as well as erection hardness and duration (Sexual Encounter Profile diary [SEP] and Global Assessment Questions [GAQ]). Safety was assessed throughout the study. Follow-up occurred at three months (V3), six months (V4) and 12 months (V5). Only treatment responders were evaluated at V4 and V5. At the three-month follow-up, mean IIEF-EF scores were significantly higher in ESWT patients than in placebo patients ( $p=0.003$ ) and IIEF-EF scores of ESWT patients remained high during the six month follow-up. A subgroup analyses indicated that the improvement of ED by ESWT detected at month three was independent of age, BMI, and ED duration. However, at six months follow-up, the beneficial effect of ESWT seemed to decline in patients with a BMI  $\geq 30$ , in patients older than 35 years and in patients with ED history  $\geq 12$  months. The percentage of patients reporting both successful penetration (SEP2) and intercourse (SEP3) in more than 50% of attempts was significantly higher in the ESWT patients compared to placebo patients ( $p=0.001$ ). One safety event (pain) and headache (1) occurred in the placebo group. A total of 12 adverse events were reported by ESWT patients ( $n=10$ ) which included headache (2), fatigue and nausea (1), nausea and headache (1), dysuria (1), fatigue and tiredness (2), headache and fatigue (1), and mild fever (1). No adverse event occurring at the sites of treatment was reported. Author noted limitations included that the study was single-blinded, (which might introduce a potential for investigator bias), small patient population and only responders were included in the open-label extension study (V4, V5). The study concluded that ESWT significantly improved the erectile function of relatively young patients with vasculogenic mild ED when compared to placebo for up to six months. However, additional research with large randomized controlled trials are needed to define subgroups of ED patients that will benefit the most from ESWT and to establish treatment protocols. No health disparities were identified by the investigators.

Shendy et al (2021) conducted a randomized controlled trial that assessed the effectiveness of low-intensity extracorporeal shockwave therapy (Li-ESWT) in the management of mixed vasculogenic and neurogenic erectile dysfunction in diabetic patients. Included patients were age 41–55 with a body mass index  $< 30$  kg/m<sup>2</sup> suffering from type-2 DM diabetic neuropathy as confirmed by nerve conduction study, and mild to moderate ED lasting for at least six months. All patients had controlled DM. Patients ( $n=42$ ) were randomly assigned to one of two groups: shock wave group ( $n=21$ ) treated with Li-ESWT plus pelvic floor exercises (PFE) and control group ( $n=21$ ) treated with pelvic floor muscle exercise and sham shock wave therapy. Li-ESWT sessions were delivered twice a week (3,000 shock waves) for three weeks, and repeated again after three weeks rest period. Pelvic floor exercises consisted of Kegel exercises three times daily for six weeks. The outcomes measured were improvement in International Index of Erectile Function (IIEF-5) and the penile Doppler parameters for the evaluation of penile perfusion of the two cavernous arteries. The assessment was done before and three months after treatment. IIEF-EF increased significantly in the study group ( $p<0.001$ ), but not in the control group ( $p=0.194$ ). Peak systolic velocity increased significantly in the two groups; however, the post-treatment peak systolic velocity was significantly higher in the study group compared to the control group ( $p<0.001$ , for both arteries). Author noted limitations were the small patient population and the short-term follow-up. The authors concluded that Li-ESWT appears to be an effective and safe noninvasive option for the treatment of diabetic



patients with erectile dysfunction. The combination of Li-ESWT and PFE can improve the vasculogenic and neurogenic elements of diabetic ED, however studies with long-term follow-up are needed to confirm the effectiveness of Li-ESWT to be approved for the treatment of DM-related ED. No health disparities were identified by the investigators.

Ladegaard et al (2021) investigated treating men with ED following robotic nerve-sparing radical prostatectomy (RARP) with Li-ESWT in a placebo randomized controlled trial. Men (n=38) from the Region of Southern Denmark (covering 1.2 million people) with ED for more than six months following nerve-sparing RP with an International Index of Erectile Function (IIEF-5) score < 22, no impaired EF prior to RP (no use of 5-PDEi, ICI or other erectogenic aid) and without active cancer or radiation therapy to the pelvic area were included. Participants were randomized into an active A group (n=20) and a placebo/sham B group (n=18). Each study arm had treatment once a week for five weeks using the handheld Duolith SD1 machine (Storz Medical, Switzerland) with 4,000 shock impulses. Outcomes were assessed by international validated questionnaires, Erection Hardness Score (EHS) and IIEF-5. Follow-up occurred at four weeks and 12 weeks following treatment. A significant increase was observed in EHS in group A at four and 12 weeks (p=0.033 and p=0.019, respectively). Group A had a significantly higher IIEF-5 score than group B at 12 weeks of follow-up. (p=0.026). However, the difference between groups in IIEF-5 score was not statistically significant at four weeks of follow-up. The study did not prohibit the use of erectile aids and the small patient population were noted as limitations by the authors. An additional limitation is that the population studied only included men from the Region of Southern Denmark and the results may not be applicable to other races or ethnic groups. The authors concluded that there was slight increase in mean IIEF-5 and EHS in men with ED following RARP treated with Li-ESWT. However, future research in Li-ESWT following RP is needed and should focus on initial timing of treatment, stratification of ED subgroups most susceptible to treatment, the efficiency of Li-ESWT in combination with other treatment modalities and defining the most optimal time for shockwave application.

Sramkova et al. (2020) evaluated the treatment outcome of Li-ESWT for ED in single-blind, randomized controlled trial. Included patients had mild to severe vasculogenic ED lasting for at least six months, were partial responders to PDE5i and had a stable partner with regular sexual activity at least twice a week. All patients were heterosexual and white European. The median patient age was 54 (range: 40–70). Patients (n=60) were randomized into two age-matched groups: Group A received treatment with PiezoWave2 unit (R.Wolf and Elvation Medical) and Group B received placebo. After a 4-week wash-out period, treatment consisted of four sessions with 6,000 shockwaves. Efficacy of treatment was measured according to the International Index of Erectile Function (IIEF-5), Erection Hardness Score (EHS), questions 2 and 3 of the Sexual Encounter Profile (SEP 2, SEP 3) and Global Assessment Question (GAQ) scores at baseline and four and 12 weeks following treatment. The patient's and partner's subjective satisfaction were also evaluated. The quality of erection according to the IIEF-5 and EHS at four and 12 weeks was clinically significant in favor of the treatment group (p=0.049 and p <0.001; p=0.030 and p<0.001, respectively). At 12 weeks, the GAQ, SEP 2, SEP 3, patient's satisfaction and partner's satisfaction reached clinical significance in favor of the treatment group (p<0.001, p=0.05, p<0.001, p <0.001, p <0.001); respectively). The author noted limitations of the study included the small patient population, short follow-up and the single blind study design. An additional limitation is that the population studied only included heterosexual white European men and the results may not be applicable to other races or ethnic groups. The authors concluded that Li-ESWT significantly improves erectile function. However, additional studies are needed, before considering this treatment as the new standard for the treatment of ED.

Baccaglini et al. (2019) conducted a randomized controlled trial that compared the early introduction of phosphodiesterase-5 inhibitor (PDE5i) with a combination of LiESWT and phosphodiesterase-5 inhibitor (PDE5i) in patients submitted to radical prostatectomy (RP). Patients aged ≤ 75 years presenting preoperatively with an IIEF-5 score > 18 and in a stable heterosexual relationship for at least three months were included. Patients underwent a bilateral nerve-sparing RP and agreed to discontinue PDE5i use at the end of the protocol for the last assessment of erectile rehabilitation. Patients (n=92) were randomized into the experimental group (n=46) or the control group (n=46). Both groups started tadalafil at a dose of 5 mg/day right after the removal of the transurethral catheter, and the experimental group received 2,400 shocks/session-week using Renova (DIREX Group). The control group did not receive placebo ESWT. The primary outcome measured the statistical difference in erectile function using the International Index of Erectile Function short form (IIEF-5), the end point was ≥ 4-point difference in favor of the experimental group at last follow-up. The authors noted that this specific cutoff has been proven to represent the minimal clinically important difference (MCID) in the EF domain of the

IIEF scale. The secondary outcome measured the rate of patients reaching orgasm and the erectile hardness score at last follow-up. Adverse events and continence status was also evaluated. At last follow-up (10 weeks), 77 patients were assessed and included in the final analysis, 36 in the experimental group 41 in the control group. A difference between groups was detected when accessing the final median IIEF-5 score (12.0 vs 10.0;  $p=0.006$ ). However, the primary clinical endpoint considering a difference  $\geq 4$ -point between the arms was not reached. When performing an exploratory analysis comparing the proportion of those individuals with an IIEF-5 score  $\geq 17$ , no difference between groups was noted. There were no significant differences noted between groups in postoperative complications or continence. The erectile hardness score and proportion of patients reaching orgasm were ruled out from the protocol because the investigators could not find any additional benefit for these tools after applying IIEF-5 in the protocol. Author acknowledged limitations included: the machine used Renova (DIREX Group) does not have the sham probe making the blinding process unfeasible; the early period after RP that assessed the IIEF-5 could have limited the spontaneous recovery of EF; PDE5i use was discontinued at the last session, which may have interfered in the penile vascular rehabilitation; some benefits of LiESWT could have disappeared after the washout period; the study protocol included only one session per week for the experimental group, possibly a more intensive application could achieved better results. Additional limitations were short term follow-up and small patient population. The population studied only included heterosexual men and the results may not be applicable to other races or ethnic groups. In conclusion, the study was unable to demonstrate clinical benefits on erectile function. The authors concluded that more studies are needed before any recommendation on this topic. No health disparities were identified by the investigators.

Dong et al. (2019) conducted a systematic review and meta-analysis of seven randomized controlled trials (RCT's) ( $n=522$  patients) to evaluate the efficacy of Li-ESWT by comparing the changes in the International Index of Erectile Function erectile function domain (IIEF-EF) and the Erection Hardness Score (EHS) to sham therapy in men with ED. The authors noted that men treated with Li-ESWT showed significant improvement in pooled mean IIEF-EF and EHS scores from baseline to follow-up compared to sham therapy ( $p<0.00001$  and  $p<0.00001$ , respectively). Study limitations included small sample size, short term follow up and no unified protocol using Li-ESWT in treating ED. The authors concluded that additional well designed RCTs are needed in the future.

A systematic review and meta-analysis of 10 RCT's ( $n=873$  patients) conducted by Sokolakis et al. (2019) concluded that when Li-ESWT was compared to sham, Li-ESWT significantly improved the IIEF-EF and EHS scores along with penile hemodynamics ( $p<0.0001$ ,  $p=0.0009$ ,  $p<0.00001$ , respectively). The mean follow-up was 5.12 months, with a range of 1–12 months. Acknowledged study limitations included the lack of consistency in outcome measures, specified dose intensities (low, medium, high ESWT) and short-term follow-up. The study concluded that larger multi-centric RCTs with longer than one year follow-up are needed.

Fojecki et al. (2017) assessed the outcome of linear low intensity-ESWT (LLi-ESWT) on ED at six and at 12 months from a randomized, double-blinded, sham-controlled trial. The advantage of linear ESWT compared with focused ESWT is that the usage of the linear probe delivers shock waves (SW) to a wider area of the corpora cavernosa, which limits movements of the probe during treatment. In this study, subjects with ED ( $n=126$ ) were randomized into two groups to compare LLi-ESWT to sham treatment for ED. Group A ( $n=63$ ) was assigned to a phase one sham for five weeks and phase two LLi-ESWT for five weeks. Group B ( $n=63$ ) was assigned to phase one and phase two of LLi-ESWT (10 weeks). The inclusion criteria were; age over 40 years, complaints of ED and in a stable relationship for greater than six months. The primary outcome measure was an increase of at least five points in the IIEF-EF ( $\Delta$ IIEF-EF score). The secondary outcome measure was an increase in the erectile hardness scale (EHS) score to at least a three in men with a score no higher than two at baseline. Patients ( $n=27$ ) were excluded from final analysis for not returning questionnaires ( $n=23$ ) or dropping out ( $n=4$ ). Linear regression of the  $\Delta$ IIEF-EF score from baseline to 12 months included 95 patients ( $n=95$ ). The success rate based on the main outcome parameter ( $\Delta$ IIEF-EF score  $\geq 5$ ) was 54% in group A vs. 47% in group B ( $p=0.28$ ). Improvements based on changes in the EHS score in groups A and B were 34% and 24%, respectively ( $p=0.82$ ). The differences between groups were not clinically significant. The limitations of the study included the lack of dosage related studies and the study design. Only the short-term effects of sham treatment were assessed, because all participants received active treatment in the second phase. The authors concluded that exposure to two cycles of linear Li-ESWT for ED is not superior to one cycle at six and at 12-month follow-ups and further studies are needed.

Kalyvianakis and Hatzichristou (2017) reported their results of a double-blinded, randomized, sham-controlled trial that assessed the effectiveness of low-intensity extracorporeal shockwave therapy (Li-ESWT) in patients with vasculogenic erectile dysfunction. Patients met inclusion criteria if they were at least 18 years old, had vasculogenic ED for at least six months, and were partial responders to PDE5 inhibitors. Following a four week washout period, the baseline International Index of Erectile Function erectile function domain (IIEF-EF) score had to be at least six (mild to moderate ED) to 21 (moderate and severe ED). Patients (n=46) were randomly assigned to receive Li-ESWT (n=30) or a sham procedure (n=16). All patients underwent penile triplex ultrasonography by the same investigator immediately before and three months after treatment. The primary outcome measured was the changes in peak systolic velocity and resistance index as measured by triplex ultrasonography at baseline and three months after treatment. Secondary outcomes measured were changes in the IIEF-EF score from baseline to one, three, six, nine, and 12 months after treatment and the percentage of patients reaching a minimal clinically important difference during the same period for the two groups. IIEF-EF minimal clinically important differences for the active vs sham group were observed for 56.7% vs 12.5% (p=0.005) at one month, 56.7% vs 12.5% (p=0.003) at three months, 63.3% vs 18.8% (p=0.006) at six months, 66.7% vs 31.3% (p=0.022) at nine months, and 75% vs 25% (p=0.008) at 12 months. Mean peak systolic velocity increased by 4.5 and 0.6 cm/s in the Li-ESWT and sham groups, respectively (p<0.001). Limitations included the small sample and strict inclusion criteria that do not reflect everyday clinical practice.

Clavijo et al. (2017) conducted a systematic review and meta-analysis of the evidence (n=7 RCTs/602 subjects) evaluating the efficacy of Li-ESWT for ED. RCTs were eligible for inclusion if they were published in the peer-reviewed literature and assessed erectile function outcomes using the IIEF-EF score. Estimates were pooled using random-effects meta-analysis. The average follow-up period was 19.8 weeks. A statistically significant improvement in pooled change in IIEF-EF score was found from baseline to follow-up in men undergoing Li-ESWT compared to those undergoing sham therapy (p<0.0001; between-group difference, p=0.047). Limitations of the review include the small sample size and short follow-up period in individual studies. It was also noted that data on PDE5i use during ESWT treatment was available in only five of seven studies. Although these results suggest that ESWT is more effective than sham therapy for ED, additional well-designed RCTs comparing ESWT to standard ED treatments are needed to establish the role in the treatment algorithm for ED.

Another systematic review and meta-analysis (n=14 studies/833 subjects) performed by Lu et al. (2017) evaluated the Li-ESWT for patients with ED. The evidence reviewed included RCTs (n=7 studies) which were used for meta-analysis and cohort studies (n=7 studies). No limitation was placed on the severity of ED or use of PDE-5i during treatment with Li-ESWT in studies. Follow-up occurred up to 12 months. The overall meta-analysis of the data showed that Li-ESWT improved the IIEF significantly overall in the treatment groups (p<0.0001). Different Li-ESWT setup parameters (energy flux density [EFD], number of pulses), and different treatment protocols, including treatment frequency and length of course, resulted in differences in reported efficacy. Results showed that studies using the highest EFD (> 0.2 mJ/mm<sup>2</sup>) reported significantly increased IIEFs (p<0.0001). Sub-group analysis (n=3 RCTs) showed that the IIEF of patients with mild ED increased significantly after Li-ESWT (p<0.0001), which did not occur for patients in the severe and moderate categories (p=0.30 and p=0.49, respectively). No adverse effects were reported. Acknowledged limitations of the meta-analysis were the inclusion of lower level evidence (i.e., cohort studies) and individual study deficiencies such as missing information as to details of randomization and treatment protocol.

Olsen et al. (2014) conducted a prospective, randomized blinded, placebo-controlled study (n=112) comparing LI-ESWT to sham treatment for ED. Both participants and physicians were blinded to treatment allocation. A total of 105 men with organic ED for at least six months who had responded to PDE-5 inhibitors were enrolled in this study. Additional inclusion requirements were an Erection Hardness Score (EHS) < 2 and an Index of Erectile Function (IIEF-15) score < 20. Exclusion criteria were psychogenic ED, neurological pathology and pelvic radiation or recovery from cancer within the past five years. The participants were assessed by EHS and IIEF-erectile function domain at baseline and at five weeks. Men in the placebo group (n=54) were offered LI-ESWT 10 weeks after study. The blinded part of the study was terminated at this point. The active placebo group (n=52) was assessed at five, 12 and 24 weeks after treatment. At five weeks of follow-up 29 men (57%) in the active group had an EHS of 3–4, which made it possible for them to have full sexual intercourse; three men (6%) had an EHS of 1–2, and 19 (37%) showed no change in ED. In the placebo group, five men (9%) had an EHS of 3–4, seven (13%) an EHS of 1–2 and 42 (78%) had experienced no change. The difference between the two groups was statistically significant at the EHS levels 0 and 3–4 (p=0.0001). The EHS response rate was 80% at week 12

and 70% at week 24 in the active group. In the active placebo group, the EHS response rate was 85% at week 12 and 75% at week 24. Between weeks 12 and 24, the number of men who achieved improved EHS scores (3–4) decreased in both groups. There was no significant difference between the two groups in terms of IIEF-erectile function domain after week five. This study is limited by its short-term follow-up and unblinding after five weeks. Study results suggest a short-term treatment effect however, this effect was not maintained throughout the follow-up period.

Vardi et al. (2012) performed a randomized, double-blind, sham controlled study (n=77) of men with organic ED who were PDE-5 inhibitor responders. After a one-month PDE-5 inhibitor washout period, 67 men were randomized in a 2:1 ratio to receive 12 sessions of Li-ESWT (n=40) or sham therapy (n=20). The primary outcome was erectile function measured by the IIEF-EF, with treatment success defined as a 5-point or greater score improvement. Secondary outcomes included an increase in EHS and an improvement in penile blood flow. The 9-week treatment period was comprised of two treatment sessions per week for three weeks that were repeated after a three-week no treatment interval. Follow-up occurred one month after the final treatment session at which time erectile function and penile hemodynamics were reassessed while the men were still not taking PDE-5 inhibitors. At 13 weeks of follow-up, men in the treated group had a 5-point or greater increase in IIEF-EF than those in the sham group (p=0.0001). The treated men had significantly improved mean scores in the IIEF subcategories of Sexual Desire (p=0.0348) and Overall Satisfaction (p=0.0054). Penile hemodynamics improved significantly in the treated group (p=0.0001). Study limitations are short-term follow-up and small sample size. Additional data are needed to confirm the efficacy suggested by these results.

Although preliminary results appear promising, additional well-designed studies with long-term follow-up are needed to establish safety and effectiveness of ESWT for the treatment of ED.

#### **Professional Societies/Organizations**

**American Academy of Family Physicians (AAFP):** According to the AAFP, oral phosphodiesterase-5 inhibitors are the first-line treatments for ED. Second-line treatments include alprostadil, vacuum devices and when all other options are ineffective surgically implanted penile prostheses are an option (Rew et al., 2016).

**American Association of Clinical Endocrinologists (AACE) and American College of Endocrinology (ACE):** AACE and ACE issued a guideline on the evaluation and treatment of male sexual dysfunction. The guideline supports the use of vacuum pump devices and internal implanted penile prostheses in the treatment of ED (AACE, 2003).

**American Urological Association (AUA):** In 2018, the AUA published revised guidelines on erectile dysfunction. According to the guidelines, men may choose to begin with the least invasive option, however any type of treatment as an initial treatment is a valid choice. The clinician is responsible to ensure that the man and his partner fully understand the benefits and risks/burdens associated with the choice and be informed of all the treatments (e.g., vacuum erection device, penile prosthesis) that are not contraindicated for the patient. The AUA also recommended against penile venous surgery and considered ESWT investigational.

**National Comprehensive Cancer Network® (NCCN):** According to the 2021 NCCN guidelines for prostate cancer, recovery of erectile function is directly related to factors such as age at radical prostatectomy, preoperative erectile function and the degree of preservation of the cavernous nerves. Replacement of resected nerves with nerve grafts has not been shown to be beneficial.

#### **Use Outside of the US**

The Canada Urology Association (CUA) recommended a stepwise progression from oral agents through second- and third-line therapies. After reversible causes of ED are ruled out, the recommended first line therapy is a trial of oral medication. If oral treatment fails or if greater insight into the etiology is desired, then specialized testing and referral are done. The second-line therapies, although more invasive than oral agents, are generally well-tolerated and effective and include injection therapy, urethral therapy and vacuum erection device. Penile implant surgery remains the third-line therapy and is an important option for men refractory to medical management, offering effective and durable ED treatment outcomes (Bella, et al., 2015).

The European Association of Urology (EAU) guidelines on male sexual dysfunction states that if drug treatment fails, patients should be offered an alternative therapy such as intracavernosal injection therapy or the use of a vacuum erection device. The surgical implantation of a penile prosthesis may be considered in patients who do not respond to pharmacotherapy or who prefer a permanent solution to their problem. In regards to shockwave therapy and erectile dysfunction, the current data is still limited and clear recommendations cannot be given (Hatzimouratidis, et al., 2018).

In addition, the EAU guidelines on penile curvature state that the mechanism of action involved in extracorporeal shock wave lithotripsy is unclear. According to the EAU, extracorporeal shock wave treatment may only be used to treat penile pain (Hatzimouratidis, et al., 2018). However, the cited evidence supporting this statement consists of a single placebo-controlled RCT (Palmieri et al., 2009) described previously.

According to the National Institute for Health and Care Excellence (NICE), (2003), “current evidence on the safety of extracorporeal shockwave therapy (ESWT) for Peyronie’s disease appears adequate. However, the evidence on the efficacy does not appear adequate to support the use of this procedure without special arrangements for consent and for audit or research.”

## Medicare Coverage Determinations

	Contractor	Policy Name/Number	Revision Effective Date
NCD	National	Diagnosis and Treatment of Impotence (230.4)	This is a longstanding national coverage determination.
LCD	CGS Administrators, LLC and Noridian Healthcare Solutions, LLC	Vacuum Erection Devices (VED) (L34824)	1/1/2020

Note: Please review the current Medicare Policy for the most up-to-date information.

## Coding/Billing Information

- Note:** 1) This list of codes may not be all-inclusive.  
 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

### Vacuum Erection Device

**Considered medically necessary when criteria in the applicable policy statements listed above are met and only when benefit coverage is available for the specific item:**

HCPCS Codes	Description
L7900	Male vacuum erection system
L7902	Tension ring, for vacuum erection device, any type, replacement only, each

### Penile Prosthesis

**Considered medically necessary when criteria in the applicable policy statements listed above are met and only when benefit coverage is available for the specific service/item:**

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
54400	Insertion of penile prosthesis; non-inflatable (semi-rigid)
54401	Insertion of penile prosthesis; inflatable (self-contained)
54405	Insertion of multi-component, inflatable penile prosthesis, including placement of pump, cylinders, and reservoir

<b>HCPCS</b> <b>Codes</b>	<b>Description</b>
C1813	Prosthesis, penile, inflatable
C2622	Prosthesis, penile, non-inflatable

### **Penile Prosthesis Removal**

**Considered medically necessary when criteria in the applicable policy statements listed above are met:**

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
54406	Removal of all components of a multi-component, inflatable penile prosthesis without replacement of prosthesis
54408	Repair of component(s) of a multi-component, inflatable penile prosthesis
54410	Removal and replacement of all component(s) of a multi-component, inflatable penile prosthesis at the same operative session
54411	Removal and replacement of all components of a multi-component inflatable penile prosthesis through an infected field at the same operative session, including irrigation and debridement of infected tissue
54415	Removal of non-inflatable (semi-rigid) or inflatable (self-contained) penile prosthesis, without replacement of prosthesis
54416	Removal and replacement of non-inflatable (semi-rigid) or inflatable (self-contained) penile prosthesis at the same operative session
54417	Removal and replacement of non-inflatable (semi-rigid) or inflatable (self-contained) penile prosthesis through an infected field at the same operative session, including irrigation and debridement of infected tissue

**Considered Experimental/Investigational/Unproven:**

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
37790	Penile venous occlusive procedure
55899 <sup>†</sup>	Unlisted procedure, male genital system
64999 <sup>††</sup>	Unlisted procedure, nervous system

**<sup>†</sup>Note:** Considered Experimental/Investigational/Unproven when used to report extracorporeal shock wave therapy for the treatment of erectile dysfunction.

**<sup>††</sup>Note:** Considered Experimental/Investigational/Unproven when used to report sural nerve grafting during radical retroperitoneal prostatectomy.

**\*Current Procedural Terminology (CPT®) ©2020 American Medical Association: Chicago, IL.**

## References

1. Adeldaeim HM, Abouyousif T, Gebaly OE, Assem A, Wahab MMA, Rashad H, Sakr M, Zahran AR. Prognostic Indicators for Successful Low-intensity Extracorporeal Shock Wave Therapy Treatment of Erectile Dysfunction. *Urology*. 2021 Mar;149:133-139.
2. Aditya I, Kwong JCC, Krakowsky Y, Grober ED. Non-conventional therapies for Peyronie's disease: what is the evidence for efficacy? *Transl Androl Urol*. 2020 Mar;9(Suppl 2):S295-S302.
3. American Association of Clinical Endocrinologists (AACE) Male Sexual Dysfunction Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the evaluation and treatment of male sexual dysfunction: a couple's problem—2003 update. *Endocr Pract*. 2003 Jan-Feb;9(1):77-95.
4. American Urological Association (AUA). Guideline on the Management of Erectile Dysfunction: Diagnosis and Treatment Recommendations. 2018. Accessed June 28, 2021. Available at URL address: [http://www.auanet.org/guidelines/erectile-dysfunction-uaa-guideline-\(2018\)](http://www.auanet.org/guidelines/erectile-dysfunction-uaa-guideline-(2018))
5. Baccaglini W, Pazeto CL, Corrêa Barros EA, Timóteo F, Monteiro L, Saad Rached RY, Navas A, Glina S. The Role of the Low-Intensity Extracorporeal Shockwave Therapy on Penile Rehabilitation After Radical Prostatectomy: A Randomized Clinical Trial. *J Sex Med*. 2020 Apr;17(4):688-694.
6. Bella, AJ, Lee JC, Carrier S, Bénard F, Brock GB. 2015 CUA Practice guidelines for erectile dysfunction. *Can Urol Assoc J* 2015;9(1-2):23-9.
7. Berardinucci D, Morales A, Heaton JP, Fenemore J, Bloom S. Surgical treatment of penile veno-occlusive dysfunction: is it justified? *Urology*. 1996;47(1):88-92.
8. Brant WO, Bella AJ, Lue TF. Peyronie's disease: Diagnosis and medical management. In: UpToDate, O'Leary MP, Givens J (Eds). Last updated January 29, 2021. UpToDate, Waltham, MA. Accessed June 25, 2021.
9. Brant WO, Bella AJ, Lue TF. Treatment options for erectile dysfunction. *Endocrinol Metab Clin North Am*. 2007 Jun;36(2):465-79.
10. Brunckhorst O, Wells L, Teeling F, Muir G, Muneer A, Ahmed K. A systematic review of the long-term efficacy of low-intensity shockwave therapy for vasculogenic erectile dysfunction. *Int Urol Nephrol*. 2019 May;51(5):773-781.
11. Burnett AL, Ramasamy R. Evaluation and Management of Erectile Dysfunction. In: Partin AW, Dmochowski RR, Kavoussi LR, Peters CA. *Campbell-Walsh-Wein Urology*. 12th ed. Philadelphia, PA: Elsevier; 2021. Ch 69. pgs 1513-1538.e8.
12. Campbell JD, Trock BJ, Oppenheim AR, Anusionwu I, Gor RA, Burnett AL. Meta-analysis of randomized controlled trials that assess the efficacy of low-intensity shockwave therapy for the treatment of erectile dysfunction. *Ther Adv Urol*. 2019 Mar 29;11:1756287219838364.
13. Carson CC. Penile prosthesis implantation: surgical implants in the era of oral medication. *Urol Clin North Am*. 2005;32(4):503-vii.
14. Cayan S. Primary penile venous leakage surgery with crural ligation in men with erectile dysfunction. *J Urol*. 2008 Sep;180(3):1056-9.

15. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed June 28, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx>
16. Centers for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed June 28, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/ncd-alphabetical-index.aspx>
17. Chitale S, Morse M, Swift L, Sethia K. Limited shock wave therapy vs sham treatment in men with Peyronie's disease: results of a prospective randomized controlled double-blind trial. *BJU Int*. 2010 Nov;106(9):1352-6.
18. Chung E. A Review of Current and Emerging Therapeutic Options for Erectile Dysfunction. *Med Sci (Basel)*. 2019;7(9):91.
19. Clavijo RI, Kohn TP, Kohn JR, Ramasamy R. Effects of Low-Intensity Extracorporeal Shockwave Therapy on Erectile Dysfunction: A Systematic Review and Meta-Analysis. *J Sex Med*. 2017 Jan;14(1):27-35.
20. Davis JW, Chang DW, Chevray P, Wang R, Shen Y, Wen S, et al. Randomized phase II trial evaluation of erectile function after attempted unilateral cavernous nerve-sparing retropubic radical prostatectomy with versus without unilateral sural nerve grafting for clinically localized prostate cancer. *Eur Urol*. 2009 May;55(5):1135-43.
21. De Berardinis E, Busetto GM, Antonini G, Giovannone R, Gentile V. Extracorporeal shock wave therapy in the treatment of Peyronie's disease: long-term results. *Arch Ital Urol Androl*. 2010 Jun;82(2):128-33.
22. Di Mauro M, Russo GI, Della Camera PA, Di Maida F, Cito G, Mondaini N, et al. Extracorporeal Shock Wave Therapy in Peyronie's Disease: Clinical Efficacy and Safety from a Single-Arm Observational Study. *World J Mens Health*. 2019 Sep;37(3):339-346.
23. Dong L, Chang D, Zhang X, Li J, Yang F, Tan K, et al. Effect of Low-Intensity Extracorporeal Shock Wave on the Treatment of Erectile Dysfunction: A Systematic Review and Meta-Analysis. *Am J Mens Health*. 2019 Mar-Apr;13(2).
24. Fazio L, Brock G. Erectile dysfunction: management update. *CMAJ*. 2004 Apr 27;170(9):1429-37.
25. Ferguson KH, Cespedes RD. Prospective long-term results and quality-of-life assessment after Dura-II penile prosthesis placement. *Urology*. 2003 Feb;61(2):437-41.
26. Fojecki GL, Tiessen S and Osther PJ: Effect of low-energy linear shockwave therapy on erectile dysfunction-a double-blinded, sham-controlled, randomized clinical trial. *J Sex Med* 2017; 14: 106.
27. Fojecki GL, Tiessen S, Osther PJ. Extracorporeal shock wave therapy (ESWT) in urology: a systematic review of outcome in Peyronie's disease, erectile dysfunction and chronic pelvic pain. *World J Urol*. 2016
28. Gao L, Qian S, Tang Z, Li J, Yuan J. A meta-analysis of extracorporeal shock wave therapy for Peyronie's disease. *Int J Impot Res*. 2016 Jun 2.
29. Gruenwald I, Appel B, Kitrey ND, Vardi Y. Shockwave treatment of erectile dysfunction. *Ther Adv Urol*. 2013 Apr;5(2):95-9.
30. Hanson GR, Borden LS Jr, Backous DD, Bayles SW, Corman JM. Erectile function following unilateral cavernosal nerve replacement. *Can J Urol*. 2008 Apr;15(2):3990-3.



31. Hatzichristodoulou G, Meisner C, Gschwend JE, Stenzl A, Lahme S. Extracorporeal shock wave therapy in Peyronie's disease: results of a placebo-controlled, prospective, randomized, single-blind study. *J Sex Med.* 2013 Nov;10(11):2815-21.
32. Hatzimouratidis K, Giuliano F, Moncada I, Muneer A, Salonia A, et al. EAU Guidelines on Male Sexual Dysfunction. 2018. Accessed on June 25, 2021. Available at URL address: <https://uroweb.org/guideline/male-sexual-dysfunction/>
33. Hauck EW, Hauptmann A, Bschiepfer T, Schmelz HU, Altinkilic BM, Weidner W. Questionable efficacy of extracorporeal shock wave therapy for Peyronie's disease: results of a prospective approach. *J Urol.* 2004 Jan;171(1):296-9.
34. Hsu GL, Chen HS, Hsieh CH, Lee WY, Chen KL, Chang CH. Clinical experience of a refined penile venous stripping surgery procedure for patients with erectile dysfunction: is it a viable option? *J Androl.* 2010 May-Jun;31(3):271-80.
35. Israilov S, Shmuely J, Niv E, Engelstein D, Livne P, Boniel J. Evaluation of a progressive treatment program for erectile dysfunction in patients with diabetes mellitus. *Int J Impot Res.* 2005 Sep-Oct;17(5):431-6.
36. Jain S, Terry TR. Penile Prosthetic Surgery and its Role in the Treatment of End-Stage Erectile Dysfunction – An Update. *Ann R Coll Surg Engl.* 2006 Jul; 88(4): 343–348.
37. Kalyvianakis D, Hatzichristou D. Low-Intensity Shockwave Therapy Improves Hemodynamic Parameters in Patients With Vasculogenic Erectile Dysfunction: A Triplex Ultrasonography-Based Sham-Controlled Trial. *J Sex Med.* 2017 Jul;14(7):891-897.
38. Kendirci M, Hellstrom WJ. Critical analysis of surgery for Peyronie's disease. *Curr Opin Urol.* 2004 Nov;14(6):381-8.
39. Khera, M. Treatment of male sexual dysfunction. Last updated June 23, 2020. In: UpToDate, Snyder PJ, O'Leary MP, Martin KA (Eds), UpToDate, Waltham, MA. Accessed on June 25, 2021
40. Kim ED, McVary KT. Long-term results with penile vein ligation for venogenic impotence. *J Urol.* 1995 Mar;153(3 Pt 1):655-8.
41. Kim ED, Nath R, Slawin KM, Kadmon D, Miles BJ, Scardino PT. Bilateral nerve grafting during radical retropubic prostatectomy: extended follow-up. *Urology.* 2001 Dec;58(6):983-7.
42. Kim ED, Scardino PT, Kadmon D, Slawin K, Nath RK. Interposition sural nerve grafting during radical retropubic prostatectomy. *Urology.* 2001 Feb;57(2):211-6.
43. Knoll LD, Henry G, Culkin D, Ohl DA, Otheguy J, Shabsigh R, Wilson SK, Delk II J. Physician and patient satisfaction with the new AMS 700 momentary squeeze inflatable penile prosthesis. *J Sex Med.* 2009 Jun;6(6):1773-1778.
44. Ladegaard PBJ, Mortensen J, Skov-Jepesen SM, Lund L. Erectile Dysfunction A Prospective Randomized Placebo-Controlled Study Evaluating the Effect of Low-Intensity Extracorporeal Shockwave Therapy (LI-ESWT) in Men With Erectile Dysfunction Following Radical Prostatectomy. *Sex Med.* 2021 Jun;9(3).
45. Lei Q, Wang D, Liu C, Ji Z, Yan S. Comparison of the efficacy and safety of low-intensity extracorporeal shock wave therapy versus on-demand sildenafil for erectile dysfunction. *Transl Androl Urol.* 2021 Feb;10(2):860-868

46. Lu Z, Lin G, Reed-Maldonado A, Wang C, Lee YC, Lue TF. Low-intensity Extracorporeal Shock Wave Treatment Improves Erectile Function: A Systematic Review and Meta-analysis. *Eur Urol*. 2017 Feb;71(2):223-233.
47. Man L, Li G. Low-intensity extracorporeal shock wave therapy for erectile dysfunction: A systematic review and meta-analysis. *Urology*. 2018 Sep;119:97-103.
48. McVary KT. Clinical practice. Erectile dysfunction. *N Engl J Med*. 2007 Dec 13;357(24):2472-81.
49. Minervini A, Ralph DJ, Pryor JP. Outcome of penile prosthesis implantation for treating erectile dysfunction: experience with 504 procedures. *BJU Int*. 2006 Jan;97(1):129-33.
50. Morales A. Erectile dysfunction: an overview. *Clin Geriatr Med*. 2003 Aug;19(3):529-38.
51. Mulhall JP, Ahmed A, Branch J, Parker M. Serial assessment of efficacy and satisfaction profiles following penile prosthesis surgery. *J Urol*. 2003 Apr;169(4):1429-33.
52. Musa ZS, El-Assmy A, Shokry AM, Shokeir AA, Zween T, Al-Kenawy MR. Long-term effectiveness and predictors of success of low-intensity shockwave therapy in phosphodiesterase type 5 inhibitors non-responders. *Arab J Urol*. 2019 Nov 11;18(1):54-58.
53. National Comprehensive Cancer Network® (NCCN). NCCN GUIDELINES™ Clinical Guidelines in Oncology™. Prostate Cancer. Version 2.2021. © National Comprehensive Cancer Network, Inc. 2021, All Rights Reserved. Accessed June 25, 2021. Available at URL address: [https://www.nccn.org/professionals/physician\\_gls/pdf/prostate.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf)
54. National Health Services (NHS). Erectile dysfunction (impotence) – Causes. Last updated 2020 March 12. Accessed June 28, 2021. Available at URL address: <https://www.nhsinform.scot/illnesses-and-conditions/sexual-and-reproductive/erectile-dysfunction-impotence#causes-of-erectile-dysfunction>
55. National Institute for Clinical Excellence (NICE). Interventional procedures guidance [IPG29]. Extracorporeal shockwave therapy for Peyronie's disease. March 2003. Accessed June 25, 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg29/chapter/1-guidance>
56. Olsen AB, Persiani M, Boie S, Hanna M, Lund L. Can low-intensity extracorporeal shockwave therapy improve erectile dysfunction? A prospective, randomized, double-blind, placebo-controlled study. *Scand J Urol*. 2014 Dec 3:1-5.
57. Ortac M, Özmez A, Cilesiz NC, Demirelli E, Kadioğlu A. The Impact of Extracorporeal Shock Wave Therapy (ESWT) for the Treatment of Young Patients with Vasculogenic Mild-Erectile Dysfunction (ED): A Prospective Randomized Single-Blind, Sham Controlled Study. *Andrology*. 2021 Mar 29.
58. Palmieri A, Arcaniolo D, Palumbo F, Verze P, Liguori G, Mondaini N, Falcone M, Scropo FI, Salonia A, Cai T; SIA-Low intensity shock wave for Erectile Dysfunction (LED) Study Group. Low intensity shockwave therapy in combination with phosphodiesterase-5 inhibitors is an effective and safe treatment option in patients with vasculogenic ED who are PDE5i non-responders: a multicenter single-arm clinical trial. *Int J Impot Res*. 2020 Jul 18.
59. Palmieri A, Imbimbo C, Longo N, Fusco F, Verze P, Mangiapia F, et al. A first prospective, randomized, double-blind, placebo-controlled clinical trial evaluating extracorporeal shock wave therapy for the treatment of Peyronie's disease. *Eur Urol*. 2009 Aug;56(2):363-9.
60. Paranhos M, Andrade E, Antunes AA, Barbieri AL, Claro JA, Srougi M. Penile prosthesis implantation in an academic institution in Latin America. *Int Braz J Urol*. 2010 Sep-Oct;36(5):591-601.

61. Porpiglia F, Ragni F, Terrone C, Renard J, Musso F, Grande S, et al. Is laparoscopic unilateral sural nerve grafting during radical prostatectomy effective in retaining sexual potency? *BJU Int.* 2005 Jun;95(9):1267-71.
62. Poulakis V, Skriapas K, de Vries R, Dillenburg W, Ferakis N, Witzsch U, et al. Extracorporeal shockwave therapy for Peyronie's disease: an alternative treatment? *Asian J Androl.* 2006 May;8(3):361-6.
63. Rao DS, Donatucci CF. Vasculogenic impotence. Arterial and venous surgery. *Urol Clin North Am.* 2001 May;28(2):309-19.
64. Rajpurkar A, Dhabuwala CB. Comparison of satisfaction rates and erectile function in patients treated with sildenafil, intracavernous prostaglandin E1 and penile implant surgery for erectile dysfunction in urology practice. *J Urol.* 2003 Jul;170:159-63.
65. Rew KT, Heidelbaugh JJ. Erectile Dysfunction. *Am Fam Physician.* 2016 Nov 15;94(10):820-827. Accessed June 28, 2021. Available at URL address <https://www.aafp.org/afp/2016/1115/p820.html>
66. Rosen RC, Wing R, Schneider S, Gendrano N 3rd. Epidemiology of erectile dysfunction: the role of medical comorbidities and lifestyle factors. *Urol Clin North Am.* 2005 Nov;32(4):403-17, v.
67. Sadeghi-Nejad H. Penile prosthesis surgery: a review of prosthetic devices and associated complications. *J Sex Med.* 2007 Mar;4(2):296-309.
68. Santucci RA, Bandukwala NQ. Penile prosthesis implantation. Updated August 31, 2016. Accessed June 25, 2021. Available at URL address: <https://emedicine.medscape.com/article/446761-overview>
69. Satkunasivam R, Appu S, Al-Azab R, Hersey K, Lockwood G, Lipa J, et al. Recovery of erectile function after unilateral and bilateral cavernous nerve interposition grafting during radical pelvic surgery. *J Urol.* 2009 Mar;181(3):1258-63.
70. Seftel AD, Mohammed MA, Althof SE. Erectile dysfunction: etiology, evaluation, and treatment options. *Med Clin North Am.* 2004 Mar;88(2).
71. Shendy WS, Elsoghier OM, El Semary MM, Ahmed AA, Ali AF, Saber-Khalaf M. Effect of low-intensity extracorporeal shock wave therapy on diabetic erectile dysfunction: Randomised control trial. *Andrologia.* 2021 May;53(4):e13997.
72. Siddiqui KM, Billia M, Mazzola CR, Alzahrani A, Brock GB, Scilley C, et al. Three-year outcomes of recovery of erectile function after open radical prostatectomy with sural nerve grafting. *J Sex Med.* 2014 Aug;11(8):2119-24.
73. Sim HG, Klot M, Lange PH, Ellis WJ, Takayama TK, Yang CC. Two-year outcome of unilateral sural nerve interposition graft after radical prostatectomy. *Urology.* 2006 Dec;68(6):1290-4.
74. Sokolakis I, Dimitriadis F, Teo P, Hatzichristodoulou G, Hatzichristou D, Giuliano F. The Basic Science Behind Low-Intensity Extracorporeal Shockwave Therapy for Erectile Dysfunction: A Systematic Scoping Review of Pre-Clinical Studies. *J Sex Med.* 2019 Feb;16(2):168-194.
75. Sramkova T, Motil I, Jarkovsky J, Sramkova K. Erectile Dysfunction Treatment Using Focused Linear Low-Intensity Extracorporeal Shockwaves: Single-Blind, Sham-Controlled, Randomized Clinical Trial. *Urol Int.* 2020;104(5-6):417-424.
76. Srirangam SJ, Manikandan R, Hussain J, Collins GN, O'Reilly PH. Long-term results of extracorporeal shockwave therapy for Peyronie's disease. *J Endourol.* 2006 Nov;20(11):880-4.

77. Strebel RT, Suter S, Sautter T, Hauri D. Extracorporeal shockwave therapy for Peyronie's disease does not correct penile deformity. *Int J Impot Res.* 2004 Oct;16(5):448-51.
78. Taylor FL, Levine LA. Peyronie's Disease. *Urol Clin N Am.* 2007 Nov; 34(4):517-34.
79. United States Food and Drug Administration (FDA). Code of Federal Regulations Title 21, Subpart D-- Sec. 876.3350 Penile inflatable implant. April 1, 2020. Page last updated November 10, 2020. Accessed June 28, 2021. Available at URL address:  
<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=876&showFR=1&subpartNode=21:8.0.1.1.25.4>
80. United States Food and Drug Administration (FDA). Class II special controls guidance document: external penile rigidity devices. Dec 28, 2004. Page last updated Jun 29, 2018. Accessed June 28, 2021. Available at URL address:  
<http://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/GuidanceDocuments/ucm072098.htm>
81. United States Food and Drug Administration (FDA). Guidance for the content of premarket notifications for penile rigidity implants. Jan 16, 2000. Page last updated March 12, 2018. Accessed June 28, 2021. Available at URL address:  
<http://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/GuidanceDocuments/ucm073769.htm>
82. Vardi Y, Appel B, Kilchevsky A, Gruenwald I. Does low intensity extracorporeal shock wave therapy have a physiological effect on erectile function? Short-term results of a randomized, double-blind, sham controlled study. *Urol.* 2012 May;187(5):1769-75.
83. Vinay J, Moreno D, Rajmil O, Ruiz-Castañe E, Sanchez-Curbelo J. Penile low intensity shock wave treatment for PDE5I refractory erectile dysfunction: a randomized double-blind sham-controlled clinical trial. *World J Urol.* 2021 Jun;39(6):2217-2222.
84. Wilson SK, Delk JR, Salem EA, Cleves MA. Long-term survival of inflatable penile prostheses: single surgical group experience with 2,384 first-time implants spanning two decades. *J Sex Med.* 2007 Jul;4(4 Pt 1):1074-9.
85. Young Academic Urologists Men's Health Group, Fode M, Hatzichristodoulou G, Serefoglu EC, Verze P, Albersen M. Low-intensity shockwave therapy for erectile dysfunction: is the evidence strong enough? *Nat Rev Urol.* 2017 Oct;14(10):593-606.
86. Zermann DH, Kutzenberger J, Sauerwein D, Schubert J, Loeffler U. Penile prosthetic surgery in neurologically impaired patients: long-term followup. *J Urol.* 2006 Mar;175(3 Pt 1):1041-4; discussion 1044.
87. Zorn KC, Bernstein AJ, Gofrit ON, Shikanov SA, Mikhail AA, Song DH, et al. Long-term functional and oncological outcomes of patients undergoing sural nerve interposition grafting during robot-assisted laparoscopic radical prostatectomy. *J Endourol.* 2008 May;22(5):1005-12.
88. Zou ZJ, Tang LY, Liu ZH, Liang JY, Zhang RC, Wang YJ, Tang YQ, Gao R, Lu YP. Short-term efficacy and safety of low-intensity extracorporeal shock wave therapy in erectile dysfunction: a systematic review and meta-analysis. *Int Braz J Urol.* 2017 Sep-Oct;43(5):805-821.

---

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.