

Medical Coverage Policy



Effective Date.....04/15/2021
Next Review Date04/15/2022
Coverage Policy Number0448

Interspinous Process Spacer Devices

Table of Contents

Coverage Policy.....	1
Overview.....	1
General Background.....	1
Medicare Coverage Determinations	10
Coding/Billing Information.....	10
References	11

Related Coverage Resources

[Intervertebral Disc \(IVD\) Prostheses](#)
[Lumbar Fusion for Spinal Instability and Degenerative Disc Conditions, Including Sacroiliac Fusion](#)
[Minimally Invasive Intradiscal/Annular Procedures and Trigger Point Injections](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgement and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Coverage Policy

Interspinous/interlaminar process spacer devices are considered experimental, investigational or unproven for all indications.

Overview

This Coverage Policy (CP) addresses interspinous/interlaminar process spacers devices (e.g., coflex®, Superior®).

Note: Dynamic spine stabilization device systems and interspinous fixation/posterior non-pedicle supplemental fixation devices (e.g., coflex-F®) are addressed in CP 0303 titled Lumbar Fusion for Spinal Instability and Degenerative Disc Conditions, Including Sacroiliac Fusion.

General Background

An interspinous/ interlaminar process spacer device may also be referred to as interspinous spacers (ISS), interspinous/ interlaminar stabilization/ distraction devices, and interspinous process decompression (IPD) systems/devices. They are proposed for patients with lumbar spinal stenosis, with or without Grade 1

spondylolisthesis, who experience relief in flexion from their symptoms of leg/buttocks/groin pain, with or without back pain, and who have undergone at least six months of non-operative treatment. The design of devices and the materials used in devices vary. The use of various devices has been proposed both as a minimally invasive surgical alternative to standard posterior lumbar decompression, with or without fusion procedures, and as an addition to decompressive surgery.

The American Academy of Orthopedic Surgeons estimates that spinal stenosis affects 8 to 11 percent of the population. Spinal stenosis is a narrowing of the vertebral canal that may lead to compression of the spinal nerves or nerve roots, especially in the lumbar vertebrae. Lumbar stenosis is commonly seen in an aging or degenerative spine. Neurogenic claudication is a combination of low back and leg pain, with numbness and motor weakness when standing or walking that is relieved by sitting or lying. Treatment for back pain may include pharmacological therapy (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], analgesics, and muscle relaxants), exercise, spinal manipulation, acupuncture, cognitive-behavioral therapy, and physical therapy. Various interventional and surgical procedures may be considered if these measures are unsuccessful. Surgical options include decompressive procedures (e.g., laminectomy) alone, or decompression and fusion. Fusion is frequently performed with rigid implant fixation systems, including pedicle screws and interbody cages.

U.S. Food and Drug Administration (FDA)

The two current interspinous/interlaminar process spacers that are FDA approved and commercially available are the coflex[®] and Superior[®] devices. Coflex[®] is intended to be implanted after a decompression of the canal has been performed at the affected levels. Superior[®] is intended to “stand-alone” (does not requiring surgical decompression). It is delivered percutaneously as a single-piece through a cannula after dilators have opened the interspinous space.

coflex[®] Interlaminar Technology (Paradigm Spine, LLC, New York, NY): The coflex[®] Interlaminar Technology received FDA approval through the PMA process on October 17, 2012. Since the original approval, there have been numerous supplemental approvals issued relating to the post-approval study. According to the FDA Summary of Safety and Effectiveness, the coflex Interlaminar Technology is an interlaminar stabilization device indicated for use in one or two level lumbar stenosis from L1-L5 in skeletally mature patients with at least moderate impairment in function, who experience relief in flexion from their symptoms of leg/buttocks/groin pain, with or without back pain, and who have undergone at least six months of non-operative treatment. The coflex is intended to be implanted midline between adjacent lamina of one or two contiguous lumbar motion segments. Interlaminar stabilization is performed after decompression of stenosis at the affected level(s) (FDA, 2012).

Description: The coflex[®] device is a U-shaped implant manufactured from medical-grade titanium alloy designed to withstand a normal physiologic load in the spine. The device is a single-piece design with 2 pairs of serrated wings: 1 pair extending from the upper long arm and the other pair extending from the lower long arm of the U. This design allows a simple press-fit insertion of the device. The U portion is positioned horizontally between 2 adjacent spinous processes and pressed into place. The wings are crimped over bone to hold the implant in place. Implantation is performed after decompression of stenosis at the affected level(s).

Superior[™] Indirect Decompression System (IDS) (Boston Scientific Corporation) / Superior[®] InterSpinous Spacer (VertiFlex[®], Inc., San Clemente, CA. Vertiflex was purchased by Boston Scientific in June 2019): The Superior InterSpinous Spacer (ISS) received FDA approval through the PMA process on May 20, 2015 (P140004). Since the original approval, there have been numerous supplemental approvals issued relating to the post-approval study. The ISS is indicated to treat skeletally mature patients suffering from pain, numbness, and/or cramping in the legs (neurogenic intermittent claudication) secondary to a diagnosis of moderate degenerative lumbar spinal stenosis, with or without Grade 1 spondylolisthesis, confirmed by X-ray, MRI and/or CT evidence of thickened ligamentum flavum, narrowed lateral recess, and/or central canal or foraminal narrowing. The Superior ISS is indicated for those patients with impaired physical function who experience relief in flexion from symptoms of leg/buttock/groin pain, numbness, and/or cramping, with or without back pain, and who have undergone at least 6 months of non-operative treatment. The Superior ISS may be implanted at one or two adjacent lumbar levels in patients in whom treatment is indicated at no more than two levels, from L1 to L5 (FDA, 2015).

Description: The Superior Interspinous Spacer is a 1-piece, fully assembled implant designed to fit between the spinous processes of the lumbar spine. It consists of the titanium alloy body with 2 Cam lobes ("wings") superior and inferior to the main body that secure the device in place. Once the implant is in place, an actuation mechanism opens the implant to provide distraction and minimize flexion in the targeted spinal region. Superior is delivered through a single small incision (12-15 millimeters [mm]) in the patient's back utilizing proprietary manual instrumentation. The device may be implanted at 1 or 2 adjacent levels from L1 to L5 to expand the space between the vertebral spinous processes. The Superior Interspinous Spacer System includes a set of single-use stainless steel instruments necessary to deliver the implant, including a dilator, cannula, reamer, interspinous gauge, inserter, and driver. The procedure can be performed on an outpatient basis under local anesthesia using fluoroscopic guidance. However, in the Investigating Superior™ In Spinal Stenosis randomized controlled trial (Patel, et al., 2015a), 82% of patients had general anesthesia, 13.2% had conscious sedation, and only 7.4% had local anesthesia. Although the Vertiflex Procedure is noted in some resources as percutaneous, in the same study only 46.8% had a percutaneous placement and 53.2% had a "miniopen" procedure.

X-STOP® Interspinous Process Decompression (IPD) System (Medtronic, Minneapolis, Minnesota): The X-Stop device received FDA premarket approval in 2005, with promising results in the short-term, but further research demonstrated minimal benefit with longer-term follow-up, along with relatively high complication rates including spinous process fracture. PMA withdrawal date was 04/30/2015. Medtronic discontinued the distribution of the X-Stop system in 2015.

DIAM® Spinal Stabilization System: The FDA recommended against approval for the DIAM system (Medtronic, P140007) in an Orthopaedic & Rehabilitation Devices panel meeting in February 2016.

Other devices undergoing study but not currently FDA-approved include:

- Wallis® Posterior Dynamic Stabilization System (Zimmer Biomet, INC., Finland/Warsaw, IN, USA)
- APERIUS™ implant used in APERIUS™ PercLID™ System (Medtronic)
- HeliFix® Interspinous Spacer System (Alphatec Spine, Carlsbad, CA)

Literature Review

coflex® Interlaminar Technology: FDA approval of the coflex was based on an Investigational Device Exemption (IDE) randomized, multicenter trial conducted by Davis et al. (2013a). A total of 322 patients who met the following criteria were included: ages of 40 and 80, at least moderate lumbar stenosis, which narrows the central spinal canal at one or two contiguous levels from L1–L5 that require surgical decompression, and BMI not > 40. Patients were followed for two years. Patients received laminectomy and coflex insertion (n=215) or posterolateral spinal fusion with pedicle screw (PS) instrumentation (n=107). The proportion of patients with spondylolisthesis was similar (coflex: 99/215 = 46.0%; PS [control]: 51/107 = 47.7%). Composite Clinical Success (CCS) criteria included some of the following: device survives 24 months, no epidural injections in 24 months, Oswestry Disability Index (ODI) improvement from baseline to month 24 visit of at least 15 points, no persistent new or worsening sensory or motor deficit, and no major device-related complications. Five year results were reported, with a five year follow-up rate of 91% (Musacchio, et al. 2016).

- Based on composite for overall success, 66.2% of coflex and 57.7% of PS succeeded (p= 0.999), thus demonstrating non-inferiority at 24 months. At 5-year follow-up, 50.3% of coflex patients met the success criteria compared with 44% of PS patients (p>0.35; not significant). Of note, patients who underwent additional surgery or injections after the study surgery were classified outcome failures in the composite assessment of success, and excluded from the analyses of individual outcome assessments such as Visual Analogue Scale (VAS) and Zurich Claudication Questionnaire (ZCQ).
- ODI (percent achieving a 15-point reduction in ODI): At 2 years, coflex 85.8%, PS 76.7%; at 5 years coflex 80.6%; PS 73.2%.
- VAS back and leg pain: Both groups showed significant improvement from baseline in both back and leg pain at all time points out to 5 years.
- ZCQ: At 5 years, the ZCQ symptom severity improvement of at least 0.5 was assessed per patient where 79.8% of coflex and 72.7% of PS met that criteria. The ZCQ physical function improvement of at least 0.5 was assessed per patient where 78.2% of coflex and 70.9% of PS patients met the criteria. The

third component of patient satisfaction was consistent from Week 6 to Month 60 in both groups (percentage of patients meeting criteria was not provided).

- Narcotic usage: There was no significant difference between the two groups at Month 60 but both groups were significantly improved from their preoperative status.
- Reoperation rates: Reoperation rate at 24 months was coflex 23/215 (10.7%) and PS 8/107 (7.5%) ($p=0.426$). At 5-year, cumulative total occurrences of reoperations/revisions were coflex group 35/215 (16.3%) and PS group 19/107 (17.8%).

Abjornson et al. (2018) reported a Davis/Musacchio sub-study on the cohort treated with decompression plus coflex at 1 or 2 levels and who did not present with spondylolisthesis preoperatively ($n=116$, 65 in the 1-level coflex group and 51 in the 2-level coflex group).

- CCS was achieved in 48.3% of 1 level and 60.9% of 2 level at 5 years.
- ODI: There was 15-point improvement in 81.6% of 1-level and 90.3% of 2-level patients at 5 years.
- VAS: 94.7% of 1 level and 100% of 2 level achieved at least a 20-mm improvement, at 5 years.
- ZCQ: Improvement of ≥ 0.5 as compared to preoperative score is calculated as improvement. In the symptom severity component, 81.6% of 1-level and 83.9% of 2-level patients reported improvement at 5 years. In the physical function component, 76.3% of 1 level and 83.9% of 2 level patients reported improvement at 5 years.
- No secondary surgery or epidural injections: 1 level (69.2%); 2 level (70.6%); at 5 years. Of the 16 patients that required a secondary surgery over the course of the study, 8 were not related, 2 unlikely, 4 possibly, and 2 definitely related to the device.

A limitation of this study is the lack of comparison with non-spondylolisthesis patients from the 'fusion with pedicle screw' cohort.

Davis et al. (2013b) reported a Davis/Musacchio sub-study on the cohort of patients with low grade (grade 1) degenerative spondylolisthesis with spinal stenosis (coflex =99, PS=51). Two years results include:

- Overall success (as described above) was similar with 59 coflex patients meeting success (62.8%), and 30 PS patients (62.5%) meeting success at 24 months.
- ODI: The percentage of patients that achieved a 15-point reduction in ODI at 2 years from baseline was 86.1% for coflex and 81.0% for PS.
- VAS back and leg pain: no significant differences noted at baseline or at 24 months.
- ZCQ: both groups improved similarly from baseline in physical function and symptom severity scores, but the coflex cohort performed significantly better than PS controls with respect to ZCQ patient satisfaction at 24 months ($p = 0.05$).
- Reoperation: The overall reoperation rate was 14.1% (14 of 99) and 5.9% (3 of 51) for the coflex and PS controls, respectively ($p = 0.18$, not statistically significant).

Simon et al. (2018) reported a Davis/Musacchio sub-study on the cohort of 116 patients who required surgical treatment at two levels. The decompression and interlaminar stabilization with coflex group consisted of 77 patients, and the posterolateral spinal fusion with pedicle screw instrumentation (PS) group consisted of 39 patients.

- CCS: the percentage of patients who achieved CCS, coflex 55.1%, PS 36.4% at 5 years.
- ODI: Of those assessed, 86.7% of coflex patients (39/45) and 92.9% of PS patients (13/14) saw an improvement of ≥ 15 points in the ODI at month 60 compared to baseline
- VAS: At 60 months, the mean VAS back pain scores decrease of 59.8 points for the ILS group was similar to the fusion group with 58.9 points.
- ZCQ: There was no significant difference between groups for Symptom Severity, Physical Function or Satisfaction score, at 60 months.
- No secondary surgery or epidural injections: The number of patients in the coflex group who did not receive a reoperation or epidural injection was 53/77 (68.8%) compared to 20/39 (51.3%) PS patients.

The European Study of Coflex And Decompression Alone (ESCADA) trial (Schmidt, et al., 2018) is a randomized controlled trial with modified intent-to-treat analysis. The trial included 230 patients seen at seven sites in Germany. Schmidt et al. compared open microsurgical decompression followed by interlaminar stabilization with coflex (D+ILS) to decompression alone (DA).

- Inclusion criteria included age > 40 years, VAS back pain score of ≥ 50 mm, at least moderate degenerative spinal stenosis, with constriction of the central spinal canal in 1 or 2 adjacent segments from L-3 to L-5 with the need for decompression. In addition, the following was allowed but not required: hypertrophy of the facet joints and subarticular recess stenosis in the relevant segment or stenosis of the foramen in the relevant segment, and/or spondylolisthesis (anterolisthesis or retrolisthesis) up to grade I.
- Exclusions were translational instability in the main segment as well as in adjacent segments (dynamic translational instability ≤ 3 mm), previous surgery at index level, and/or vertebral or pars fracture.

At 24 months, with an overall 91% follow up rate, results demonstrated no significant differences between the groups in the patient reported outcomes: ODI scores, ZCQ, and VAS back and neck pain scores ($p > 0.05$). The CCS was calculated using 1) ODI success with improvement > 15 points; 2) survivorship with no SSIs or lumbar injections; 3) neurological maintenance or improvement without worsening; and; 4) no device- or procedure-related severe adverse events. The DA arm had 228% more lumbar injections ($p = 0.0065$) than the D+ILS arm (epidural steroid injections for D+ILS, 5/110 [4.5%]; for DA, 17/115 [14.8%]). When this measurement was included in the CCS, the result became significant. Authors conclude the use of coflex extends the durability and sustainability of a decompression procedure.

A Hayes Technology Brief (September 21, 2018) on the use of the coflex Interlaminar stabilization device with decompression for the treatment of lumbar spinal stenosis in adult patients with and without spondylolisthesis noted an overall low-quality body of evidence. Hayes suggested areas of uncertainty that remain include definitive patient selection and demonstration of a distinct benefit of the coflex device over traditional surgical intervention.

A randomized controlled double-blind 'FELIX' trial (Moojen, et al., 2013) was conducted at five neurosurgical centers in the Netherlands to assess whether interspinous process device implantation is more effective in the short term than conventional surgical decompression for patients age 40 and 85 years with NIC due to lumbar spinal stenosis. Patients with lumbar spinal stenosis at one or two levels with an indication for surgery were randomized to treatment with coflex device (no bony decompression was done) ($n=80$) or surgical decompression ($n=79$). The difference in ZCQ scores coflex group and the standard decompression group at eight weeks (63% vs. 72%, $p=0.44$) or one year (66% vs. 69%, $p=0.77$) is not significant. However, the repeat surgery rate in the coflex was significantly higher than in the standard decompression group, at 29% vs. 8% ($p<0.001$). The authors stated "the number of reoperations in the interspinous process device treatment arm is very worrisome, especially because reoperations do not reach the success rate of primary surgeries; use of interspinous process devices might even prevent recovery in 20% of patients".

Richter et al. (2014) conducted a prospective, controlled study to assess the outcome of symptomatic lumbar spinal stenosis (LSS) treated with decompressive surgery alone ($n=31$) compared to decompressive surgery with implantation of the coflex interspinous device ($n=31$). Included patients had signs, symptoms and MRI findings of lumbar spinal stenosis and a minimum of three months of conservative treatment, were age 45-80 with one or two level stenosis, and had not undergone previous surgery of the lumbar spine. There was no formal randomization procedure. There was a significant improvement in both groups ($p>0.001$) in the clinical outcome assessed in the ODI, the Roland-Morris Disability Questionnaire, the VAS, and the pain-free walking distance at all time of assessment compared to baseline. Up to two years after surgery, there were no significant differences between the two groups in all measured parameters, including patient satisfaction and subjective operation decision. In the coflex group, three revisions with pedicle screw fusion of the segment were necessary. In surgery only group, two patients had to be instrumented and fused. The authors concluded that the additional placement of a coflex interspinous device does not improve the already good clinical outcome after decompressive surgery for lumbar spinal stenosis in the 24-month follow-up interval.

In a prospective study, Kumar et al. (2014) compared decompression plus coflex ($n=22$) to decompression alone ($n=24$). The included 46 patients were 40–74 years old with symptomatic lumbar spinal stenosis. The mean ODI score for both the coflex and the comparison group showed significant improvement at six months, one year, and two years as compared to the preoperative score. The mean improvement in ODI scores of patients in the coflex group was significantly greater than the comparison group ($p<0.001$). The incidence of complications in the two groups was not significantly different ($p=0.35$). The authors support the implantation of coflex after spinal decompression.

Li et al. (2019) reported on a retrospective study including 99 patients with degenerative lumbar disease (DLD) at L3–L5. A total of 45 patients underwent ‘Topping off’ surgery or ‘hybrid’ surgery (L4–5 posterior lumbar interbody fusion [PLIF] + L3–4 Coflex) and 54 patients underwent PLIF = L3–5 PLIF. Patients were excluded if they had degenerative lumbar scoliosis or kyphosis, lumbar spine fracture, spondylolisthesis at L3–L4 of grade II and above, severe osteoporosis, and history of lumbar spine surgery. The primary study outcome was to assess the efficacy of preventing adjacent segment degeneration (ASD).

- Outcomes showed both two groups had a significant improvement in VAS and ODI scores for lower back/leg pain at 3 years after surgery than before ($p < 0.05$). But there was no significant difference in the pairwise comparison ($p > 0.05$).
- The two groups had no significant difference in intervertebral mobility (L2–L3) before surgery ($p > 0.05$). After surgery, it was lower in the Topping-off group than in the PLIF group ($p < 0.05$). At 3 years after surgery, general adjacent segment mobility (GASM) (L2–4) was not significantly different between the two groups ($p > 0.05$).
- At 3 years after surgery, the modified Pfirrmann grade of disc was increased by 1 grade in 2 cases of the Topping-off group (4.44%). In contrast, disc degeneration was more severe in the PLIF group, with increased Pfirrmann grade in 14 cases (25.93%) (including 2 cases with intervertebral mobility $> 10^\circ$). Among them, 11 cases had an increase by 1 grade, 2 cases 2 grade, and 1 case 3 grade (this patient received a revision surgery). The difference was of statistical significance between the groups ($p < 0.05$).

In the Topping-off group, one case had intraspinal hematoma after surgery; in the PLIF group, one case had subcutaneous incision infection and another case intraspinal hematoma. The clinical efficacy and the incidence of adjacent segment degeneration (ASD) of Topping-off surgery for degenerative lumbar disease (DLD) remain to be verified by trials with a larger sample size and longer follow-up.

In a long term retrospective study, patients receiving coflex implant were followed for a minimum of 8 years (Du, et al., 2020). A total of 56 patients who underwent single segment interlaminar decompression and coflex implantation were followed for 107.6 ± 13.3 months. Results showed ROM of adjacent segments increased at 6 months and at the last follow-up compared with that before surgery ($P > 0.05$). At 6 months after surgery, intervertebral space height (ISH) and intervertebral foramen height (IFH) of implanted segment was significantly higher than that before surgery ($P < 0.05$). At the last follow-up, there was a decrease in ISH and IFH ($P > 0.05$). During the follow-up period, a total of 11 patients (19.6%) experienced complications and 6 patients (10.7%) underwent secondary surgery.

Two small retrospective studies report greater than five year results. A small retrospective study reported at least five year results on 87 patients (Yuan, et al., 2017) with a total of 42 patients who underwent decompression and coflex interspinous stabilization. A total of 45 patients had decompression and posterior lumbar interbody fusion (PLIF). The mean ODI and VAS scores in the coflex group were significantly lower compared with the PLIF group initially. However, at final follow-up, the mean ODI scores between the two groups had no significant difference. At final follow-up, the index level ROM was significantly higher in the coflex group. At the final follow-up, two (4.8%) patients in the coflex group required revision surgery for ASD, five (11.1%) patients in the PLIF group underwent a revision surgery for ASD (did not reach statistical significance; $p = 0.277$). Errico et al. (2009) reported retrospective results from one orthopedic spine surgeon who followed 127 patients for a mean of 6.3 years. A patient satisfaction query demonstrated that 7% were unsatisfied, 46% were satisfied, and 46% were very satisfied with their clinical outcome. Based on the follow-up radiographs, 92 of patients had no device related issues and 8% had device-related issues. Both studies are limited by their small, retrospective design.

A small, retrospective cohort of adults with lumbar stenosis and grade 1 stable spondylolisthesis ($n=83$) were evaluated to compare postoperative outcomes following single-level decompression and implantation of coflex ($n=46$) and single-level laminectomy alone ($n=37$). Mean follow-up was 516 to 677 days. Patients who received coflex were older, had a higher anesthesia grade but similar comorbidities. Results demonstrated the coflex cohort had higher estimated blood loss ($p = 0.004$), longer operative time ($p = 0.001$), and longer length of stay ($p = 0.001$). Total perioperative complications (21.7% vs 5.4%, $p = 0.035$) and instrumentation related complication was higher in the coflex cohort (10.9% vs 0% laminectomy group, $p = 0.039$). Similar post-op complications, revision, and neurologic complication rates were seen between the two cohorts at last follow up (Zhong, et al., 2020).

coflex® Literature Review Summary: Studies in the published peer reviewed scientific literature include small populations, especially considering the prevalence of lumbar stenosis. One trial demonstrated a significantly higher reoperation rate that may actually prevent a better recovery owing to the lower recovery rate after a second operation. Published studies do not demonstrate any long-term health outcome advantage with the additional use of coflex. Large population sufficiently powered randomized controlled trials that demonstrate long-term health outcome advantages with the addition of coflex are needed.

Superion® InterSpinous Spacer

Patel et al. (2015a) conducted a prospective, multicenter, randomized controlled investigational device exemption trial to compare two year outcomes in patients with NIC secondary to moderate lumbar spinal stenosis (LSS) who were treated with the Superior spacer or a control spacer (X-STOP). Eligible patients were at least 45 years of age and reported symptoms of NIC secondary to a confirmed diagnosis of LSS at one or two contiguous levels from L1 to L5, despite at least six months of nonsurgical management. A total of 391 randomized patients were implanted with Superior (n = 190) or X-STOP (n = 201) spacers at 29 sites in the United States. At study end, participation was 280, Superior (n = 136) or X-STOP (n = 144) spacers. A total of 28% were lost in follow-up: 111 withdrawn due to a protocol-defined secondary intervention, including device explant, revision surgery at the index level without explant, rhizotomy, rehospitalization for deep infection, or lumbar injection at the index level. The primary endpoint of this study was a composite treatment success outcome at the two year follow-up visit, defined as: (1) clinically significant improvement in at least 2 of 3 ZCQ domain scores compared with baseline (2) freedom from reoperation, revision, removal, or supplemental fixation at the index level, (3) freedom from epidural steroid injection or nerve block at the index level within 12 weeks of the 2-year visit, (4) freedom from rhizotomy or spinal cord stimulator at any level, and (5) freedom from major implant or procedure-related complications.

At two years follow up, the authors stated that the primary composite endpoint of this study was met, which demonstrated that the Superior spacer was non-inferior to the X-Stop spacer. Leg pain, the predominant patient complaint, decreased in severity by 70% during 2 years in each group. Most (77%) patients achieved leg pain clinical success (improvement ≥ 20 mm) at 2 years. Back pain clinical success (improvement ≥ 20 mm) was 68%, with no differences between groups. Oswestry Disability Index clinical success ($\geq 15\%$ point improvement) was achieved in 65% of patients. There were a total of 44 (23.2%) reoperations or revisions in the Superior group compared with 38 (18.9%) in the X-STOP (control) group ($p = 0.32$). The authors noted that the long-term durability of interspinous process spacers is currently unknown and requires further investigation.

- Laurysen et al. (2015) performed a qualitative comparison of the published two-year clinical findings from Patel et al. (2015a) with historical laminectomy literature, (19 studies) for similar outcome measurements associated with decompressive laminectomy (N=1045). The 19 studies included retrospective, prospective, and randomized trials. Back and leg pain, ODI, and ZCQ values were compared. Following treatment with either spacer or laminectomy, patients attained clinically substantial gains across all outcome measures at 12 months with durable improvement through 24 months, postoperatively. The authors of this literature review that included retrospective studies concluded that both treatments provide effective and durable symptom relief of claudicant symptoms.
- Patel et al. (2015b) reported three year outcomes. All outcomes were reported using a modified intention-to-treat population. At year three, 36.4% are lost in follow up (Superion = 120 or X-STOP = 129). The 'primary composite endpoint' was individual patient success based on four components: improvement in two of three domains of the Zurich Claudication Questionnaire, no reoperations at the index level, no major implant/ procedure-related complications, and no clinically significant confounding treatments. The proportion of subjects achieving the 'primary composite endpoint' was greater for Superior (63/120, 52.5%) than for X-STOP (49/129, 38.0%) ($p=0.023$). Comparing the 24-month data with the 36-month data, there was a higher increase in X-STOP reoperations, revisions, and removals (n=15 out of 44 total) compared to the Superior device (n=11 out of 49 total).

Nunley et al. (2017) reported five year outcomes on the Superior arm of the Patel et al. (2015a) trial. Of the original 190 patients randomly assigned to receive treatment with Superior, 88 were free from reoperation or steroid injection at 5-year follow-up and able to provide complete clinical outcome evaluations (46.3%). Authors' report 74 of 88 patients (84%) demonstrated clinical success on at least 2 of 3 ZCQ domains (symptom severity,

physical function, and patient satisfaction). A limitation of this study is the loss of participation at five year follow-up (88 of 190 = 46.3%).

- In a quality of life sub-study, Nunley et al. (2018a) reported that of 189 patients initially randomized to Superion treatment, SF-12 questionnaire responses were captured in 68 study subjects at 5 years. Physical component summary and mental component summary (PCS, MCS) scores were computed preoperatively and the percentage improvement in PCS and MCS at the 5-year follow-up interval compared to preoperative values was computed. The mean PCS score improved from 29.4 ± 8.1 preoperatively to 43.8 ± 11.6 at 5 years, representing average percentage improvements of 49%, ($p < 0.001$). 87% (59 of 68) of subjects who provided 5-year SF-12 responses continued to maintain or improve their PCS score. The mean MCS score improved from 50.0 ± 12.7 preoperatively to 54.7 ± 8.6 at 5 years, representing a 9% improvement ($p > 0.10$ for both comparisons). At 5 years, 57% (39 of 68) of subjects showed maintenance or improvement in PCS scores.
- Nunley et al. (2018b) reported an opioid-medication analysis of the Superion arm of the Patel et al. (2015a) trial. At baseline, almost 50% (94 of 190) of subjects were using opioid medication. Thereafter, there was a sharp decrease in opioid-medication prevalence from 25.2% (41 of 163) at 12 months to 13.3% (20 of 150) at 24 months to 7.5% (8 of 107) at 60 months. A similar pattern was also observed among subjects with a history of opiates prior to entering the trial.

Tekmyster et al. (2019) reported on a prospective registry of 445 patients at multiple US sites. The objective was to report 12 month outcomes of pain severity data and patient satisfaction after interspinous process decompression (IPD) with a stand-alone interspinous spacer (Superion). The maximum number of patients providing pain severity data was 2,090, 759, 1,553 and 445 at baseline, 3 weeks, 6 and 12 months, respectively. For patient satisfaction and treatment approval, the maximum number of patients providing follow-up data was 751, 1,542 and 443 at 3 weeks, 6 and 12 months, respectively.

- Mean leg pain severity decreased from 76.6 ± 22.4 mm preoperatively to 33.0 ± 29.9 mm at 3 weeks, 33.1 ± 34.0 mm at 6 months, and 30.4 ± 34.6 mm at 12 months, reflecting an overall 60% improvement.
- Back pain severity improved from 76.8 ± 22.2 mm preoperatively to 37.5 ± 29.6 mm at 3 weeks, 41.9 ± 32.5 mm at 6 months, and 39.9 ± 32.3 mm at 12 months (48% improvement).
- For patient satisfaction at 3 weeks, 6 and 12 months, 89%, 80%, and 80% were satisfied or somewhat satisfied with their treatment and 90%, 75%, and 75% would definitely or probably undergo the same treatment again.

In the phone survey, the rate of revision was 3.6% (51 of 1,426).

In a prospective study, Bini et al. (2011) observed 121 patients following insertion of the Superion device. Patients had a diagnosis of moderate lumbar spinal stenosis, failed 3 months conservative treatment, and persistent pain relieved by lumbar flexion. A total of 22 (18%) of the patient's patients presented with concomitant grade I spondylolisthesis. A total of 52 were observed at 12 months. ODI improved 64% ($p < 0.001$) through 12 months and clinical success was 92%. Extremity and axial pain improved 53% and 49% (both $p < 0.001$), respectively, through 12 months with clinical success of 76% for axial pain and 86% for extremity pain. The follow-up period in the current study extends only through 12 months so no direct comparison of complication and revision rates can be made with certainty.

Superion® Literature Review Summary: There is a lack of large well-designed studies in the peer review scientific literature comparing stand alone use of Superion device to established surgical decompression. Published studies do not demonstrate any long-term health outcome advantage with the use of Superion as an alternative to standard surgical treatment. Large population sufficiently powered randomized controlled trials that demonstrate long-term health outcome advantages are needed.

Professional Societies/Organizations

American Academy of Orthopaedic Surgeons (AAOS): At this time, there are no AAOS Clinical Practice Guidelines or AAOS Appropriate Use Criteria addressing the use of interspinous/interlaminar spacer devices.

On the AAOS website, under Disease & Conditions, under Lumbar Spinal Stenosis, the AAOS states: Interspinous process devices, or spacers, are inserted between the spinous processes in the back of the spine. These devices spread the vertebrae apart and keep the space for the nerves open and functioning. This

procedure is a minimally invasive surgical option for lumbar spinal stenosis. Interspinous process spacers were approved in 2005. Many procedures have been performed since then. In some studies, success rates are greater than 80 percent. Numerous spacer devices are currently being evaluated. They may be a safe alternative to an open laminectomy for some patients. Limited bone (lamina) is removed with this procedure, and it may be performed under local anesthesia. The key to success with this procedure is appropriate selection of the patients. The appropriate candidate must have relief of buttock and leg pain when sitting or bending forward. The pain returns upon standing.

North American Spine Society (NASS):

Lumbar Interspinous Device without Fusion and Decompression Coverage Policy Recommendations (May 2018) apply to interspinous process ISP devices that are intended to be used in conjunction with a direct decompressive procedure and include:

- Stabilization with an ISP without fusion in conjunction with laminectomy may be indicated as an alternative to lumbar fusion for degenerative lumbar stenosis with or without low-grade spondylolisthesis (less than or equal to 3 mm of anterolisthesis on a lateral radiograph) with qualifying criteria when appropriate:
 - significant mechanical back pain is present (in addition to those symptoms associated with neural compression) that is felt unlikely to improve with decompression alone. Documentation should indicate that this type of back pain is present at rest and/or with movement while standing and does not have characteristics consistent with neurogenic claudication.
 - a lumbar fusion is indicated post-decompression for a diagnosis of lumbar stenosis with a Grade 1 degenerative spondylolisthesis as recommended in the NASS Coverage Recommendations for Lumbar Fusion.
 - a lumbar laminectomy is indicated as recommended in the NASS Coverage Recommendations for Lumbar Laminectomy.
 - previous lumbar fusion has not been performed at an adjacent segment.
 - previous decompression has been performed at the intended operative segment.

- ISP devices are NOT indicated in cases that do not fall within the above parameters. In particular, they are not indicated in the following scenarios and conditions:
 - degenerative spondylolisthesis of Grade 2 or higher.
 - degenerative scoliosis or other signs of coronal instability.
 - dynamic instability as detected on flexion-extension views demonstrating at least 3 mm of change in translation.
 - iatrogenic instability or destabilization of the motion segment.
 - a fusion is otherwise not indicated for a Grade 1 degenerative spondylolisthesis and stenosis as per the NASS Coverage Recommendations for Lumbar Fusion.
 - a laminectomy for spinal stenosis is otherwise not indicated as per the NASS Coverage Recommendations for Lumbar Laminectomy.

The NASS Evidence-Based Clinical Guideline Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis (Kreiner, et al., 2013) states “There is insufficient evidence at this time to make a recommendation for or against the placement of an interspinous process spacing device in patients with lumbar spinal stenosis”.

The NASS Evidence-Based Clinical Guideline Diagnosis and Treatment of Degenerative Lumbar Spondylolisthesis (Matz, et al., 2016) states “There is insufficient and conflicting evidence to make a recommendation for or against the efficacy of interspinous spacers versus medical/interventional treatment in the management of degenerative lumbar spondylolisthesis patients. Grade of Recommendation: I (Insufficient Evidence)”.

The NASS Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis & Treatment of Low Back Pain (last updated 8/10/2020 and endorsed by the American Academy of Physical Medicine and Rehabilitation and American Association of Neurological Surgeons and Congress of Neurological Surgeons) does not address Coflex, Superior or X-Stop.

International Society for the Advancement of Spine Surgery (ISASS): The ISASS Policy Statement Decompression with Interlaminar Stabilization (Guyer, et al., 2016) states that patients who have all of the following criteria may be eligible for decompression with interlaminar stabilization:

1. Radiographic confirmation of at least moderate lumbar stenosis, which narrows the central spinal canal at 1 or 2 contiguous levels from L-1 to L-5 that require surgical decompression. Moderate stenosis is defined as > 25% reduction of the anteroposterior dimension compared with the next adjacent normal level, with nerve root crowding compared with the normal level, as determined by the surgeon on CT scanning or MRI.
2. Radiographic confirmation of the absence of gross angular or translatory instability of the spine at index or adjacent levels (instability as defined by White and Panjabi: sagittal plane translation >4.0 mm or 15% or local sagittal plane rotation > 15° at L1–2, L2–3, and L3–4; >20° at L4–5 based on standing flexion-extension radiographs). Improved imaging technologies are able to better refine/detect previously undetected instability and as these technologies become more established, surgeons should expect to refine with specificity and clear delineation of appropriate surgical candidates requiring stabilization.
3. Patients who experience relief in flexion from their symptoms of leg/buttocks/groin pain, with or without back pain, and who have undergone at least 12 weeks of non-operative treatment consisting of non-steroidal anti-inflammatory drugs and at least one of the following: rest, restriction of activities of daily living, physical therapy, or steroid injections.

The American Board of Internal Medicine’s (ABIM) Foundation Choosing Wisely® Initiative
No relevant information.

Use Outside the U.S.

National Institute for Health and Clinical Excellence (NICE) (United Kingdom): NICE issued guidance in November 2010 on Interspinous distraction procedures for lumbar spinal stenosis causing neurogenic claudication. The revised guidance states that current evidence shows that these procedures are efficacious for carefully selected patients in the short and medium term, although failure may occur and further surgery may be needed. The guidance states that there are no major safety concerns, and that these procedures may be used provided that normal arrangements are in place for clinical governance, consent and audit. The guidance also states that patient selection should be carried out by specialist spinal surgeons who are able to offer patients a range of surgical treatment options.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No National Coverage Determination found	
LCD		No Local Coverage Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.

Coding/Billing Information

- Note:** 1) This list of codes may not be all-inclusive.
2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Considered Experimental/Investigational/Unproven:

CPT®* Codes	Description
22867	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level

CPT®* Codes	Description
22868	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (List separately in addition to code for primary procedure)
22869	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level
22870	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (List separately in addition to code for primary procedure)

HCPCS Codes	Description
C1821	Interspinous process distraction device (implantable)

*Current Procedural Terminology (CPT®) ©2020 American Medical Association: Chicago, IL.

References

1. Abjornson C, Yoon BV, Callanan T, Shein D, Grinberg S, et al. Spinal Stenosis in the Absence of Spondylolisthesis: Can Interlaminar Stabilization at Single and Multi-levels Provide Sustainable Relief? *Int J Spine Surg.* 2018 Mar 30;12(1):64-69. eCollection 2018 Jan.
2. American Academy of Orthopedic Surgeons. All guidelines. Accessed January 2021. Available at URL address: <http://www.orthoguidelines.org/guidelines>
3. American Academy of Orthopedic Surgeons. Appropriate Use Criteria. Accessed January 2021. Available at URL address: <http://www.orthoguidelines.org/go/auc/>
4. American Academy of Orthopedic Surgeons website. Diseases & Conditions. Lumbar spinal stenosis. Not dated. Accessed January 2021. Available at URL address: <https://orthoinfo.aaos.org/en/diseases--conditions/> <https://orthoinfo.aaos.org/en/diseases--conditions/lumbar-spinal-stenosis/>
5. Badve SA, Kurra S, Geisler FH, Metkar U, Tallarico R, Lavelle W. Nerve Root Sedimentation Sign: Can It Predict the Success for Surgical Intervention in Patients With Symptomatic Lumbar Spinal Stenosis? *Cureus.* 2020 Aug 17;12(8):e9803.
6. Bini W, Miller LE, Block JE. Minimally invasive treatment of moderate lumbar spinal stenosis with the Superion interspinous spacer. *Open Orthop J.* 2011;5:361-7.
7. Block JE, Lavelle WF, Nunley PD. Toward a cure for lumbar spinal stenosis: The potential of interspinous process decompression. *Med Hypotheses.* 2019 Nov;132:109357.
8. Boston Scientific. The Vertiflex™ Procedure. Superion™ Indirect Decompression System. Accessed January 2021. Available at URL address: <https://www.bostonscientific.com/en-US/products/indirect-decompression-system/superion-indirect-decompression-system.html>
9. Cai Y, Luo J, Huang J, Lian C, Zhou H, et al. Interspinous spacers versus posterior lumbar interbody fusion for degenerative lumbar spinal diseases: a meta-analysis of prospective studies. *Int Orthop.* 2016 Jun;40(6):1135-42.
10. Cairns K, Deer T, Sayed D, van Noort K, Liang K. Cost-effectiveness and Safety of Interspinous Process Decompression (Superion). *Pain Med.* 2019 Dec 1;20(Suppl 2):S2-S8.

11. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed January 2021. Available at URL address: https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx?Cntrctr=373&ContrVer=1&CntrctrSelected=373*1&DocType=Active%7cFuture&s=All&bc=AggAAAQAAAA&
12. Centers for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed January 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/nc>
13. Clinicaltrials.gov. accessed January 2021. Available at URL address: <https://www.clinicaltrials.gov/>
14. Davis R, Auerbach JD, Bae H, Errico TJ. Can low-grade spondylolisthesis be effectively treated by either coflex interlaminar stabilization or laminectomy and posterior spinal fusion? Two-year clinical and radiographic results from the randomized, prospective, multicenter US investigational device exemption trial: clinical article. *J Neurosurg Spine*. 2013(b) Aug;19(2):174-84.
15. Davis RJ, Errico TJ, Bae H, Auerbach JD. Decompression and Coflex interlaminar stabilization compared with decompression and instrumented spinal fusion for spinal stenosis and low-grade degenerative spondylolisthesis: two-year results from the prospective, randomized, multicenter, Food and Drug Administration Investigational Device Exemption trial. *Spine (Phila Pa 1976)*. 2013(a) Aug 15;38(18):1529-39.
16. Du MR, Wei FL, Zhu KL, Song RM, Huan Y, Jia B, Gu JT, Pan LX, Zhou HY, Qian JX, Zhou CP. Coflex interspinous process dynamic stabilization for lumbar spinal stenosis: Long-term follow-up. *J Clin Neurosci*. 2020 Nov;81:462-468.
17. Errico TJ, Kamerlink JR, Quirno M, Samani J, Chomiak RJ. Survivorship of coflex Interlaminar-Interspinous Implant. *SAS J*. 2009 Jun 1;3(2):59-67.
18. Falowski SM, Sayed D, Deer TR, Brescacin D, Liang K. Biomechanics and Mechanism of Action of Indirect Lumbar Decompression and the Evolution of a Stand-alone Spinous Process Spacer. *Pain Med*. 2019 Dec 1;20(Suppl 2):S14-S22.
19. Fan Y, Zhu L. Decompression alone versus fusion and Coflex in the treatment of lumbar degenerative disease: A network meta-analysis. *Medicine (Baltimore)*. 2020 Mar;99(11):e19457.
20. Gala R, Russo GS, Whang PG. Interspinous implants to treat spinal stenosis. *Curr Rev Musculoskelet Med*. 2017 Jun;10(2):182-188.
21. Gazzeri R, Galarza M, Neroni M, Fiore C, Faiola A, Puzilli F4, et al. Failure rates and complications of interspinous process decompression devices: a European multicenter study. *Neurosurg Focus*. 2015 Oct;39(4):E14.
22. Golish SR, Groff MW, Araghi A, Inzana JA. Superiority Claims for Spinal Devices: A Systematic Review of Randomized Controlled Trials. *Global Spine J*. 2020 May;10(3):332-345.
23. Guyer R, Musacchio M, Cammisa FP Jr, Lorio MP. ISASS Recommendations/Coverage Criteria for Decompression with Interlaminar Stabilization - Coverage Indications, Limitations, and/or Medical Necessity. *Int J Spine Surg*. 2016 Dec 5;10:41.
24. Hartman J, Granville M, Jacobson RE. The Use of Vertiflex® Interspinous Spacer Device in Patients With Lumbar Spinal Stenosis and Concurrent Medical Comorbidities. *Cureus*. 2019 Aug 12;11(8):e5374.

25. Hayes, Inc. Health Technology Brief. coflex Interlaminar Stabilization Device for Treatment of Lumbar Spinal Stenosis. Hayes, Inc.; Sep 21, 2018. Annual Review Oct 26, 2020.
26. Hayes, Inc. Evolving Evidence Review. Superior Interspinous Spacer System for Treatment of Neurogenic Claudication Caused by Spinal Stenosis. Hayes, Inc.; Oct 06, 2020.
27. International Society for the Advancement of Spine Surgery (ISASS). Policy Statements. Accessed January 2021. Available at URL address: <https://www.isass.org/policy-statements/>
28. Kim DH, Shanti N, Tantorski ME, Shaw JD, Li L, Martha JF, et al. Association between degenerative spondylolisthesis and spinous process fracture after interspinous process spacer surgery. *Spine J.* 2012 Jun;12(6):466-72.
29. Kondrashov DG, Hannibal M, Hsu KY, Zucherman JF. Interspinous process decompression with the X-STOP device for lumbar spinal stenosis: a 4-year follow-up study. *J Spinal Disord Tech.* 2006 Jul;19(5):323-7.
30. Kreiner DS, Shaffer WO, Baisden JL, Gilbert TJ, North American Spine Society, et al. An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spinal stenosis (update). *Spine J.* 2013 Jul;13(7):734-43.
31. Kuchta J, Sobottke R, Eysel P, Simons P. Two-year results of interspinous spacer (X-Stop) implantation in 175 patients with neurologic intermittent claudication due to lumbar spinal stenosis. *Eur Spine J.* 2009 Jun;18(6):823-9.
32. Kumar N, Shah SM, Ng YH, Pannierselvam VK, Dasde S, Shen L. *Asian Spine J.* 2014 Apr;8(2):161-9.
33. Lauryssen C, Jackson RJ, Baron JM, Tallarico RA, Lavelle WF, et al. Stand-alone interspinous spacer versus decompressive laminectomy for treatment of lumbar spinal stenosis. *Expert Rev Med Devices.* 2015;12(6):763-9.
34. Li D, Hai Y, Meng X, Yang J, Yin P. Topping-off surgery vs posterior lumbar interbody fusion for degenerative lumbar disease: a comparative study of clinical efficacy and adjacent segment degeneration. *J Orthop Surg Res.* 2019 Jun 28;14(1):197.
35. Li AM, Li X, Yang Z. Decompression and coflex interlaminar stabilisation compared with conventional surgical procedures for lumbar spinal stenosis: A systematic review and meta-analysis. *Int J Surg.* 2017 Apr;40:60-67.
36. Lønne G, Johnsen LG, Rossvoll I, Andresen H, Storheim K, et al. Minimally invasive decompression versus x-stop in lumbar spinal stenosis: a randomized controlled multicenter study. *Spine (Phila Pa 1976).* 2015 Jan 15;40(2):77-85.
37. Machado GC, Ferreira PH, Harris IA, Pinheiro MB, Koes BW, et al. Effectiveness of surgery for lumbar spinal stenosis: a systematic review and meta-analysis. *PLoS One.* 2015 Mar 30;10(3):e0122800.
38. Machado GC, Ferreira PH, Yoo RI, Harris IA, Pinheiro MB, Koes BW, et al. Surgical options for lumbar spinal stenosis. *Cochrane Database Syst Rev.* 2016 Nov 1;11:CD012421.
39. Matz PG, Meagher RJ, Lamer T, Tontz WL Jr, Annaswamy TM, et al. Guideline summary review: An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spondylolisthesis. *Spine J.* 2016 Mar;16(3):439-48.
40. Merkow J, Varhabhatla N, Manchikanti L, Kaye AD, Urman RD, Yong RJ. Minimally Invasive Lumbar Decompression and Interspinous Process Device for the Management of Symptomatic Lumbar Spinal Stenosis: a Literature Review. *Curr Pain Headache Rep.* 2020 Feb 18;24(4):13.

41. Miller LE, Block JE. Interspinous spacer implant in patients with lumbar spinal stenosis: preliminary results of a multicenter, randomized, controlled trial. *Pain Res Treat.* 2012;2012:823509.
42. Mo Z, Li D, Zhang R, Chang M, Yang B, Tang S. Comparative effectiveness and safety of posterior lumbar interbody fusion, Coflex, Wallis, and X-stop for lumbar degenerative diseases: A systematic review and network meta-analysis. *Clin Neurol Neurosurg.* 2018 Sep;172:74-81.
43. Moojen WA, Arts MP, Jacobs WC, van Zwet EW, van den Akker-van Marle ME, Koes BW, Vleggeert-Lankamp CL, Peul WC; Leiden-The Hague Spine Intervention Prognostic Study Group. Interspinous process device versus standard conventional surgical decompression for lumbar spinal stenosis: randomized controlled trial. *BMJ.* 2013 Nov 14;347:f6415.
44. Musacchio MJ, Laurysen C, Davis RJ, Bae HW, Peloza JH, et al. Evaluation of Decompression and Interlaminar Stabilization Compared with Decompression and Fusion for the Treatment of Lumbar Spinal Stenosis: 5-year Follow-up of a Prospective, Randomized, Controlled Trial. *Int J Spine Surg.* 2016 Jan 26;10:6.
45. Nandakumar A, Clark NA, Smith FW, Wardlaw D. Two-year results of X-stop interspinous implant for the treatment of lumbar spinal stenosis: a prospective study. *J Spinal Disord Tech.* 2013 Feb;26(1):1-7.
46. North American Spine Society (NASS). Coverage Policy Recommendations. Accessed January 2021. Available at URL address: <https://www.spine.org/coverage>
47. North American Spine Society (NASS). Clinical Guidelines. Accessed January 2021. Available at URL address: <https://www.spine.org/researchclinicalcare/qualityimprovement/clinicalguidelines.aspx>
<https://www.spine.org/Research-Clinical-Care/Quality-Improvement/Clinical-Guidelines>
48. National Institute for Health and Clinical Excellence (NICE). Interspinous distraction procedures for lumbar spinal stenosis causing neurogenic claudication. London, UK: NICE; 2010 Nov. Accessed January 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg365>
49. Nunley PD, Patel VV, Orndorff DG, Lavelle WF, Block JE, Geisler FH. Five-year durability of stand-alone interspinous process decompression for lumbar spinal stenosis. *Clin Interv Aging.* 2017 Sep 6;12:1409-1417.
50. Nunley PD, Patel VV, Orndorff DG, Lavelle WF, Block JE, et al. Interspinous Process Decompression Improves Quality of Life in Patients with Lumbar Spinal Stenosis. *Minim Invasive Surg.* 2018 Jul 2;2018:1035954. eCollection 2018a.
51. Nunley PD, Deer TR, Benyamin RM, Staats PS, Block JE. Interspinous process decompression is associated with a reduction in opioid analgesia in patients with lumbar spinal stenosis. *J Pain Res.* 2018 Nov 20;11:2943-2948. eCollection 2018.
52. Park SC, Yoon SH, Hong YP, Kim KJ, Chung SK, et al. Minimum 2-year follow-up result of degenerative spinal stenosis treated with interspinous u (coflex). *J Korean Neurosurg Soc.* 2009 Oct;46(4):292-9.
53. Patel VV, Whang PG, Haley TR, Bradley WD, Nunley PD, et al. Superior interspinous process spacer for intermittent neurogenic claudication secondary to moderate lumbar spinal stenosis: two-year results from a randomized controlled FDA-IDE pivotal trial. *Spine (Phila Pa 1976).* 2015a Mar 1;40(5):275-82. (NCT00692276)
54. Patel VV, Nunley PD, Whang PG, Haley TR, Bradley WD, Davis RP, et al. Superior(®) InterSpinous Spacer for treatment of moderate degenerative lumbar spinal stenosis: durable three-year results of a randomized controlled trial. *J Pain Res.* 2015b Oct 3;8:657-62.

55. Patel VV, Whang PG, Haley TR, Bradley WD, Nunley PD, et al. Two-year clinical outcomes of a multicenter randomized controlled trial comparing two interspinous spacers for treatment of moderate lumbar spinal stenosis. *BMC Musculoskelet Disord*. 2014 Jul 5;15:221. (NCT00692276)
56. Phan K, Rao PJ, Ball JR, Mobbs RJ. Interspinous process spacers versus traditional decompression for lumbar spinal stenosis: systematic review and meta-analysis. *J Spine Surg*. 2016 Mar;2(1):31-40.
57. Pintauro M, Duffy A, Vahedi P, Rymarczuk G, Heller J. Interspinous implants: are the new implants better than the last generation? A review. *Curr Rev Musculoskelet Med*. 2017 Jun;10(2):189-198.
58. Puzzilli F, Gazzeri R, Galarza M, Neroni M, Panagiotopoulos K, et al. Interspinous spacer decompression (X-STOP) for lumbar spinal stenosis and degenerative disk disease: a multicenter study with a minimum 3-year follow-up. *Clin Neurol Neurosurg*. 2014 Sep;124:166-74.
59. Richter A, Halm HFH, Hauck M, Quante M. Two-year follow-up after decompressive surgery with and without implantation of an interspinous device for lumbar spinal stenosis: a prospective controlled study. *J Spinal Disord Tech*. 2014 Aug;27(6):336-41.
60. Schmidt S, Franke J, Rauschmann M, Adelt D, Bonsanto MM, Sola S. Prospective, randomized, multicenter study with 2-year follow-up to compare the performance of decompression with and without interlaminar stabilization. *J Neurosurg Spine*. 2018 Apr;28(4):406-415.
61. Siddiqui M, Smith FW, Wardlaw D. One-year results of X-Stop interspinous implant for the treatment of lumbar spinal stenosis. *Spine*. 2007 May 20;32(12):1345-8.
62. Simon RB, Dowe C, Grinberg S, Cammisa FP Jr, Abjornson C. The 2-Level Experience of Interlaminar Stabilization: 5-Year Follow-Up of a Prospective, Randomized Clinical Experience Compared to Fusion for the Sustainable Management of Spinal Stenosis. *Int J Spine Surg*. 2018 Aug 31;12(4):419-427. eCollection 2018 Aug.
63. Strömquist BH, Berg S, Gerdhem P, Johnsson R, Möller A, et al. X-stop versus decompressive surgery for lumbar neurogenic intermittent claudication: randomized controlled trial with 2-year follow-up. *Spine (Phila Pa 1976)*. 2013 Aug 1;38(17):1436-42.
64. Surgalign Spine Technologies. coflex® Interlaminar Stabilization®. Accessed January 2021. Available at URL address: <https://coflexsolution.com/clinical-research/> <https://coflexsolution.com/> <https://www.surgalign.com/?s=coflex>
65. Tekmyster G, Sayed D, Cairns KD, Raso LJ, Kim C, Block JE. Interspinous Process Decompression With The Superior® Spacer For Lumbar Spinal Stenosis: Real-World Experience From A Device Registry. *Med Devices (Auckl)*. 2019 Oct 3;12:423-427.
66. U. S. Food and Drug Administration (FDA) Center for Devices and Radiological Health. Premarket approval database. X-Stop® Interspinous Process Decompression System. P040001. Decision Date: 11/21/2005. Withdrawal Date: 04/30/2015. Accessed January 2021. Available at URL address: https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma_template.cfm?id=p040001 <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P040001>
67. U. S. Food and Drug Administration (FDA) Center for Devices and Radiological Health. Premarket approval database. Coflex® Interlaminar Technology P110008. Decision Date: 10/17/2012. Accessed January 2021. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf11/P110008A.pdf
68. U. S. Food and Drug Administration (FDA) Center for Devices and Radiological Health. Premarket approval database. PMA Decision date: 05/20/2015. Superior InterSpinous Spacer (VertiFlex®, Inc.)

P140004 Accessed January 2021. Available at URL address:
http://www.accessdata.fda.gov/cdrh_docs/pdf14/P140004a.pdf

69. Wu AM, Zhou Y, Li QL, Wu XL, Jin YL, et al. Interspinous spacer versus traditional decompressive surgery for lumbar spinal stenosis: a systematic review and meta-analysis. *PLoS One*. 2014 May 8;9(5):e97142.
70. Yuan W, Su QJ, Liu T, Yang JC, Kang N, Guan L, et al. Evaluation of Coflex interspinous stabilization following decompression compared with decompression and posterior lumbar interbody fusion for the treatment of lumbar degenerative disease: A minimum 5-year follow up study. *J Clin Neurosci*. 2017 Jan;35:24-29.
71. Zhang JX, Jing XW, Cui P, He X, Hao DJ, Li SJ. Effectiveness of dynamic fixation coflex treatment for degenerative lumbar spinal stenosis. *Exp Ther Med*. 2018;15(1):667-672.
72. Zhao XW, Ma JX, Ma XL, Li F, He WW, Jiang X, et al. Interspinous process devices(IPD) alone versus decompression surgery for lumbar spinal stenosis(LSS): A systematic review and meta-analysis of randomized controlled trials. *Int J Surg*. 2017 Mar;39:57-64.
73. Zhao H, Duan LJ, Gao YS, Yang YD, Zhao DY, et al. Retraction Note: Comparison of two FDA-approved interspinous spacers for treatment of lumbar spinal stenosis: Superior versus X-STOP-a meta-analysis from five randomized controlled trial studies. *J Orthop Surg Res*. 2018 Jun 4;13(1):138.(Retracted article)
74. Zhong J, O'Connell B, Balouch E, Stickley C, Leon C, O'Malley N, Protopsaltis TS, Kim YH, Maglaras C, Buckland AJ. Patient Outcomes After Single Level Coflex ® Interspinous Implants versus Single Level Laminectomy. *Spine (Phila Pa 1976)*. 2020 Dec 31;Publish Ahead of Print.
75. Zucherman JR, Hsu KY, Hartjen CA, Mehalic TF, Implicito DA, Martin MJ, et al. A multicenter, prospective, randomized trial evaluating the X-STOP interspinous process decompression system for the treatment of neurogenic intermittent claudication: two-year follow-up results. *Spine*. 2005 Jun 15;30(12):1351-8.

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.