



Medical Coverage Policy

Effective Date06/15/2025

Next Review Date06/15/2026

Coverage Policy Number..... 0470

Redundant Skin Surgery

Table of Contents

Overview	2
Coverage Policy	2
Health Equity Considerations.....	3
General Background.....	3
Medicare Coverage Determinations	6
Coding Information	6
References.....	7
Revision Details.....	9

Related Coverage Resources

[Bariatric Surgery and Procedures](#)
[Blepharoplasty, Reconstructive Eyelid Surgery, and Brow Lift](#)
[Breast Reconstruction Following Mastectomy or Lumpectomy](#)
[Gender Dysphoria Treatment](#)
[Panniculectomy and Abdominoplasty](#)
[Surgical Treatments for Lymphedema and Lipedema](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted

for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses surgical procedures for the excision of redundant or excessive skin.

Coverage Policy

Coverage for redundant skin surgery varies across plans. Refer to the customer's benefit plan document for coverage details.

If coverage for the specific service is available, the following conditions of coverage apply.

Rhytidectomy, or procedures for excision of redundant or excessive skin of other anatomical areas* (e.g., neck, upper and lower extremities, buttocks), is considered medically necessary when ALL of the following criteria are met:

- There is presence of a functional deficit due to a severe physical deformity or disfigurement resulting from the redundant or excessive skin.
- The surgery is expected to restore or improve the functional deficit.
- The redundant or excessive skin is demonstrated on preoperative photographs.
- The redundant or excessive skin is interfering with activities of daily living.
- For areas other than the face, there is evidence, including photographs, that the redundant or excessive skin is causing persistent intertriginous dermatitis, cellulitis, or skin ulceration, which is refractory to at least three (3) months of medical management, including all applicable treatments. In addition to good hygiene practices, applicable treatment should include topical antifungals; topical and/or systemic corticosteroids; and/or local or systemic antibiotics.

Rhytidectomy, or procedures for excision of redundant or excessive skin of other anatomical areas* (e.g., upper and lower extremities, buttocks), for ANY of the following is considered cosmetic in nature and not medically necessary:

- The surgery is being performed to treat psychological symptomatology or psychosocial complaints, in the absence of significant physical, objective signs.
- The surgery is being performed for the primary purpose of improving appearance.
- The suction-assisted lipectomy is performed alone and not as part of another medically necessary procedure.
- The surgery is for glabellar frown lines.

***Please refer to Cigna Medical Coverage Policies for Panniculectomy and Abdominoplasty, and Blepharoplasty, Reconstructive Eyelid Surgery, and Brow Lift for criteria relating to these surgical procedures. This policy is intended to address redundant skin surgery for anatomical areas not addressed in those Coverage Policies.**

Labiaplasty is considered cosmetic in nature and not medically necessary. Please refer to the Cigna Coverage Policy for Treatment of Gender Dysphoria for information regarding labiaplasty procedure as part of initial gender reassignment surgery.

Note: If the procedure is being performed following significant weight loss, in addition to meeting the criteria noted above, there should be evidence that the individual has maintained a stable weight for at least six months. If the weight loss is the result of bariatric surgery, excision of redundant skin should not be performed until at least 18 months after bariatric surgery and only when the weight has been stable for at least the last six months.

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

General Background

A rhytidectomy is the surgical treatment of wrinkles. It may include removing excess skin, tightening of muscles and redraping of the skin. When the procedure is performed on the facial area, it is also referred to as a facelift. A cervicofacial rhytidectomy involves neck and face components. When it is performed on the forehead area, it may be referred to as a forehead lift or brow lift. (For information regarding redundant skin surgery from the brow area, refer to the Cigna Medical Coverage Policy: Blepharoplasty, Reconstructive Eyelid Surgery, and Brow Lift.) Initially, facelifts consisted mainly of subcutaneous undermining. With increased knowledge of anatomy, the procedure progressed and may involve the underlying structures, including the superficial musculoaponeurotic system (SMAS) and platysma muscle. The SMAS is a fascial layer that underlies the skin in the parotid and cheek areas. The platysma is a large muscle that is located in the subcutaneous plane of the neck, extending from the lower cheek and mandible to the upper chest (Friedman, 2006). It is thought that advancement of the SMAS and platysma muscle will result in an enhanced outcome compared to excisions of skin alone. These procedures are usually performed on an outpatient basis. A cervicoplasty is removal of excess skin from the neck area.

Rhytidectomy procedures and other procedures, including cervicoplasty, that involve the excision of redundant skin and fat are usually performed solely for cosmesis, with the primary purpose being to improve appearance. The most common reason for performing a rhytidectomy procedure is to reverse the signs of aging. There are rare situations where these procedures may be performed to correct a functional impairment and would be considered medically necessary. In the facial area, a functional impairment may be due to facial paralysis or palsy. The functional impairment may involve difficulty with eating, swallowing and achieving oral continence (e.g., an inability to take in food or liquids orally).

Rhytidectomy procedures for glabellar frown lines, or the area that is above the nose and between the eyebrows, are cosmetic in nature. The primary purpose of this procedure is to remove wrinkles, with no functional deficit being corrected.

Removal of excessive or redundant skin may be performed in other areas of the body, including the thigh, leg, hip, buttock, abdomen and arm areas. (For information on redundant skin removal from the abdominal area, refer to the CIGNA Medical Coverage Policy: Panniculectomy and Abdominoplasty.) Depending on the body area, these procedures may be referred to as buttock lift, thigh lift, leg lift, or arm lift (i.e., brachioplasty). These procedures may also involve the removal of subcutaneous tissue, including lipectomy (i.e., removal of fat deposits). Lipectomy may be performed with the open approach or may be performed through suction-assisted procedures or liposuction. This involves the use of liposuction cannula to remove the fat deposits. When suction-assisted lipectomy is performed alone, not as part of another procedure, for the removal of excessive skin, it is considered cosmetic in nature and not medically necessary.

Significant weight loss may result in the presence of redundant skin. This is one of the most common complaints of patients seeking excision of excessive skin and fat. Procedures for removal of redundant skin in the extremities and after weight loss are often performed in conjunction with other procedures, such as an abdominoplasty and/or panniculectomy procedure.

The presence of massive redundant or excessive skin, in rare situations, may result in chronic and persistent local skin conditions in the skin folds. These conditions may include intertrigo, intertriginous dermatitis, cellulitis, ulcerations or tissue necrosis, or they may lead to painful inflammation of the subcutaneous adipose tissue. When these skin conditions are severe, there may be interference with activities of daily living, such as personal hygiene and ambulation. In addition to excellent personal hygiene practices, treatment of these skin conditions generally involves topical or systemic corticosteroids, topical antifungals, and topical or systemic antibiotics.

Labiaplasty

Labiaplasty, or labia reduction (Current Procedural Terminology [CPT] codes 15839 or 56620), is a surgical procedure that removes tissue from the labia, and/or reshapes the labia. The procedure may be performed for asymmetrical, enlarged, or hypertrophic labia minora and/or labia majora. In general, labiaplasty is performed for reduction of labia minora; however, the procedure may also involve labia majora. The procedure is generally cosmetic in nature and is performed to improve appearance. The diagnosis of labia minora hypertrophy is based upon the presence of symptoms and/or distress associated with labia minora that are within the size range considered hypertrophic (Laufer, 2022). There are no standard diagnostic criteria for labia minora hypertrophy. Clinicians generally use labial width (measurement of medial-lateral axis of the labia minora when gently stretched to full width) measurements. There are no standards for normal labial width, and there can be a wide variation in what is considered normal. A stretch width of greater than six cm is generally felt to be consistent with hypertrophy (Laufer, 2022).

In situations where there is discomfort from the condition, the documentation by the provider should include signs of chronic irritation such as skin abrasions, redness, or hyperkeratosis if present. These symptoms can usually be managed with non-surgical measures including: personal hygiene measures (such as use of emollients), supportive garments, arrangement of labia minora during exercise and other activities, and avoidance of form-fitting clothes.

Vulvectomy (CPT code 56620) is a surgical procedure where part or all of the vulva is removed. A simple, complete vulvectomy includes removal of all of the labia majora, labia minora, and clitoris, while a simple, partial vulvectomy may include removal of part or all of the labia majora and labia minora on one side and the clitoris. The removal of vulvar tissue (vulvectomy or partial

vulvectomy) may be considered medically necessary for diagnosis or treatment of benign, premalignant or malignant lesions.

In the absence of the above conditions, vulvar surgery performed to change the appearance or reduce the size of the labia would be considered cosmetic as well as not medically necessary. (For information regarding a labiaplasty procedure as part of initial gender reassignment surgery, refer to the CIGNA Medical Coverage Policy: Treatment of Gender Dysphoria.)

Professional Societies/Organizations

American College of Obstetricians and Gynecologists (ACOG): ACOG published a Committee Opinion for Elective Female Genital Cosmetic Surgery (ACOG, 2023). Recommendations and conclusions include:

- Patients should be made aware that surgery or procedures to alter sexual appearance or function (excluding procedures performed for clinical indications, such as clinically diagnosed female sexual dysfunction, pain with intercourse, interference in athletic activities, previous obstetric or straddle injury, reversing female genital cutting, vaginal prolapse, incontinence, or gender affirmation surgery) are not medically indicated, pose substantial risk, and their safety and effectiveness have not been established.
- Women should be informed about the lack of high-quality data that support the effectiveness of genital cosmetic surgical procedures and counseled about their potential complications, including pain, bleeding, infection, scarring, adhesions, altered sensation, dyspareunia, and need for reoperation.
- Obstetrician–gynecologists should have sufficient training to recognize women with sexual function disorders as well as those with depression, anxiety and other psychiatric conditions. Individuals should be assessed, if indicated, for body dysmorphic disorder. In women who have suspected psychological concerns, a referral for evaluation should occur before considering surgery.
- In responding to a patient’s concern about the appearance of her external genitalia, the obstetrician–gynecologist can reassure her that the size, shape, and color of the external genitalia vary considerably from woman to woman. These variations are further modified by pubertal maturity, aging, anatomic changes resulting from childbirth, and atrophic changes associated with menopause or hypoestrogenism, or both.

American College of Obstetricians and Gynecologists (ACOG): ACOG published a Committee Opinion for Breast and Labial Surgery in Adolescents (ACOG, 2017; 2020; reaffirmed 2024). Recommendations include:

- The obstetrician-gynecologist caring for adolescents should have good working knowledge of nonsurgical alternatives for comfort and appearance as well as knowledge of indications and timing of surgical intervention and referral.
- When adolescents seek medical treatment, the first step is often education and reassurance regarding normal variation in anatomy, growth and development.
- Appropriate patient counseling and assessment of the adolescent’s physical maturity and emotional readiness are necessary before surgical management or referral.
- Individuals should be screened for body dysmorphic disorder. If an obstetrician-gynecologist suspects an adolescent has body dysmorphic disorder, referral to a mental health professional is appropriate.

The opinion also notes that, “When adolescents seek medical treatment, the first step is often education and reassurance regarding normal variation in anatomy, growth, and development. Nonsurgical comfort and cosmetic measures may be offered, including supportive garments, personal hygiene measures [such as use of emollients], arrangement of the labia minora during exercise, and use of formfitting clothing. Surgical correction (labiaplasty) in girls younger than 18 years should be considered only in those with significant congenital malformation, or persistent

symptoms that the physician believes are caused directly by labial anatomy, or both. Physicians should be aware that surgical alteration of the labia that is not necessary to the health of the adolescent, who is younger than 18 years, is a violation of federal criminal law. At least half of the states also have laws criminalizing labiaplasty under certain circumstances, and some of these laws apply to minors and adults. Obstetrician–gynecologists should be aware of federal and state laws that affect this and similar procedures.”

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No National Coverage Determination found	
LCD	Noridian Healthcare Solutions, LLC	Plastic Surgery (L35163 & L37020)	2/14/2025
LCD	Wisconsin Physicians Service Insurance Corporation	Cosmetic and Reconstructive Surgery (L39051)	10/13/2024

Note: Please review the current Medicare Policy for the most up-to-date information.
(NCD = National Coverage Determination; LCD = Local Coverage Determination)

Coding Information

Notes:

1. This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare & Medicaid Services (CMS) code updates may occur more frequently than policy updates.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
15819	Cervicoplasty
15824	Rhytidectomy; forehead
15825	Rhytidectomy; neck with platysmal tightening (platysmal flap, P-flap)
15828	Rhytidectomy; cheek, chin, and neck
15829	Rhytidectomy; superficial musculoaponeurotic system (SMAS) flap
15832	Excision, excessive skin and subcutaneous tissue (includes lipectomy); thigh
15833	Excision, excessive skin and subcutaneous tissue (includes lipectomy); leg
15834	Excision, excessive skin and subcutaneous tissue (includes lipectomy); hip
15835	Excision, excessive skin and subcutaneous tissue (includes lipectomy); buttock
15836	Excision, excessive skin and subcutaneous tissue (includes lipectomy); arm
15837	Excision, excessive skin and subcutaneous tissue (includes lipectomy); forearm or hand
15838	Excision, excessive skin and subcutaneous tissue (includes lipectomy); submental fat pad

15839 [†]	Excision, excessive skin and subcutaneous tissue (includes lipectomy); other area
--------------------	---

[†]Note: Considered cosmetic/not medically necessary when used to report labiaplasty

Suction-assisted lipectomy

Considered Cosmetic/Not Medically Necessary when performed alone and not as part of a medically necessary procedure:

CPT®* Codes	Description
15876	Suction assisted lipectomy; head and neck
15877	Suction assisted lipectomy; trunk
15878	Suction assisted lipectomy; upper extremity
15879	Suction assisted lipectomy; lower extremity

Rhytidectomy for glabellar frown lines:

Considered Cosmetic/Not Medically Necessary:

CPT®* Codes	Description
15826	Rhytidectomy; glabellar frown lines

Labiaplasty

Considered Cosmetic/Not Medically Necessary when used to report labiaplasty:

CPT®* Codes	Description
56620	Vulvectomy, simple; partial

***Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.**

References

1. American College of Obstetricians and Gynecologists (ACOG), Committee Opinion. ACOG Committee Opinion No. 686 Summary: Breast and Labial Surgery in Adolescents. Obstet Gynecol. 2017 Jan;129(1):235. (Reaffirmed 2024). Accessed April 23, 2025. Available at URL address: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2017/01/breast-and-labial-surgery-in-adolescents>
2. American College of Obstetricians and Gynecologists (ACOG), Committee on Gynecologic Practice. ACOG Committee Opinion No. 795. Elective Female Genital Cosmetic Surgery. Jan 2020 (reaffirmed 2023). Accessed April 23, 2025. Available at URL address: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2020/01/elective-female-genital-cosmetic-surgery>
3. American Society of Plastic Surgeons. Surgical Treatment of Skin Redundancy Following Obese and Massive Weight Loss Patients. Jun 2017. Accessed April 23, 2025. Available at

URL address: <https://www.plasticsurgery.org/for-medical-professionals/health-policy/recommended-insurance-coverage-criteria>

4. American Society of Plastic Surgeons. Cosmetic Plastic Surgery. Procedures at a glance. ©2007. Accessed April 23, 2025. Available at URL address: http://www1.plasticsurgery.org/include/pdf/shopping/brochures/Cosmetic_Plastic_Surgery.pdf
5. American Society of Plastic Surgeons (ASPS). Practice Parameter for Surgical Treatment of Skin Redundancy for Obese and Massive Weight Loss Patients. 2017 Jun. Accessed April 26, 2024. Available at URL address: <https://www.plasticsurgery.org/documents/Health-Policy/Guidelines/guideline-2017-skin-redundancy.pdf>
6. Carniol PJ, Ganc DT. Is there an ideal facelift procedure? Curr Opin Otolaryngol Head Neck Surg. 2007 Aug;15(4):244-52.
7. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed April 23, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/reports/local-coverage-proposed-lcds-alphabetical-report.aspx?proposedStatus=A&sortBy=title>
8. Chang P, Salisbury MA, Narsete T, Buckspan R, Derrick D, Ersek RA. Vaginal labiaplasty: defense of the simple "clip and snip" and a new classification system. Aesthetic Plast Surg. 2013 Oct;37(5):887-91.
9. Davison SP. Labiaplasty and Labia Minora Reduction. Emedicine. Updated: Feb 25, 2022. Accessed April 23, 2025. Available at URL address: <http://emedicine.medscape.com/article/1372175-overview>
10. Eremia S, Willoughby MA. Rhytidectomy. Dermatol Clin. 2005 Jul;23(3):415-30.
11. Friedman O. Facelift surgery. Facial Plast Surg. 2006 May;22(2):120-8.
12. Goodman MP, Placik OJ, Benson RH 3rd, Miklos JR, Moore RD, Jason RA, et al. A large multicenter outcome study of female genital plastic surgery. J Sex Med. 2010 Apr;7(4 Pt 1):1565-77.
13. Grossman JA. Thigh and Knee Liposuction. Emedicine. Updated: March 30, 2023. Accessed April 23, 2025. Available at URL address: <http://emedicine.medscape.com/article/1272071-overview>
14. Hunter JG. Labia Minora, Labia Majora, and Clitoral Hood Alteration: Experience-Based Recommendations. Aesthet Surg J. 2016 Jan;36(1):71-9.
15. Iverson RE, Lynch DJ; American Society of Plastic Surgeons Committee on Patient Safety. Practice advisory on liposuction. Plast Reconstr Surg. 2004 Apr 15;113(5):1478-90; discussion 1491-5.
16. Kilpatrick JK, LaFerriere KL. Deep Plane Rhytidectomy. Emedicine. Updated: December 17, 2024. Accessed April 23, 2025. Available at URL address: <http://www.emedicine.com/ent/topic131.htm>

17. Lamont J, Bajzak K, Bouchard C, Burnett M, Byers S, Cohen T, et al. No. 279-Female Sexual Health Consensus Clinical Guidelines. J Obstet Gynaecol Can. 2018 Jun;40(6):e451-e503.
18. Laufer MR, Reddy J. Labia minora hypertrophy. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Literature review current through: Mar 2025. | This topic last updated: May 3, 2024. Accessed on April 23, 2025.
19. Liao LM, Michala L, Creighton SM. Labial surgery for well women: a review of the literature. BJOG. 2010 Jan;117(1):20-5.
20. Oranges CM, Sisti A, Sisti G. Labia minora reduction techniques: a comprehensive literature review. Aesthet Surg J. 2015 May;35(4):419-31.
21. Reddy J, Laufer MR. Hypertrophic labia minora. J Pediatr Adolesc Gynecol. 2010 Feb;23(1):3-6.
22. Reisman NR. Buttocks Contouring. Emedicine. Updated: May 9, 2018. Updated Jan 12, 2023. Accessed April 23, 2025. Available at URL address: <http://emedicine.medscape.com/article/1271806-treatment#showall>
23. Rodriguez-Bruno K, Papel ID. Rhytidectomy: principles and practice emphasizing safety. Facial Plast Surg. 2011 Feb;27(1):98-111.
24. Rohrich RJ, Rios JL, Smith PD, Gutowski KA. Neck rejuvenation revisited. Plast Reconstr Surg. 2006 Oct;118(5):1251-63.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Review	<ul style="list-style-type: none"> No clinical policy statement changes. 	6/15/2025
Annual Review	<ul style="list-style-type: none"> No clinical policy statement changes. 	6/15/2024

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.