



Medical Coverage Policy

Effective Date.....11/15/2021
Next Review Date..... 9/15/2022
Coverage Policy Number 0545

Angioplasty (Extracranial, Intracranial) and Intracranial Aneurysm Repair

Table of Contents

| | |
|--|----|
| Overview | 1 |
| Coverage Policy | 1 |
| Angioplasty (Extracranial, Intracranial) | 2 |
| Intracranial Aneurysm Repair | 3 |
| General Background | 4 |
| Angioplasty (Extracranial, Intracranial) | 4 |
| Intracranial Aneurysm Repair | 8 |
| Medicare Coverage Determinations | 15 |
| Coding/Billing Information | 15 |
| References | 17 |

Related Coverage Resources

[Venous Angioplasty and/or Stent Placement in Adults](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses endovascular treatment of intracranial aneurysm and atherosclerosis (intracranial, extracranial), using angioplasty, with or without stent placement. Angioplasty is a minimally invasive procedure performed to restore blood flow through a blocked or narrowed artery. It is an alternative to carotid endarterectomy. The coverage criteria in this Medical Coverage Policy are primarily based on recommendations from published practice parameters, recommendations and professional society/organization consensus guidelines.

Carotid angioplasty and stenting is usually safely performed as an ambulatory/outpatient procedure (i.e., less than twenty four hour stay). When clinically appropriate, however, some individuals undergoing carotid artery angioplasty and stenting may require an inpatient hospital stay greater than twenty four hours.

Coverage Policy

Angioplasty (Extracranial, Intracranial)

Extracranial carotid artery angioplasty with stent placement as treatment for carotid artery stenosis is considered medically necessary when EITHER of the following criteria is met:

- The individual is or has been symptomatic within the previous six months (i.e., stroke or transient ischemic attack [TIA]), has 50-99% stenosis of the common or internal carotid artery, and is considered high risk for carotid endarterectomy because of the presence of ANY of the following comorbidities or anatomic features:
 - knowledge of two or more proximal or major diseased coronary arteries with $\geq 70\%$ stenosis that have not or cannot be revascularized
 - currently on a list for major organ transplantation (i.e., heart, lung, liver, kidney) or is being evaluated for such
 - left ventricular ejection fraction $< 30\%$
 - New York Heart Association (NYHA) Functional Class III or higher congestive heart failure
 - uncontrolled diabetes defined as fasting glucose > 400 mg/dl and ketones $> 2+$
 - restenosis after previous carotid artery endarterectomy (CEA) or stenting
 - patient is status/post-radiation treatment to the neck
 - patient is status/post-radical neck surgery
 - surgically inaccessible lesions (e.g., lesions above the level of C2 or below the clavicle, lesions obstructed by tumors in the neck)
 - spinal immobility (i.e., inability to flex neck beyond neutral or kyphotic deformity)
 - contralateral laryngeal nerve paralysis
 - age ≥ 80 years
 - severe chronic obstructive pulmonary disease (e.g., FEV less than 50% predicted)
 - coronary artery bypass within six weeks
 - tracheostomy
 - recent myocardial infarction (≤ 30 days)
 - recent coronary artery bypass grafting or valve repair
 - high or low placed carotid lesions
 - dialysis-dependent renal failure
 - unstable angina defined as rest angina with electrocardiogram (ECG) changes
 - contralateral internal carotid artery occlusion

- The individual is asymptomatic, has 60-99% stenosis of the common or internal carotid artery, and ANY of the following imaging characteristics or other findings suggestive of increased stroke risk while on medical therapy:
 - stenosis progression
 - silent infarction on computerized tomography (CT)
 - large plaque area
 - plaque echolucency
 - intra-plaque hemorrhage on magnetic resonance imaging (MRI)
 - impaired cerebral vascular reserve
 - spontaneous embolization on transcranial Doppler
 - history of contralateral TIA

Extracranial carotid artery angioplasty with stent placement is considered not medically necessary for ANY other indication, including the following:

- carotid stenosis with angiographically visible intraluminal thrombus
- total vessel occlusion
- the stenosis cannot be safely reached or accessed by endovascular approach

Inpatient Admission

When the above medical necessity criteria has been met for extracranial carotid artery angioplasty and stenting, an inpatient admission is considered medically necessary when EITHER of the following criteria is met:

- **The individual is considered to be high risk for peri or post-procedural complications resulting from one of the following significant comorbidities:**
 - poorly controlled diabetes mellitus
 - severe pulmonary disease
 - end stage renal disease
 - New York Heart Association (NYHA) Functional Class III or higher congestive heart failure
 - severe carotid tortuosity
 - contralateral carotid stenosis > 50%
 - age ≥ 80 years
 - multiple significant co-morbidities
 - carotid artery pseudoaneurysm
 - presence of moderate or severe aortic stenosis
 - severe plaque calcification, circumferential carotid plaque
 - heavily calcified aortic arch
 - near occlusion of the carotid artery (i.e., string sign)
 - ulcerated plaque
 - ≥ 7 age-related white matter changes on magnetic resonance imaging
 - carotid arteriovenous (AV) fistula formation
 - oral anticoagulant therapy (i.e., warfarin, direct oral anticoagulants [DOACs])

- **Occurrence of ANY of the following intra or post-procedural complications:**
 - hemodynamic instability (e.g., tachycardia, hypotension, inadequate perfusion)
 - presence of neurological deficit (e.g., difficulty swallowing, facial drooping, stroke)
 - bleeding or hematoma at the operative site
 - myocardial compromise (e.g., ischemia, infarct)
 - inability to deploy a cerebral protection device
 - intravascular procedural complications such as acute vessel closure, stent malposition, or vessel dissection
 - carotid artery stent fracture
 - inability to use a closure device
 - cerebral hyperperfusion syndrome

Extracranial vertebral artery angioplasty with stent placement is considered medically necessary when BOTH the following criteria are met:

- recurrent vertebrobasilar territory symptoms refractory to maximum medical management
- 50-99% extracranial vertebral artery stenosis

Extracranial vertebral artery angioplasty with stent placement is considered experimental, investigational or unproven for treatment of ANY other indication, including asymptomatic vertebral artery stenosis.

Intracranial angioplasty, with stent placement, for treatment of atherosclerosis of intracranial arteries is considered experimental, investigational or unproven.

Intracranial Aneurysm Repair

Either of the following is considered medically necessary for repair of an intracranial aneurysm:

- surgical clipping
- endovascular treatment using either embolization coiling or balloon–assisted coiling

Intracranial Stent Placement

Intracranial stent placement (e.g., stent assisted coil, endoluminal flow diversion) using a stent approved by the U.S. Food and Drug Administration for the intended use, is considered medically necessary for treatment of a wide-necked intracranial aneurysm (e.g., neck diameter of > 4mm or a dome-to-neck ratio of < 2) located in the internal carotid artery when the aneurysm is not amenable to surgical or standard endovascular treatment (e.g., clipping, coil embolization).

Intrasaccular Flow Disruption

Endovascular repair of an intracranial aneurysm using an intrasaccular flow-disruption device (e.g., Woven EndoBridge [WEB] Aneurysm Embolization System [MicroVention, Inc.]) is considered experimental, investigational or unproven.

General Background

Atherosclerosis is the buildup of plaque inside arteries. When plaque buildup occurs in the extracranial or intracranial arteries, the vessel becomes blocked or severely narrowed and may lead to stroke and associated neurologic impairment. Risk factors associated with extracranial and intracranial atherosclerosis are generally the same as those for atherosclerotic disease that occurs in other areas of the body.

Stroke is the fifth leading cause of death and leading cause of disability in the United States; on average the incidence is 800,000 cases/year with a majority resulting from acute ischemic stroke (AIS) from cerebrovascular occlusion (Eskey, et al., 2018). Intravenous thrombolytic therapy with recombinant tissue plasminogen activator (tPA) is a well-established treatment of stroke and associated with improved clinical outcomes when administered within 3-4.5 hours of the onset of symptoms.

Carotid endarterectomy is an established treatment for symptomatic carotid artery stenosis. Symptomatic is defined as focal neurologic symptoms caused by ischemic stroke in the carotid artery territory or TIAs, and ipsilateral to significant carotid atherosclerotic pathology. Endovascular treatment may be performed as a viable alternative for some patients.

Angioplasty (Extracranial, Intracranial)

Carotid and cerebral angiography are considered the gold standard for imaging vasculature in the head and neck region and can identify arterial dissections, arteriovenous malformations or fistulas, intracranial occlusive disease, and traumatic vascular injuries. Angiography is the radiographic visualization of blood vessels following injection of a radiopaque substance. Other techniques such as duplex ultrasonography, computed tomography (CT) angiography and magnetic resonance (MR) imaging angiography may also be used to determine the degree of stenosis.

Carotid Artery: The common carotid arteries are each located on the side of the neck and supply blood to the brain, neck and face. Each common artery branch has two divisions: the internal carotid and external carotid. Treatment for stenosis occurring in the carotid artery depends on the degree of blockage and the presence of symptoms. Asymptomatic patients (no history of ischemic stroke or TIA) with stenosis are treated medically with antiplatelet therapy (e.g., aspirin) to decrease the likelihood of a blood clot and decrease the risk of stroke. Patients with severe symptomatic stenosis are referred for surgery. Carotid artery endarterectomy (CEA) is considered the treatment of choice for significant blockage or stenosis that occurs in the carotid arteries and is symptomatic. CEA has been shown to reduce risk of stroke in both asymptomatic and symptomatic patients in

several randomized controlled trials (RCTs) (Brothers, et al., 2015). However, CEA has been associated with an increased risk of morbidity and mortality resulting in the presence of significant comorbidities and anatomic factors, resulting in both neurological and non-neurological complications (Brott, et al., 2013).

For individuals who meet criteria for revascularization, carotid artery angioplasty and stenting (CAS) may be considered an alternative (Burton and Lindsay, 2005) to CEA. In contrast to CEA, angioplasty of the carotid artery is not associated with potential for cranial nerve injury and can be performed under local anesthesia. This procedure is a minimally invasive endovascular procedure performed to restore blood flow through a narrowed artery, reduce the chance of embolization, and prevent stenosis and stroke. Stent implantation involves the permanent placement of a small mesh tube within the narrowed artery to compress the obstructive material and maintain patency of the artery, therefore restoring blood flow. The balloon is then deflated and the catheter is removed. In some circumstances, an embolic protection device may be inserted to accompany stent placement. The embolic protection device typically consists of a small wire mesh or basket that is used to capture any embolic debris that may dislodge from the lesion in order to prevent the debris from reaching the brain or other intracranial areas. Such devices are purported to further decrease the neurologic event risk from carotid angioplasty and similar procedures. Selection of device depends on the lesion characteristics and anatomic considerations (Ricotta, et al., 2011).

Similar to CEA, CAS is associated with risks. Risk stratification for CAS is similarly divided into two categories: anatomic (e.g., lesion location, lesion type) and physiologic characteristics (e.g., comorbid conditions). CAS has been recommended for patients who have high perioperative coronary risk or anatomic risk for CEA (Murad, et al., 2011; Gurm, et al., 2008a). For normal risk patients, studies suggest CEA has resulted in lower perioperative and longer term stroke and death rates when compared with carotid artery stenting (Brothers, et al., 2015). A meta-analysis published by Murad and colleagues (2011) evaluated efficacy and safety of CEA versus CAS in subjects with carotid artery disease. Thirteen RCTs were included in the review, 7,484 subjects in total, and 80% were symptomatic. Compared with CEA, CAS was associated with an increased risk of any stroke, decreased risk of perioperative myocardial infarction (MI), and nonsignificant increase in mortality. When the analysis was limited to the most recent RCTs published, which were described by the authors as having better methodology and 56% of the total cohort (two RCTs), results demonstrated CAS was associated with significant increased risk of any stroke and mortality, and a nonsignificant decrease for risk of MI.

High Risk Indicators: Patients at high risk for CEA are defined as having significant comorbidities and/or anatomic risk factors who would be poor candidates for CEA in the opinion of a surgeon (CMS, 2005). Comorbid conditions and/or anatomic features that should be considered prior to selection of endovascular management can be found in each of the manufacturers' U.S. Food and Drug Administration (FDA) Summary of Safety and Effectiveness Data for their CAS system. These are consistent with the inclusion and exclusion criteria for subjects in the CAS trials and studies (e.g., ARCHER, CABERNET, SAPPHERE trials). For example, in order for patients to qualify as a high-risk or nonsurgical candidate in any of the three ARCHER trials, two or more of the criteria listed in a-e OR one or more of the criteria listed in f-q had to be met:

- a) knowledge of two or more proximal or major diseased coronary arteries with $\geq 70\%$ stenosis that have not or cannot be revascularized
- b) unstable angina defined as rest angina with electrocardiogram (ECG) changes
- c) MI within the previous 30 days and current need for carotid artery revascularization
- d) concurrent requirement for aortocoronary bypass or cardiac valve surgery within 30 days
- e) contralateral occlusion of the internal carotid artery
- f) currently on a list for major organ transplantation (i.e., heart, lung, liver, kidney) or is being evaluated for such
- g) ejection fraction $< 30\%$ or New York Heart Association (NYHA) Functional Class III or higher
- h) $FEV_1 < 30\%$ (predicted)
- i) dialysis-dependent renal failure
- j) uncontrolled diabetes defined as fasting glucose > 400 mg/dl and ketones $> 2+$
- k) restenosis after previous CEA
- l) patient is status/post-radiation treatment to the neck
- m) patient is status/post-radical neck surgery

- n) surgically inaccessible lesions (e.g., lesions above the level of C2 or below the clavicle, lesions obstructed by tumors in the neck)
- o) spinal immobility (i.e., inability to flex neck beyond neutral or kyphotic deformity)
- p) presence of tracheostomy stoma
- q) contralateral laryngeal nerve paralysis

In addition, published guidelines (Brott, et al., 2013), textbooks and various other sources indicate that for the management of individuals with extracranial and vertebral artery disease, although no adequate studies have validated high risk criteria, generally accepted criteria for high risk includes at least one of the following (based on SAPHIRE 2004, SAPHIRE 2008, and CREST 2010 trials):

- clinically significant cardiac disease (congestive heart failure, abnormal stress test, or need for open heart surgery)
- severe pulmonary disease
- contralateral carotid occlusion
- contralateral laryngeal nerve palsy
- previous radical neck surgery or radiation therapy to the neck
- recurrent stenosis after endarterectomy
- age \geq 80 years
- chronic obstructive pulmonary disease
- prior CEA or CAS
- prior coronary artery bypass surgery

In 2018, Naylor and colleagues reported within the European Vascular Surgery Clinical Practice Guidelines that data is conflicting, and there is no general consensus regarding how ‘high risk’ is defined. These authors recommend management decisions based on an individual patient basis with consideration of comorbidities as well as anatomical features and the experience of the CAS surgeon (Naylor, et al., 2018).

Risks associated with angioplasty include restenosis (although uncommon) after implantation of the stent, non-neurologic complications during the procedure (e.g., hemodynamic instability), and complications resulting from embolic debris that become dislodged from the site of the lesion either during or after the procedure which may lead to stroke or death. Stroke following CAS may be the result of embolization at the time of the procedure, delayed embolization, or hemorrhagic stroke related to hyperperfusion (Gurm, et al., 2008a). In general however, the overall postoperative neurologic complication rates for angioplasty and stenting of the extracranial carotids for the treatment of stenosis range from 0% to 10%.

Absolute contraindications to CAS include carotid stenosis with an angiographically visible intraluminal thrombus, complete vessel occlusion, endovascular inaccessible stenosis, or other significant contraindications for angiography.

Vertebral Artery: The vertebral arteries are located on each side of the neck, rise from the subclavian artery, branches into four segments (three are extracranial and one is intracranial) as it rises superiorly, and then joins to form the basilar artery at the skull. The basilar artery carries blood to the brain. Atherosclerosis of the vertebral or basilar arteries accounts for 20-25% of strokes (Naylor, et al., 2018).

Non-invasive imaging using contrast enhanced magnetic resonance angiography or computed tomography is recommended for diagnosing vertebral artery disease due to the risk for angiography-related stroke (Naylor, et al., 2018). The risk of stroke remains high in symptomatic patients despite best medical management; in asymptomatic patients risk is much lower. Padalia and colleagues (2018) reported that for symptomatic patients according to the WASID (Warfarin Aspirin Symptomatic Intracranial Disease) trial, one year risk of ischemic stroke in the territory of a 50% intracranial stenosis was about 11-12% on high dose aspirin and warfarin. The rate increased to 23% when the degree of stenosis was 70% (Padalia, et al., 2018).

Symptomatic blockage that occurs within the intracranial arteries is standardly treated with medical therapy. According to the 2012 Society of Neuro Interventional Surgery (Hussain, et al., 2012), medical management of

symptomatic intracranial atherosclerotic disease includes aspirin and clopidogrel for three months with aggressive risk factor modification (i.e., hypertension, hyperlipidemia, diabetes and smoking cessation) as first line therapy. Angioplasty and stenting may be recommended for patients who remain symptomatic despite maximal medical management; however, anatomically these arteries are more difficult to access compared to the extracranial arteries. Evidence in the peer-reviewed published scientific literature has demonstrated a high risk of ischemic stroke with intracranial vertebral artery stenting, and treatment is reserved for individuals with stenosis who are hemodynamically unstable and are refractory to maximal medical management (Padalia, et al., 2018).

Recommendations and clinical practice guidelines from professional societies for intracranial angioplasty and stenting vary. Brott et al. (2013) published guidelines on the management of patients with extracranial carotid and vertebral artery disease and reported that randomized controlled trials are lacking for vertebral artery stenting, and there is insufficient evidence to support endovascular management is superior to best medical management for treatment of vertebral artery stenosis. In 2012, Hussain and associates published a standard of practice for endovascular treatment of intracranial atherosclerosis. Within this document, the authors report endovascular angioplasty with or without stenting is a possible therapeutic option for selected individuals with symptomatic intracranial arterial disease, defined as subjects with symptomatic 70-99% intracranial stenosis when aggressive maximal medical therapy has failed (Hussain, et al., 2012).

The American Heart Association Stroke Council, Council on Cardiovascular Nursing, Council on Peripheral Vascular Disease and Council on Clinical Cardiology published guidelines for early management of patients with acute ischemic stroke (Jauch, et al., 2013). Within these guidelines, the authors acknowledge the following:

- Emergent intracranial angioplasty and/or stenting is not well established and should only be used in a clinical trial setting (Class IIb, Level of Evidence C)
- Emergent angioplasty and/or stenting of the extracranial carotid or vertebral arteries in unselected patients is not well established (Class IIb, Level of Evidence C). Use of these techniques may be considered in the presence of acute ischemic stroke resulting from cervical atherosclerosis or dissection (Class IIb, Level of Evidence C).

Within these guidelines Class IIb evidence is defined as: Benefit \geq Risk, Additional studies with broad objectives are needed, additional registry data would be helpful; Procedure /Treatment may be considered.

Level of evidence "C" is defined as: Very limited populations evaluated (Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as sex, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use), Only consensus opinion of experts, case studies or standard of care.

The European Society for Vascular Surgery supports extracranial angioplasty and stenting for patients with recurrent vertebrobasilar symptoms (despite best medical management) and who have 50-99% extracranial vertebral artery stenosis (Naylor, et al., 2018). The evidence is insufficient to support safety and efficacy of intracranial angioplasty and stenting.

A scientific statement from the American Heart Association (Eskey, et al., 2018) supports intracranial stenting with the Wingspan or Pharos stent system for individuals with 70-99% stenosis when there is progressing symptoms, recurrent TIA or stroke despite medical management with dual antiplatelet therapy, systolic BP of < 140, and high intensity statin therapy. These criteria are in accordance with the FDA Humanitarian Device Exemption (HDE) approval. Within this guideline, the authors note in detail the SAMMPRIS (Chimowitz, et al., 2011) and VISSIT (Zaidat, et al., 2015) trials; both trials were halted early because the 30-day endpoint of stroke, death or intracranial hemorrhage occurred in more patients who underwent stenting than medical management. In addition, at one year there was an increased risk of stroke or TIA in the stent group.

U.S. Food and Drug Administration (FDA)

The FDA has listed numerous stents and stent systems that have received Premarket Approval (PMA) for treatment of carotid stenosis and embolic protection devices from various manufacturers. The first carotid stenting system was approved in 2004 (i.e., ACCULINK™ Carotid Stent System and the RX ACCULINK™ Carotid Stent System [Guidant Corporation, Santa Clara, CA] used in conjunction with carotid embolic protection

systems [ACCUNET™ and RX ACCUNET™ Embolic Protection Systems, Guidant Corporation, Santa Clara, CA]); however, several stenting systems have been approved since then. The FDA-approved stents and distal embolic protection devices differ in the deployment methods once they reach the targeted lesion. The FDA-approved stents and distal embolic protection devices were initially approved based on either RCTs (i.e., Precise and AngioGuard), or the devices were approved based on uncontrolled trials, single-arm trials or registries, and comparison to historical controls. Additional stents and embolic protection devices have since been approved, and various PMA supplements have been approved since the initial approval on most devices.

Contraindications for each CAS system and distal embolic protection device are included in the FDA Summary of Safety and Effectiveness Data. These include but are not limited to the following:

- contraindication to anticoagulant and/or antiplatelet therapy
- severe vascular tortuosity or anatomy that would preclude the safe introduction of a guide catheter, sheath, embolic protection system, or stent system
- known hypersensitivity to nickel-titanium
- uncorrected bleeding disorders
- lesions in the ostium of the common carotid artery

Literature Review

Early studies were conflicting and did not show a benefit of CAS compared with CEA for reduction of stroke or death in the treatment of atherosclerosis of the carotid arteries. However, there is a growing body of evidence to support CAS is not inferior to CEA. A number of randomized clinical trials, nonrandomized trials, systematic reviews and meta-analyses support safety and efficacy of carotid artery angioplasty and stenting for select individuals requiring revascularization (Murad, et al., 2011; Yavin, et al., 2011; Meier, et al., 2010; Eckstein, et al., 2008; Eisenhauer and White, 2008; Gurm, et al., 2008a; Gurm, et al., 2008b; Iyer, et al., 2008; Gray, et al., 2007; Katzen, et al., 2007; White, et al., 2006; Burton and Lindsay, 2005; Groschel, et al., 2005; Zahn, et al., 2005). The preponderance of the evidence supports the conclusion that CAS with embolic protection is not inferior to CEA in either symptomatic or asymptomatic patients at increased risk for surgical complications of CEA (White, et al., 2008). Individuals with surgical high risk features (anatomic and comorbid) for CEA have been proven to have outcomes similar to CEA; this was reported in a randomized controlled trial (SAPPHIRE). The SAPPHIRE study is an accepted study by the FDA as evidence supporting CAS device approval (Yadav, 2004). Three year outcomes data from the SAPPHIRE study have confirmed the long-term safety and efficacy in this high risk subset of patients. The author reported that 73.8% of patients in the stenting group and 69.7% in the endarterectomy group were free of major adverse events at three years (the pre-specified major end point, defined as death, myocardial infarction, or stroke within 30 days or death or ipsilateral stroke between 31 days and 1,080 days). A total of 80.0% of patients in the stenting group and 75.8% in the endarterectomy group were alive at three years. A total of 92.0% of patients in the stenting group and 93.3% in the endarterectomy group were free of stroke at three years (defined as stroke within 30 days or ipsilateral stroke between 31 days and 1,080 days) (Gurm, et al., 2008b).

The medical literature does not lend firm support to treatment of asymptomatic candidates. For individuals with asymptomatic disease, the American Heart Association/American Stroke Association (AHA/ASA) has noted that advances in optimal medical therapy have resulted in uncertainty about the need for, and benefit of, either treatment modality in the asymptomatic subgroup with carotid artery stenosis. The recommendations for selection of asymptomatic patients for carotid revascularization indicate decisions should be guided by an assessment of comorbid conditions and life expectancy, as well as other individual factors and a thorough discussion regarding risks and benefits of the procedure. The authors concluded that more data is needed to compare long-term outcomes following carotid artery endarterectomy and angioplasty in asymptomatic individuals with carotid artery stenosis (Goldstein, et al., 2011).

The updated Society for Vascular Surgery Guidelines for Management of Extracranial Carotid Disease (Ricotta, et al., 2011), the AHA/ASA guidance regarding asymptomatic disease (Goldstein, et al., 2011), and the National Institute for Health and Clinical Excellence (NICE, 2011a) agree the current evidence for safety of carotid artery stent placement in asymptomatic carotid stenosis shows well-documented risk, particularly for stroke.

Intracranial Aneurysm Repair

Clipping, Coiling, Stenting, Endoluminal Flow Diversion: Aneurysms occur when there is a weakened area within the arterial wall and the artery becomes distended, often resulting from atherosclerosis, trauma, infection, or from other medical conditions. They may occur anywhere anatomically, but most commonly occur in the aorta or brain (i.e., cerebral). Cerebral aneurysms are associated with morbidity and mortality resulting from subarachnoid hemorrhage following rupture. Treatment is indicated for ruptured and nonruptured cerebral aneurysms and may include either clipping, balloon-assisted coiling, stent-assisted coiling, or flow diversion (Eskey, et al., 2018). Clipping involves placing an implantable clip over the neck of the aneurysm, isolating it from circulation. Coiling involves endovascular placement of embolization coils into the aneurysm sac to exclude it from the circulation; however, the coil may protrude into the parent artery. More recently, stent-assisted coiling and the use of flow-diverting stents have been investigated as an alternative for patients with cerebral aneurysms whose anatomy is not amenable to simple coiling, or for treatment of a wide-neck aneurysm. Wide-neck aneurysms are defined as having a neck of 4mm or a dome-to-neck ratio of < 2 (FDA). Stent-assisted coiling or endoluminal flow diversion using flow diverting stents is considered an alternative treatment only in select cases and in accordance with FDA indications (Eskey, et al., 2018).

U.S. Food and Drug Administration:

Two intracranial stents have received approval from the FDA through the humanitarian device exemption (HDE) program for treatment of intracranial aneurysms. The Neuroform™ Microdelivery Stent System (Stryker, Kalamazoo, MI) received approval in 2002 for use with embolic coils for treatment of wide-neck intracranial aneurysms that cannot be treated by surgical clipping (FDA, H020002). A second device, the Enterprise™ Vascular Reconstruction Device and Delivery System (Cordis Neurovascular Inc., Miami Lakes, FL) received approval in 2007 and is intended for use with embolic coils for treatment of wide-neck, intracranial, saccular or fusiform aneurysms (FDA, H060001).

A flow diversion device, the Pipeline® Embolization Device (Micro Therapeutics, Irvine, CA), received premarket approval (PMA) from the U.S. Food and Drug Administration (FDA, P100018) in 2011 for endovascular treatment of large or giant wide-necked intracranial aneurysms in the intracranial artery. It is a braided multi-alloy mesh cylinder designed for placement in the neck of an intracranial aneurysm. Additional flow diversion devices that have received FDA PMA approval for treatment of wide-necked intracranial aneurysms include Surpass Streamline Flow Diverter (Stryker Neurovascular, Fremont, CA), Flow Re-Direction Endoluminal Device (FRED) System (Microvention, Inc, Aliso Viejo, CA) and an expanded approval in 2019 for the Pipeline Flex device. The FDA recalled the Pipeline Flex Embolization Device on September 20, 2021 due to a risk of the delivery system's wire and tubes fracturing and breaking off when the system is being used to place, retrieve or move the stent inside the patient. Information available at clinicaltrials.gov indicate another device, the SILK Vista Baby Flow Device (Balt Extrusion, Montmorency France) is currently being investigated although an FDA approval was not found on the FDA site.

According to the PMA labeling indications, the Pipeline® Embolization Device is indicated for the endovascular treatment of adults (age 22 and above) with large or giant wide-necked intracranial aneurysms in the internal carotid artery from the petrous to the superior hypophyseal segments (FDA, PMA P100018A). In the clinical trials for the PMA, inclusion criteria were individuals with a single target intracranial aneurysm located on the petrous, cavernous, or paraophthalmic region of the internal carotid artery (n=108); a neck of the target aneurysm of ≥ 4 mm or no discernible neck, and a maximum fundus diameter ≥ 10 mm. The parent artery diameter was defined as 2.5 to 5.0 mm distal and proximal to the target aneurysm. Subjects were not enrolled if they had a stenosis of the extracranial carotid artery or of the IA parent artery of $> 50\%$, or if they had a subarachnoid hemorrhage within the prior 60 days, or intracerebral hemorrhage or major surgery in the preceding 42 days. The expanded approval for the Pipeline Flex device indicates the device is intended for patients with small or medium, wide-necked brain aneurysms in the territory from the petrous to the terminus of the internal carotid artery (ICA).

According to the FDA labeling, the Surpass Streamline Flow Diverter (Stryker Neurovascular, Fremont, CA) device is indicated for use in the endovascular treatment of patients (18 years of age and older) with unruptured large or giant saccular wide-neck (neck width ≥ 4 mm or dome-to-neck ratio < 2) or fusiform intracranial aneurysms in the internal carotid artery from the petrous segment to the terminus arising from a parent vessel with a diameter ≥ 2.5 mm and ≤ 5.3 mm (FDA, PMA P170024).

FDA labeling information for the Flow Re-Direction Endoluminal Device (FRED®) System indicates the intended use of this device is for the internal carotid artery from the petrous segment to the terminus, for the endovascular treatment of adult patients (22 years of age or older) with wide-necked (neck width \geq 4 mm or dome-to-neck ratio $<$ 2) saccular or fusiform intracranial aneurysms arising from a parent vessel with a diameter \geq 2.0 mm and \leq 5.0 mm (FDA, PMA 180027).

Literature Review: Evidence in the peer-reviewed scientific literature and professional society recommendations tend to support clinical safety and efficacy for the use of flow diversion devices as treatment of an intracranial aneurysm, as an alternative to coil embolization, consistent with FDA indications when there is a wide-neck bifurcation aneurysm (4 mm or more) or when the sack-to-neck ratio is less than 2:1 (Eskey, et al., 2018; Beckse, et al., 2017; Kallmes, et al., 2017; Kallmes, et al., 2015; Thompson, et al., 2015). Endoluminal flow diversion is considered a reasonable option for patients with intracranial aneurysms that, due to their morphology, complexity, or location, are not amenable to treatment with other endovascular techniques or standard surgery, or for patients who are not appropriate candidates for open surgery due to comorbidity.

Hayes published a review for the use of the Pipeline Embolization Device (PED) as an endovascular treatment for adult patients with intracranial aneurysms not amenable to standard therapies and who do not have acute subarachnoid hemorrhage or other contraindications to treatment, and noted that there is a large, low-quality body of evidence that PED offers benefit in the selected patient population, with generally acceptable rates of occlusion and neurologic morbidity and mortality. Although ample data provide estimates of occlusion and overall neurologic morbidity and mortality rates associated with flow diversion devices, comparative data are insufficient to draw conclusions regarding whether these outcomes are higher, lower, or the same as standard interventions. Hayes noted a concern for patients whose aneurysm characteristics are amenable to treatment by standard therapies (Hayes, 2020).

Intrasaccular Flow Disruption: Intrasaccular flow diversion devices are currently under investigation for treatment of a bifurcating intracranial aneurysm. A bifurcating aneurysm is defined as a weakness located at a junction between two diverging vessels. This device is used to block the flow of blood into the aneurysm, to reduce the chance of it rupturing, or to stop further bleeding from an aneurysm that has already ruptured. Antiplatelet premedication is not required for WEB aneurysm treatment. During the procedure, a catheter is inserted into the femoral artery and then advanced into the cerebral circulation under X-ray guidance. A second, smaller catheter is then placed inside the first and is inserted into the aneurysm. A basket-like device made of fine wire mesh is then pushed through the second catheter and placed into the aneurysm sac. The mesh device covers the aneurysm neck and obstructs blood flow into the aneurysm sac, creating blood stasis and promoting endothelial growth across the neck of the aneurysm. The appropriate device size is selected according to the aneurysm width and height.

U.S. Food and Drug Administration:

One device that recently received FDA PMA approval is the Woven Endobridge (WEB) Aneurysm Embolization System (MicroVention, Inc.). The device is intended for use at the middle cerebral artery (MCA) bifurcation, internal carotid artery (ICA) terminus, anterior communicating artery (AComm) complex, or basilar artery apex for the endovascular treatment of adult patients with saccular, wide neck, bifurcating intracranial aneurysms with dome diameter from 3 mm to 10 mm, and either neck size 4 mm or greater, or the dome-to-neck ratio is greater than 1 and less than 2. According to the FDA, one-year follow-up data from the WEB Intrasaccular Therapy Study (WEB-IT) was used to support the approval and a PMA five year post approval study is required by the FDA. The WEB-IT study (Arthur, et al., 2019) is a prospective, multi-center non-randomized pivotal study that was conducted under IDE G130286 and was initiated prior to device approval.

Literature review: In 2019, Arthur and colleagues published the one year outcomes of the WEB-IT study, a prospective, multicenter, single-arm, interventional study conducted at 21 United States and six international centers. The study enrolled 150 adults with a wide-necked bifurcating aneurysm of the anterior and posterior intracranial circulation. Clinical follow-up occurred at 30 days, six months and one year, and is scheduled to occur every year for five years total. The one year primary endpoint is the proportion of patients with complete aneurysm occlusion without re-treatment, recurrent subarachnoid hemorrhage, or significant parent artery stenosis (defined as $>$ 50% stenosis). The one year primary safety endpoint is the proportion of patients with primary safety events (i.e., death from any non-accidental cause or any major stroke, defined as an ischemic or

hemorrhagic stroke resulting in an increase of ≥ 4 points on the National Institutes of Health Stroke Scale and persisting for seven days after the procedure) within the first 30 days after treatment, or a major ipsilateral stroke or neurologic death from day 31 to one year subsequent to treatment. Required characteristics of the aneurysm noted by the authors included the following: ruptured or unruptured, saccular in shape, located at the basilar apex (BA), middle cerebral artery (MCA) bifurcation, internal carotid artery terminus, or anterior communicating artery complex, dome-to-neck ratio ≥ 1 , wide-neck intracranial aneurysm with neck size ≥ 4 mm or dome-to-neck ratio < 2 , and diameter appropriate for treatment with the WEB device according to the device instructions for use. Exclusion criteria included vascular tortuosity or morphology that could preclude safe access and support during treatment with WEB and a modified Rankin Scale score of ≥ 2 at baseline or before rupture. The authors reported one primary safety event occurred; the event occurred on postoperative day 22 and was a delayed ipsilateral parenchymal hemorrhage. Angiograph results at 12 month follow up demonstrated 53.8% of subjects (77/143) had complete occlusion of the aneurysm and adequate occlusion of the aneurysm occurred in 84.6% of subjects (121/143). During the 12 months following treatment, 5.6% of subjects underwent or had planned target aneurysm retreatment (one with coils, four with stent assisted coiling, and three with flow diversion). When coupled with an additional six subjects considered failures within the first six months, the retreatment rate was 9.8% (14/143 subjects). The authors concluded that the WEB device provides an option for patients with wide-neck bifurcation aneurysms that when compared to other reported outcomes, is safer and more effective than currently available therapies. While comparative trials are limited, according to clinicaltrials.gov, a randomized trial evaluating the intrasaccular endobridge device (RISE Trial) for treatment of intracranial aneurysm is currently recruiting subjects. The study model description is the use of a standard best conventional treatment option (surgical or endovascular) versus the WEB embolization device (Clinicaltrials.gov, NCT03936647). The estimated completion of the study is July 2024.

There is a growing body of evidence in the peer-reviewed scientific literature evaluating WEB treatment for bifurcating intrasaccular aneurysms as an alternative to coiling. A majority of evidence is in the form of prospective case series (Goertz, et al., 2020; Pierot, et al., 2020; Pierot, et al., 2018; Fiorella, et al., 2017; van Rooji, et al., 2017; Papagiannaki, et al., 2014), retrospective case series (Kabbasch, et al., 2019; Sauvigny, et al., 2019; Maurer, et al., 2019; Lawson, et al., 2018; Mine, et al., 2018; Popielski, et al., 2018; Liebig, et al., 2015) and systematic reviews (Zhang, et al., 2020; Lv, et al., 2018; Tau, et al., 2018; Armoiry, et al., 2016). Outcomes are short-term and there is lack of high-quality evidence evaluating the long-term effectiveness and safety of the Woven Endobridge (WEB) device for use in the endovascular treatment of an intracranial bifurcating aneurysm. Although preliminary studies lend support to satisfactory results in the short term, randomized controlled trials and long-term clinical outcomes (e.g., 3-5 years post procedure) are lacking and further study is needed.

A number of systematic reviews have been published, many of which involve an overlapping of subjects/trials. In 2020, Zhang and colleagues conducted a systematic review and meta-analysis evaluating the effectiveness, safety and risk factors of the Woven EndoBridge (WEB) device in the treatment of wide-neck intracranial aneurysms. A total of 36 studies involving 1,759 patients with 1,749 aneurysms were included with a mean follow-up of 9.34 months. Eligible studies included those with five or more patients undergoing WEB for wide-neck intracranial aneurysms, reported angiographic or clinical outcomes and risk factors, and were published after December 1, 2012. The authors concluded that WEB has a satisfactory safety profile and shows promising efficacy in treating wide-neck intracranial aneurysms, adequate occlusion rates were nearly 80%, the most common complication was thromboembolism (9%), and morbidity and mortality were 6% and 7%, respectively. However, the risk factors for the long-term angiographic results and complications after WEB treatment in wide-neck aneurysms should be further studied by well-designed clinical trials (Zhang, et al., 2020).

Tau and associates (2018) published a systematic review with meta-analysis for the Woven EndoBridge (WEB) for endovascular therapy of intracranial aneurysms. In total (initial review and update), 12 uncontrolled case-series were included reporting outcomes for 940 patients, studies were published between January 2010 through September 2015 (initial report), and October 2015 through December 2017 (update). The median clinical follow-up was seven months (range 3-27.9 months), adequate occlusion rate was 81%, 9% of subjects required additional treatment, 14% had periprocedural complications, and mortality was 5% at seven months follow-up. The authors concluded that although WEB showed high rates of adequate aneurysm occlusion at mid-term, procedure-related complications and mortality rates were not negligible, and that additional studies are needed to compare the WEB device with other established treatment options.

Lv et al. (2018) conducted a systematic review of the published literature to evaluate the complications, complete occlusion rate, and morbidity and mortality of embolization results for WNBAs using the WEB device. A total of 19 studies, including 935 patients, were identified. The technical success rate was 97%, adequate occlusion rate was 81%, morbidity during follow-up was 3%, and mortality was 2%. Mid-term occlusion rates were slightly lower when compared to single-stent coiling.

At present, investigators continue to evaluate the safety and efficacy of intrasaccular flow diversion as a treatment alternative to coiling for treatment of an intracranial bifurcating aneurysm. Although preliminary studies lend support to satisfactory results in the short term, patient selection criteria is not firmly established and comparative, controlled trials with long-term clinical outcomes are lacking in the peer reviewed literature base; further trials are necessary to safety and efficacy.

Professional Society Recommendations

The American Association of Neurological Surgeons (AANS): The AANS provides general information regarding treatment of cerebral aneurysm using stent-assisted coiling, flow diversion stents and intrasaccular flow disruption. The AANS suggests stent-assisted coiling is performed to prevent herniation of the coiled mass into the parent artery which can lead to stroke. The stent is permanently placed across the aneurysm neck; anti-platelet medication is required. Flow diversion is generally performed for the treatment of aneurysms that are challenging and less amenable to traditional endovascular coiling, such as complex aneurysms, including large and giant aneurysms, wide-neck aneurysms, fusiform aneurysms and recanalized aneurysms after previous coiling. Stent-assisted coiling and balloon-assisted coiling are alternative endovascular options for such aneurysms; however, some studies reported their limited efficacy due to high recanalization rates and flow disruption. For treatment using intrasaccular flow disruption, a device such as the WEB device is used for management of wide-necked aneurysms at the bifurcation of an artery. The WEB device is placed within an aneurysm in contrast to the flow diversion devices which are placed in the parent artery. One of the biggest advantages is the reduced need for antiplatelet medications, especially for the ruptured aneurysms, which is particularly helpful in patients with subarachnoid hemorrhage; however, the clinical experience with its use is currently limited.

American Heart Association: The American Heart Association published a scientific statement regarding indications for the performance of intracranial endovascular neurointerventional procedures (Eskey, et al., 2018). Within this document, the AHA notes the document is not a clinical practice guideline. The treatments are changing rapidly, and as such, recommendations and levels of evidence are not included. The AHA makes the following suggestions regarding treatment of individuals with stroke or transient ischemic attack (TIA) resulting from stenosis of a major intracranial artery:

- For 50-69% stenosis, treatment with medical therapy
- For 70-99% stenosis, optimal medical therapy (aspirin, clopidogrel, systolic <140, statin therapy) and risk factor modification
- For 70-99% stenosis, intracranial stenting with the Wingspan or Pharos stent system should not be initial therapy, even in individuals on antithrombotic therapy at the time of stroke or TIA
- For severe stenosis (70-99%) and progressing symptoms, recurrent TIA or stroke, despite dual antiplatelet therapy, systolic <140, and high intensity statin therapy, angioplasty with or without Wingspan stent may be warranted
- The utility of angioplasty alone or placement of other than Wingspan or Pharos stent is unknown and investigational.

The American Heart Association / American Stroke Association published guidelines (Thompson, et al., 2015) for management of patients with unruptured intracranial aneurysms. The guideline is endorsed by the American Association of Neurological Surgeons, the Congress of Neurological Surgeons, and the Society of Noninterventional Surgery. Within these guidelines, the authors report that endoluminal flow diversion represents a new treatment and should be considered in carefully select cases for treatment of unruptured intracranial aneurysms (Class IIb, Level of Evidence B). Strict adherence to U.S. FDA indications is recommended. Class IIb Level B recommendation is defined as “usefulness / efficacy is less well established, greater conflicting evidence from a single randomized trial or nonrandomized studies”.

European Society for Vascular Surgery: Guidelines for the management of atherosclerotic carotid and vertebral artery disease (Naylor, et al., 2018) provide evidence based recommendations for treatment of extracranial atherosclerosis. Recommendations are defined as follows:

Class I: evidence and/or general agreement a given treatment or procedure is beneficial, useful, and effective

Class II: conflicting evidence and/or divergence of opinion about the usefulness/efficacy of the given treatment or procedure

Class IIa: weight of evidence /opinion is in favor of usefulness/efficacy

Class IIb: usefulness/efficacy is less well established by evidence/opinion

Class III: evidence or general agreement that the given treatment or procedure is not useful/effective, and in some cases may be harmful

Levels of evidence are defined as follows:

Level A: data derived from multiple randomized clinical trials or meta-analyses

Level B: data derived from a single randomized clinical trial or large non-randomised studies

Level C: consensus of opinion of the experts and/or small studies, retrospective studies, registries

Regarding management of vertebral artery stenosis the guideline states:

- Asymptomatic vertebral artery atherosclerosis should not be treated by open or endovascular interventions (Class III, Level C)
- Patients with recurrent vertebrobasilar territory symptoms despite best medical therapy, and who have a 50-99% extracranial vertebral artery stenosis may be considered for vascularization (Class IIb, Level B)

Regarding carotid artery stenosis the guideline states:

- It is recommended most patients who have suffered carotid territory symptoms within the preceding 6 months and who are aged >70 years and who have 50-99% stenosis should be treated by carotid endarterectomy rather than stenting (Class I, Level A)
- When revascularization is indicated in patients who have suffered carotid territory symptoms within the preceding 6 months and who are aged <70 years, carotid stenting may be considered an alternative to endarterectomy provided the documented procedural death/stroke rate is <6% (Class IIb, Level A)
- Carotid endarterectomy or carotid artery stenting are not recommended in symptomatic patients with a chronic internal carotid near-occlusion, unless associated with recurrent ipsilateral symptoms (despite optimal medical therapy) and following multidisciplinary team review (Class III, Level C)
- Carotid artery endarterectomy or stenting may be considered in recently symptomatic patients with <50% stenosis if they suffer recurrent symptoms despite best medical therapy and following multidisciplinary team review (Class IIb, Level C)
- In recently symptomatic patients with 50-99% stenosis and anatomical and/or medical comorbidities that are considered by the multidisciplinary team to make them "higher risk for carotid endarterectomy", carotid stenting should be considered as an alternative to endarterectomy provided the documented procedural death/stroke rate is < 6% (Class IIa, Level B).
- In average or high surgical risk patients with asymptomatic stenosis 60-99% and in the presence of one or more imaging characteristics that may be associated with an increased risk of ipsilateral stroke, CAS may be an alternative when the perioperative stroke/death rates are < 3% and the patient's life expectancy is greater than 5 years. Imaging characteristics of high stroke risk may include stenosis progression, silent infarction on CT, large plaque area, plaque echolucency, intra-plaque hemorrhage on magnetic resonance imaging, impaired cerebral vascular reserve, spontaneous embolization on transcranial Doppler, or history of contralateral TIA) (Class IIB, Level B)
- Prophylactic carotid artery endarterectomy and carotid stenting are not recommended in patients with asymptomatic carotid stenosis prior to major non-cardiac, non-vascular surgical procedures (Class III, Level B)

American Stroke Association (ASA)/American College of Cardiology Foundation (ACCF)/American Association of Neuroscience Nurses (AANN)/American Association of Neurological Surgeons (AANS)/American College of Radiology (ACR), American Society of Neuroradiology (ASNR), Congress of Neurological Surgeons (CNS), Society of Atherosclerosis Imaging and Prevention (SAIP), Society for Cardiovascular Angiography and Interventions (SCAI), Society of Interventional Radiology (SIR), Society

of NeuroInterventional Surgery (SNIS), Society for Vascular Medicine (SVM), and Society for Vascular Surgery (SVS) (2011): Brott et al. (2013) published joint consensus guidelines regarding the management of extracranial carotid and vertebral artery disease. Regarding the selection of patients for carotid revascularization, the authors made the following statements for carotid angioplasty and stenting (CAS):

- CAS is indicated as an alternative to CEA for symptomatic patients at average or low risk of complications associated with endovascular intervention when the diameter of the lumen of the internal carotid artery is reduced by > 70% (documented by noninvasive imaging) or > 50% (documented by catheter angiography) and the anticipated rate of periprocedural stroke or mortality is < 6% (level of evidence B [defined as data from a single randomized trial or non-randomized studies]). (Class I, level of evidence B recommendation [procedure is useful/effective, evidence from single RCT or nonrandomized studies])
- CAS is indicated for asymptomatic patients with 70-99% stenosis (Class IIb, level of evidence B recommendation [recommendation in favor of procedure being useful/effective, some conflicting evidence from single RCT or nonrandomized trials])

Regarding extracranial vertebral artery stenosis the authors concluded there is insufficient evidence to demonstrate that endovascular management is superior to medical therapy.

American College of Cardiology: In 2007, the American College of Cardiology (ACC) issued a joint expert consensus document on CAS (American College of Cardiology Foundation, et al., 2007). The document states, "Carotid artery stenting is a reasonable alternative to CEA, particularly in patients at high risk for CEA. Although there are no randomized studies comparing CAS with and without embolic protection devices, the use of embolic protection devices appears to be important in reducing the risk of stroke during CAS. Careful neurological assessment is required before and after CAS. At the present time, there is insufficient evidence to support CAS in high-risk patients with asymptomatic stenosis less than 80% or in any patient without high-risk features. Operators should previously have achieved a high level of proficiency in catheter-based intervention, complete dedicated training in CAS, and be credentialed at their hospital."

The document outlines the high-risk criteria for CEA into anatomical criteria and medical comorbidities. Anatomical criteria includes: lesion at C-2 or higher, lesion below clavicle, prior radical neck surgery or radiation, prior ipsilateral CEA, contralateral laryngeal nerve palsy, and tracheostoma. Medical comorbidities include: age \geq 80 years, Class III/IV congestive heart failure, Class III/IV angina pectoris, left main/ \geq two vessel coronary disease, urgent (< 30 days) heart surgery, left ventricle ejection fraction \leq 30%, recent (< 30 days) MI, severe chronic lung disease, and severe renal disease. There has been no update to this consensus document since 2007.

The Society of Cardiac Angiography and Interventions (SCAI), the Society for Vascular Medicine and Biology (SVMB), and the Society for Vascular Surgery (SVS): Carotid stenting is a technically complex procedure. Minor embolic events can lead to major complications. The Society of Cardiac Angiography and Interventions (SCAI), the Society for Vascular Medicine and Biology (SVMB), and the Society for Vascular Surgery (SVS) issued a clinical competence statement on carotid stenting addressing the training and credentialing for carotid stenting. This multispecialty consensus recommendation states that physicians who perform carotid stenting with embolic protection must meet or exceed minimum qualifications deemed necessary to offer safe and effective therapy. The qualifications must include proficiency in the cognitive, technical, and clinical skills necessary to care for patients with carotid artery disease (Rosenfield, et al., 2005).

The American Board of Internal Medicine's (ABIM) Foundation Choosing Wisely® Initiative American Academy of Neurology: The AAN Choosing Wisely recommendation (2013) does not recommend carotid endarterectomy for asymptomatic carotid stenosis unless the complication rate is low (< 3%). According to the AAN statement "Based on studies reporting an upfront surgical complication rate ranging from 2.3% (ACAS) to 3.1% (ACST) among patients undergoing carotid endarterectomy (CEA) for asymptomatic stenosis of >60%, and an absolute risk reduction for stroke or death of roughly 5-6% in the surgical group at 5 years, several specialty societies (Brott, et al., 2013; Goldstein et al., 2011; Ricotta et al., 2011; Chaturvedi et al., 2005) have recommended that surgery for asymptomatic patients should be reserved for those with a perioperative complication risk of < 3% and a life expectancy of greater than 3-5 years. The cited 3% threshold for complication rates may be high because more recent studies have reported lower stroke rates with

improvements in both surgical (Brott, et al., 2010) and medical (Marquardt, et al., 2010) management. However, there are no recent randomized trials comparing these treatments. Given this, the more recent AHA guidelines (Brott, et al., 2013) state that it is “reasonable” to perform CEA for asymptomatic patients with >70% stenosis if the surgical complication rate is “low.” Reported complication rates vary widely by location (Kresowik, et al., 2004), and are dependent on how complications are tracked (self-report vs. neurologist’s evaluation vs. administrative data (Wolff, et al., 2007). Despite calls for rigorous monitoring 15 years ago (Goldstein, et al., 2011), most patients will likely need to rely on the surgeon’s self-reported rates.”

Use Outside of the US

European Society for Vascular Surgery: Guidelines for the management of atherosclerotic carotid and vertebral artery disease (Naylor, et al., 2018) provide evidence based recommendations for treatment of extracranial atherosclerosis. Recommendations are described above.

National Institute for Health and Clinical Excellence (NICE) (United Kingdom): NICE Interventional Procedure Guidance for carotid artery stent placement for asymptomatic extracranial carotid stenosis shows well documented risks, in particular for risk of stroke (NICE, 2011a). NICE Interventional Procedure Guidance for carotid artery stent placement for symptomatic extracranial carotid stenosis states, “Current evidence on the safety and efficacy of carotid artery stent placement for symptomatic extracranial carotid stenosis is adequate to support the use of this procedure” (NICE, 2011b).

NICE Medical technologies guidance for Pipeline Flex embolisation device with Shield Technology for the treatment of complex intracranial aneurysms states, “The case for adopting the Pipeline Flex embolisation device with Shield Technology in the NHS is supported by the current evidence when it is used in patients with complex giant or large intracranial aneurysms which are unsuitable for surgery and being considered for stenting, and where large numbers of coils would be needed during stent-assisted coiling” (NICE, 2012).

In August 2019, NICE published interventional procedure guidance for “Endovascular insertion of an intrasaccular wire-mesh blood-flow disruption device for intracranial aneurysms” and concluded that the current evidence on the safety and efficacy of endovascular insertion of an intrasaccular wire-mesh blood-flow disruption device for intracranial aneurysms is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit. Patient selection should be done by a multidisciplinary team, except for emergency situations, when this may be replaced by a discussion between an interventional neuroradiologist and neurosurgeon. Evidence reviewed by the authors included one systematic review, two non-randomized comparative studies, nine case series and one case report (NICE, 2019).

Choosing Wisely Canada

Vascular Surgery recommendations from Choosing Wisely Canada state: Don’t perform carotid endarterectomies or stenting in most asymptomatic high risk patients with limited life expectancy. The purpose of carotid artery surgery and stenting is to prevent stroke and, when combined with appropriate medical therapy, is a successful strategy in selected, mainly symptomatic, patients. Medical therapy alone is an effective alternative in many asymptomatic patients and safer in those who are elderly or at high risk for surgery and stenting and don’t have the life expectancy to benefit from such a prophylactic procedure.

Medicare Coverage Determinations

| | Contractor | Policy Name/Number | Revision Effective Date |
|-----|------------|--|-------------------------|
| NCD | National | Vertebral Artery Surgery (20.1) | Longstanding |
| NCD | National | Percutaneous Transluminal Angioplasty (PTA) (20.7) | 3/11/2013 |
| LCD | | No Local Coverage Determination found | |

Note: Please review the current Medicare Policy for the most up-to-date information.

Coding/Billing Information

Note: 1) This list of codes may not be all-inclusive.

2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Extracranial Angioplasty and Stenting

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

| CPT®* Codes | Description |
|--------------------|--|
| 37215 | Transcatheter placement of intravascular stent(s), cervical carotid artery, open or percutaneous, including angioplasty, when performed, and radiological supervision and interpretation; with distal embolic protection |
| 37216 | Transcatheter placement of intravascular stent(s), cervical carotid artery, open or percutaneous, including angioplasty, when performed, and radiological supervision and interpretation; without distal embolic protection |
| 37217 | Transcatheter placement of intravascular stent(s), intrathoracic common carotid artery or innominate artery by retrograde treatment, open ipsilateral cervical carotid artery exposure, including angioplasty, when performed, and radiological supervision and interpretation |
| 37218 | Transcatheter placement of intravascular stent(s), intrathoracic common carotid artery or innominate artery, open or percutaneous antegrade approach, including angioplasty, when performed, and radiological supervision and interpretation |
| 0075T | Transcatheter placement of extracranial vertebral artery stent(s), including radiologic supervision and interpretation, open or percutaneous; initial vessel |
| 0076T | Transcatheter placement of extracranial vertebral artery stent(s), including radiologic supervision and interpretation, open or percutaneous; each additional vessel (List separately in addition to code for primary procedure) |

Intracranial Aneurysm Repair

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

| CPT®* Codes | Description |
|--------------------|---|
| 61700 | Surgery of simple intracranial aneurysm, intracranial approach; carotid circulation |
| 61702 | Surgery of simple intracranial aneurysm, intracranial approach; vertebrobasilar circulation |
| 61703 | Surgery of intracranial aneurysm, cervical approach by application of occluding clamp to cervical carotid artery (Selverstone-Crutchfield type) |
| 61705 | Surgery of aneurysm, vascular malformation or carotid-cavernous fistula; by intracranial and cervical occlusion of carotid artery |
| 61708 | Surgery of aneurysm, vascular malformation or carotid-cavernous fistula; by intracranial electrothrombosis |
| 61710 | Surgery of aneurysm, vascular malformation or carotid-cavernous fistula; by intra-arterial embolization, injection procedure, or balloon catheter |

Intracranial Angioplasty and Stenting

Experimental, Investigational or Unproven when used to report treatment of atherosclerosis of intracranial arteries:

| CPT®* Codes | Description |
|--------------------|---|
| 61630 | Balloon angioplasty, intracranial (eg, atherosclerotic stenosis), percutaneous |
| 61635 | Transcatheter placement of intravascular stent(s), intracranial (eg, atherosclerotic stenosis), including balloon angioplasty, if performed |

Intracranial Stent Placement

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

| CPT®* Codes | Description |
|-------------|--|
| 61624 | Transcatheter permanent occlusion or embolization (eg, for tumor destruction, to achieve hemostasis, to occlude a vascular malformation), percutaneous, any method; central nervous system (intracranial, spinal cord) |

Experimental, Investigational, Unproven when used to report intrasaccular flow diversion:

| CPT®* Codes | Description |
|-------------|--|
| 61624 | Transcatheter permanent occlusion or embolization (eg, for tumor destruction, to achieve hemostasis, to occlude a vascular malformation), percutaneous, any method; central nervous system (intracranial, spinal cord) |

*Current Procedural Terminology (CPT®) ©2020 American Medical Association: Chicago, IL.

References

1. American College of Cardiology Foundation; American Society of Interventional & Therapeutic Neuroradiology; Society for Cardiovascular Angiography and Interventions; Society for Vascular Medicine and Biology; Society of Interventional Radiology, Bates ER, Babb JD, Casey DE Jr, Cates CU, Duckwiler GR, Feldman TE, Gray WA, Ouriel K, Peterson ED, Rosenfield K, Rundback JH, Safian RD, Sloan MA, White CJ. ACCF/SCAI/SVMB/SIR/ASITN 2007 clinical expert consensus document on carotid stenting: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents (ACCF/SCAI/SVMB/SIR/ASITN Clinical Expert Consensus Document Committee on Carotid Stenting). *J Am Coll Cardiol*. 2007 Jan 2;49(1):126-70.
2. Armoiry X, Turjman F, Hartmann DJ, et al. Endovascular Treatment of Intracranial Aneurysms with the WEB Device: A Systematic Review of Clinical Outcomes. *AJNR Am J Neuroradiol*. 2016;37(5):868-872.
3. Arthur AS, Molyneux A, Coon AL, et al. The safety and effectiveness of the Woven EndoBridge (WEB) system for the treatment of wide-necked bifurcation aneurysms: final 12-month results of the pivotal WEB Intrasaccular Therapy (WEB-IT) Study. *J Neurointerv Surg*. 2019;11(9):924-930.
4. Barr JD, Connors JJ 3rd, Sacks D, Wojak JC, Becker GJ, Cardella JF, Chopko B, Dion JE, Fox AJ, Higashida RT, Hurst RW, Lewis CA, Matalon TA, Nesbit GM, Pollock JA, Russell EJ, Seidenwurm DJ, Wallace RC; SIR Standards of Practice Committees. Quality improvement guidelines for the performance of cervical carotid angioplasty and stent placement. *AJNR Am J Neuroradiol*. 2003 Nov-Dec;24(10):2020-34.
5. Becske T, Brinjikji W, Potts MB, et al. Long-Term Clinical and Angiographic Outcomes Following Pipeline Embolization Device Treatment of Complex Internal Carotid Artery Aneurysms: Five-Year Results of the Pipeline for Uncoilable or Failed Aneurysms Trial. *Neurosurgery*. 2017 Jan 1;80(1):40-48.
6. Blackshear JL, Cutlip DE, Roubin GS, Hill MD, Leimgruber PP, Begg RJ, et al; CREST Investigators. Myocardial infarction after carotid stenting and endarterectomy: results from the carotid revascularization endarterectomy versus stenting trial. *Circulation*. 2011 Jun 7;123(22):2571-8. Epub 2011 May 23.
7. Bonati LH, Ederle J, McCabe DJ, Dobson J, Featherstone RL, Gaines PA, et al; CAVATAS Investigators. Long-term risk of carotid restenosis in patients randomly assigned to endovascular

treatment or endarterectomy in the Carotid and Vertebral Artery Transluminal Angioplasty Study (CAVATAS): long-term follow-up of a randomised trial. *Lancet Neurol.* 2009 Oct;8(10):908-17. Epub 2009 Aug 28.

8. Bonati LH, Jongen LM, Haller S, Flach HZ, Dobson J, Nederkoorn PJ, et al; ICSS-MRI study group. New ischaemic brain lesions on MRI after stenting or endarterectomy for symptomatic carotid stenosis: a substudy of the International Carotid Stenting Study (ICSS). *Lancet Neurol.* 2010 Apr;9(4):353-62. Epub 2010 Feb 25.
9. Brothers TE, Ricotta JJ 2nd, Gillespie DL, et al. Contemporary results of carotid endarterectomy in "normal-risk" patients from the Society for Vascular Surgery Vascular Registry. *J Vasc Surg.* 2015 Oct;62(4):923-8.
10. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. Developed in collaboration with the American Academy of Neurology and Society of Cardiovascular Computed Tomography. *Catheter Cardiovasc Interv.* 2013 Jan 1;81(1):E76-123.
11. Brott TG, Hobson RW 2nd, Howard G, Roubin GS, Clark WM, Brooks W, et al; CREST Investigators. Stenting versus endarterectomy for treatment of carotid-artery stenosis. *N Engl J Med.* 2010 Jul 1;363(1):11-23. Epub 2010 May 26. Erratum in: *N Engl J Med.* 2010 Jul 8;363(2):198. *N Engl J Med.* 2010 Jul 29;363(5):498.
12. Brown MM, Rogers J, Bland JM. Endovascular versus surgical treatment in patients with carotid stenosis in the Carotid and Vertebral Artery Transluminal Angioplasty Study (CAVATAS): a randomized trial. *Lancet.* 2001;357(9270):1729-37.
13. Burton KR, Lindsay TF. Assessment of short-term outcomes for protected carotid angioplasty with stents using recent evidence. *J Vasc Surg.* 2005 Dec;42(6):1094-100.
14. Caroff J, Mihalea C, Klisch J, et al. Single-Layer WEBs: Intracascular Flow Disrupters for Aneurysm Treatment—Feasibility Results from a European Study. *AJNR Am J Neuroradiol.* 2015;36(10):1942-1946.
15. Centers for Medicare and Medicaid Services (CMS). Decision memo for Carotid Artery Stenting (CAG-0085R). National Coverage Analyses. Medical Coverage Database. Baltimore, MD: CMS: Mar 17, 2005. Accessed Jun 30, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=157&keyword=0085r&keywordType=any&areald=all&docType=NCA&contractOption=all&sortBy=relevance&KeywordLookup=Doc&KeywordSearchType=Or&bc=AAAAAAQACAAAAAA&>
16. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed Jun 30, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx>
17. Center for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed Jun 30, 2021. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/ncd-alphabetical-index.aspx>

18. Chaturvedi S, Bruno A, Feasby T, et al. Carotid endarterectomy--an evidence-based review: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2005 Sep 27;65(6):794-801.
19. Chimowitz MI, Lynn MJ, Derdeyn CP, et al., SAMPRIS Trial Investigators. Stenting versus aggressive medical therapy for intracranial arterial stenosis. *N Engl J Med*. 2011 Sep 15;365(11):993-1003.
20. Choosing Wisely. American Academy of Neurology. Feb 21, 2013. Accessed Jun 30, 2021. Available at URL address: <http://www.choosingwisely.org/societies/american-academy-of-neurology/>
21. Choosing Wisely Canada. Canadian Society for Vascular Surgery. Updated Aug 2020. Accessed Jun 30, 2021. Available at URL address: <https://choosingwiselycanada.org/vascular-surgery/>
22. Cremonesi A, Setacci C, Bignamini A, Bolognese L, Briganti F, Di Sciascio G, Inzitari D, Lanza G, Lupattelli L, Mangiafico S, Pratesi C, Reimers B, Ricci S, de Donato G, Ugolotti U, Zaninelli A, Gensini GF. Carotid artery stenting: first consensus document of the ICCS-SPREAD Joint Committee. *Stroke*. 2006 Sep;37(9):2400-9.
23. Eckstein HH. European Society for Vascular Surgery Guidelines on the Management of Atherosclerotic Carotid and Vertebral Artery Disease. *Eur J Vasc Endovasc Surg*. 2018 Jan;55(1):1-2.
24. Eckstein HH, Ringleb P, Allenberg JR, Berger J, Fraedrich G, Hacke W, et al. Results of the Stent-Protected Angioplasty versus Carotid Endarterectomy (SPACE) study to treat symptomatic stenoses at 2 years: a multinational, prospective, randomised trial. *Lancet Neurol*. 2008 Oct;7(10):893-902.
25. Ederle J, Bonati LH, Dobson J, Featherstone RL, Gaines PA, Beard JD, et al; CAVATAS Investigators. Endovascular treatment with angioplasty or stenting versus endarterectomy in patients with carotid artery stenosis in the Carotid and Vertebral Artery Transluminal Angioplasty Study (CAVATAS): long-term follow-up of a randomised trial. *Lancet Neurol*. 2009 Oct;8(10):898-907. Epub 2009 Aug 28.
26. Ederle J, Featherstone RL, Brown MM. Randomized controlled trials comparing endarterectomy and endovascular treatment for carotid artery stenosis: a Cochrane systematic review. *Stroke*. 2009 Apr;40(4):1373-80. Epub 2009 Feb 19. Review.
27. Eisenhauer AC, White CJ. Endovascular Treatment of Noncoronary Obstructive Vascular Disease. In: Zipes DP, Libby P, Bonow RO, Braunwald E, editors. *Braunwald's Heart Disease. A Textbook of Cardiovascular Medicine*. 8th ed. St. Louis, MO: W.B. Saunders Company; 2008. Ch.59.
28. Eskey CJ, Meyers PM, Nguyen TN, et al. Indications for the Performance of Intracranial Endovascular Neurointerventional Procedures: A Scientific Statement From the American Heart Association. *Circulation*. 2018;137(21):e661-e689.
29. Fiorella D, Molyneux A, Coon A, et al. Demographic, procedural and 30-day safety results from the WEB Intra-saccular Therapy Study (WEB-IT). *Journal of Neurointerventional Surgery*. 2017; 9: 1191–6.
30. Goertz L, Liebig T, Siebert E, et al. Risk Factors of Procedural Complications Related to Woven EndoBridge (WEB) Embolization of Intracranial Aneurysms. *Clin Neuroradiol*. 2020;30(2):297-304.
31. Goldstein LB, Bushnell CD, Adams RJ, et al. Guidelines for the primary prevention of stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 2011 Feb; 42(2):517-84.
32. Gray WA, Yadav JS, Verta P, Scicli A, Fairman R, Wholey M, et al. The CAPTURE registry: results of carotid stenting with embolic protection in the post approval setting. *Catheter Cardiovasc Interv*. 2007 Feb 15;69(3):341-8.

33. Groschel K, Riecker A, Schulz JB, Ernemann U, Kastrup A. Systematic review of early recurrent stenosis after carotid angioplasty and stenting. *Stroke*. 2005 Feb;36(2):367-73.
34. Gurm HS, Nallamotheu BK, Yadav J. Safety of carotid artery stenting for symptomatic carotid artery disease: a meta-analysis. *Eur Heart J*. 2008a Jan;29(1):113-9.
35. Gurm HS, Yadav JS, Fayad P, et al., SAPPHIRE Investigators. Long-term results of carotid stenting versus endarterectomy in high-risk patients. *N Engl J Med*. 2008b Apr 10;358(15):1572-9.
36. Hayes, Inc. Evidence Analysis Research Brief. Woven EndoBridge Aneurysm Embolization System for Treatment of Intracranial Aneurysms. Hayes, Inc.; Aug 28, 2020.
37. Hayes, Inc. Health Technology Assessment. Flow Diverter Devices for Intracranial Aneurysms: A review of reviews. Hayes, Inc. March 30, 2017.
38. Hobson RW 2nd, Howard VJ, Roubin GS, Brott TG, Ferguson RD, Popma JJ, et al; CREST Investigators. Carotid artery stenting is associated with increased complications in octogenarians: 30-day stroke and death rates in the CREST lead-in phase. *J Vasc Surg*. 2004 Dec;40(6):1106-11.
39. Hussain MS, Fraser JF, Abruzzo T, et al. Standard of practice: endovascular treatment of intracranial atherosclerosis. *J Neurointerv Surg*. 2012;4(6):397-406.
40. Iyer SS, White CJ, Hopkins LN, Katzen BT, Safian R, Wholey MH, et al.; BEACH Investigators. Carotid artery revascularization in high-surgical-risk patients using the Carotid WALLSTENT and FilterWire EX/EZ: 1-year outcomes in the BEACH Pivotal Group. *J Am Coll Cardiol*. 2008 Jan 29;51(4):427-34.
41. Jauch EC, Saver JL, Adams HP Jr, Bruno A, Connors JJ, Demaerschalk BM, Khatri P, McMullan PW Jr, Qureshi AI, Rosenfield K, Scott PA, Summers DR, Wang DZ, Wintermark M, Yonas H; American Heart Association Stroke Council; Council on Cardiovascular Nursing; Council on Peripheral Vascular Disease; Council on Clinical Cardiology. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013 Mar;44(3):870-947.
42. Kabbasch C, Goertz L, Siebert E, et al. WEB embolization versus stent-assisted coiling: comparison of complication rates and angiographic outcomes. *J Neurointerv Surg*. 2019;11(8):812-816.
43. Kallmes DF, Brinjikji W, Cekirge S, et al. Safety and efficacy of the Pipeline embolization device for treatment of intracranial aneurysms: a pooled analysis of 3 large studies. *J Neurosurg*. 2017 Oct;127(4):775-780.
44. Kallmes DF, Hanel R, Lopes D, et al. International retrospective study of the pipeline embolization device: a multicenter aneurysm treatment study. *AJNR Am J Neuroradiol*. 2015 Jan;36(1):108-15.
45. Katzen BT, Criado FJ, Ramee SR, Massop DW, Hopkins LN, Donohoe D, et al; on behalf of the CASES-PMS Investigators. Carotid artery stenting with emboli protection surveillance study: thirty-day results of the CASES-PMS study. *Catheter Cardiovasc Interv*. 2007 Aug 1;70(2):316-23.
46. Kresowik TF, Bratzler DW, Kresowik RA, Hendel ME, Grund SL, Brown KR, Niladena DS. Multistate improvement in process and outcomes of carotid endarterectomy. *J Vasc Surg [Internet]*. 2004 [cited 2013 Jan 3];39:372-380.
47. Lawson A, Molyneux A, Sellar R, et al. Safety results from the treatment of 109 cerebral aneurysms using the Woven EndoBridge technique: preliminary results in the United Kingdom. *Journal of Neurosurgery*. 2018 Jan; 128: 144–53.

48. Li K, Barras CD, Chandra RV, et al. A review on the management of cerebral vasospasm following aneurysmal subarachnoid haemorrhage. *World Neurosurg*. 2019 Mar 18. pii: S1878-8750(19)30751-X.
49. Liebig T, Kabbasch C, Strasilla C, et al. Intracascular Flow Disruption in Acutely Ruptured Aneurysms: A Multicenter Retrospective Review of the Use of the WEB. *AJNR Am J Neuroradiol*. 2015;36(9):1721-1727.
50. Lv X, Zhang Y, Jiang W. Systematic Review of Woven EndoBridge for Wide-Necked Bifurcation Aneurysms: Complications, Adequate Occlusion Rate, Morbidity, and Mortality. *World Neurosurg*. 2018;110:20-25.
51. Mantese VA, Timaran CH, Chiu D, Begg RJ, Brott TG; CREST Investigators. The Carotid Revascularization Endarterectomy versus Stenting Trial (CREST): stenting versus carotid endarterectomy for carotid disease. *Stroke*. 2010 Oct;41(10 Suppl):S31-4.
52. Maragos GA, Dmytriw AA, Salem MM, et al. Overview of Different Flow Diverters and Flow Dynamics. *Neurosurgery*. 2020;86(Suppl 1):S21-S34.
53. Marquardt L, Geraghty OC, Mehta Z, Rothwell PM. Low risk of ipsilateral stroke in patients with asymptomatic carotid stenosis on best medical treatment: a prospective, population-based study. *Stroke* [Internet]. 2010 [cited 2013 Jan 1];41:e11-e7.
54. Maurer C, König I, Berlis A, Weber W, Fischer S. Two-Center Experience in the Endovascular Treatment of Intracranial Aneurysms Using the Woven EndoBridge 17 Device Including Midterm Follow-Up Results: A Retrospective Analysis. *AJNR Am J Neuroradiol*. 2019;40(9):1517-1522
55. Meier P, Knapp G, Tamhane U, Chaturvedi S, Gurm HS. Short term and intermediate term comparison of endarterectomy versus stenting for carotid artery stenosis: systematic review and meta-analysis of randomised controlled clinical trials [published correction appears in *BMJ*. 2010;340:c1798]. *BMJ*. 2010;340:c467. Published 2010 Feb 12.
56. Mine B, Goutte A, Brisbois D, et al. Endovascular treatment of intracranial aneurysms with the Woven EndoBridge device: mid term and long term results. *Journal of Neurointerventional Surgery*. 2018 Feb; 10: 127–32.
57. Murad MH, Shahrour A, Shah ND, Montori VM, Ricotta JJ. A systematic review and meta-analysis of randomized trials of carotid endarterectomy vs stenting. *J Vasc Surg*. 2011 Mar;53(3):792-7. Epub 2011 Jan 8.
58. National Institute for Health and Clinical Excellence (NICE). Carotid artery stent placement for asymptomatic extracranial carotid stenosis. IPG388. Apr 27, 2011a. Accessed Jun 30, 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg388>
59. National Institute for Health and Clinical Excellence (NICE). Carotid artery stent placement for symptomatic extracranial stenosis. IPG389. Apr 27, 2011b. Accessed Jun 30, 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg389>
60. National Institute for Health and Clinical Excellence (NICE). Endovascular insertion of an intracascular wire-mesh blood-flow disruption device for intracranial aneurysms. IPG658. Aug 14, 2019. Accessed Jun 30, 2021. Available at URL address: <https://www.nice.org.uk/guidance/ipg658>
61. National Institute for Health and Clinical Excellence (NICE). Pipeline Flex embolisation device with Shield Technology for the treatment of complex intracranial aneurysms. MTG10. May 30, 2012. Last updated Jan 9, 2019. Accessed Jun 30, 2021. Available at URL address: <https://www.nice.org.uk/guidance/mtg10>

62. Naylor AR, Ricco JB, deBorst GJ, et al., Editor's Choice: Management of Atherosclerotic Carotid and Vertebral Artery Disease: 2017 Clinical Practice guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg* (2018) 55, 3-81.
63. North American Symptomatic Carotid Endarterectomy Trial (NASCET) Collaborators. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade carotid stenosis. *N Engl J Med*. 1991;325:445-53.
64. Padalia A, Sambursky JA, Skinner C, Moureiden M. Percutaneous Transluminal Angioplasty with Stent Placement versus Best Medical Therapy Alone in Symptomatic Intracranial Arterial Stenosis: A Best Evidence Review. *Cureus*. 2018 Jul 16;10(7):e2988.
65. Papagiannaki C, Spelle L, Januel AC, et al. WEB intrasaccular flow disruptor-prospective, multicenter experience in 83 patients with 85 aneurysms. *AJNR Am J Neuroradiol*. 2014;35(11):2106-2111.
66. Pierot L, Liebig T, Sychra V, et al. Intrasaccular flow-disruption treatment of intracranial aneurysms: preliminary results of a multicenter clinical study. *AJNR Am J Neuroradiol*. 2012;33(7):1232-1238.
67. Pierot L, Moret J, Barreau X, et al. Aneurysm Treatment With Woven EndoBridge in the Cumulative Population of 3 Prospective, Multicenter Series: 2-Year Follow-Up. *Neurosurgery*. 2020;87(2):357-367.
68. Pierot L, Moret J, Barreau X, et al. Safety and efficacy of aneurysm treatment with WEB in the cumulative population of three prospective, multicenter series. *Journal of Neurointerventional Surgery*, 2018 Jun; 10: 553–9.
69. Popielski J, Berlis A, Weber W, Fischer S. Two-Center Experience in the Endovascular Treatment of Ruptured and Unruptured Intracranial Aneurysms Using the WEB Device: A Retrospective Analysis. *AJNR Am J Neuroradiol*. 2018;39(1):111-117.
70. Ricotta JJ, Aburahma A, Ascher E, et al; Society for Vascular Surgery. Updated Society for Vascular Surgery guidelines for management of extracranial carotid disease: executive summary. *J Vasc Surg*. 2011 Sep;54(3):832-6.
71. Rosenfield K, Babb JD, Cates CU, et al. Clinical competence statement on carotid stenting: training and credentialing for carotid stenting--multispecialty consensus recommendations: a report of the SCAI/SVMB/SVS Writing Committee to develop a clinical competence statement on carotid interventions. *J Am Coll Cardiol*. 2005;45(1):165-174.
72. Sauvigny T, Nawka MT, Schweingruber N, et al. Early clinical course after aneurysmal subarachnoid hemorrhage: comparison of patients treated with Woven EndoBridge, microsurgical clipping, or endovascular coiling. *Acta Neurochir (Wien)*. 2019;161(9):1763-1773.
73. Tau N, Sadeh-Gonik U, Aulagner G, et al. The Woven EndoBridge (WEB) for endovascular therapy of intracranial aneurysms: Update of a systematic review with meta-analysis. *Clinical Neurology & Neurosurger*, 2018 Mar; 166: 110–5.
74. Thompson BG, Brown RD Jr, Amin-Hanjani S, et al. Guidelines for the Management of Patients With Unruptured Intracranial Aneurysms: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2015 Aug;46(8):2368-400.
75. U.S. Food and Drug Administration Center for Devices and Radiological Health (CDRH). CDRH Consumer Information. New Device Approval: ACCULINK Carotid Stent System RX ACCULINK Carotid Stent System-P040012. Approval Aug 30, 2004. Accessed Aug 22, 2011. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pma&id=320201>

76. U.S. Food & Drug Administration (FDA). Premarket Approval Database. Surpass Streamline Flow Diverter. Approval Jul 13, 2018. PMA: P170024. Accessed Jun 30, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pma&id=410001>
77. U.S. Food & Drug Administration (FDA). Premarket Approval Database. Pipeline® Embolization Device, Approval Apr 06, 2011. PMA: P100018. Accessed Jun 30, 2021. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pma&id=320441>
78. U.S. Food & Drug Administration (FDA). Summary of Safety and Probably Benefit. NEUROLINK® System, Sep 5, 2001. Humanitarian Device Exemption H010004. Accessed Jun 30, 2021. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf/H010004b.pdf
79. U.S. Food & Drug Administration (FDA). Summary of Safety and Probably Benefit. CORDIS ENTERPRISE™ Vascular Reconstruction Device and Delivery System. May 26, 2005. Humanitarian Device Exemption H060001. Accessed Jun 30, 2021. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf6/H060001B.pdf
80. UpToDate® Management of asymptomatic extracranial carotid atherosclerotic disease. Last updated Jun 29, 2021. Accessed Jun 30, 2021. Available at URL address: https://www.uptodate.com/contents/management-of-asymptomatic-extracranial-carotid-atherosclerotic-disease?search=Management%20of%20asymptomatic%20carotid%20atherosclerotic%20disease&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
81. UpToDate® Management of symptomatic carotid atherosclerotic disease. Last updated Aug 11, 2020. Accessed Jun 30, 2021. Available at URL address: https://www.uptodate.com/contents/management-of-symptomatic-carotid-atherosclerotic-disease?search=Management%20of%20symptomatic%20carotid%20atherosclerotic%20disease&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
82. UpToDate® Overview of carotid artery stenting. Last updated Jun 29, 2021. Accessed Jun 30, 2021. Available at URL address: https://www.uptodate.com/contents/overview-of-carotid-artery-stenting?search=Carotid%20artery%20stenting%20and%20its%20complications&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
83. Usman AA, Tang GL, Eskandari MK. Metaanalysis of procedural stroke and death among octogenarians: carotid stenting versus carotid endarterectomy. *J Am Coll Surg.* 2009 Jun;208(6):1124-31.
84. van Rooij S, Sprengers ME, Peluso JP, et al. A systematic review and meta-analysis of Woven EndoBridge single layer for treatment of intracranial aneurysms. *Interv Neuroradiol.* 2020;26(4):455-460.
85. van Rooij SBT, van Rooij WJ, Peluso JP, et al. WEB Treatment of Ruptured Intracranial Aneurysms: A Single-Center Cohort of 100 Patients. *AJNR Am J Neuroradiol.* 2017;38(12):2282-2287.
86. Wang T, Luo J, Wang X, Yang K, Jadhav V, Gao P, Ma Y, Zhao N, Jiao L. Endovascular therapy versus medical treatment for symptomatic intracranial artery stenosis. *Cochrane Database of Systematic Reviews* 2020, Issue 8. Art. No.: CD013267.
87. White CJ, Beckman JA, Cambria RP, Comerota AJ, Gray WA, Hobson RW 2nd, Iyer SS; American Heart Association Writing Group 5. Atherosclerotic Peripheral Vascular Disease Symposium II: controversies in carotid artery revascularization. *Circulation.* 2008 Dec 16;118(25):2852-9.
88. White CJ, Iyer SS, Hopkins LN, Katzen BT, Russell ME; BEACH Trial Investigators. Carotid stenting with distal protection in high surgical risk patients: the BEACH trial 30 day results. *Catheter Cardiovasc Interv.* 2006 Apr;67(4):503-12.

89. Wolff T, Guirguis-Blake J, Miller T, Gillespie M, Harris R. Screening For Asymptomatic Carotid Artery Stenosis. Rockville (MD): Agency for Healthcare Research and Quality (US); December 2007.
90. Yadav JS. Carotid stenting in high-risk patients: design and rationale of the SAPPHIRE trial. *Cleve Clin J Med.* 2004;71 Suppl 1:S45-S46.
91. Yavin D, Roberts DJ, Tso M, Sutherland GR, Eliasziw M, Wong JH. Carotid endarterectomy versus stenting: a meta-analysis of randomized trials. *Can J Neurol Sci.* 2011 Mar;38(2):230-5.
92. Zahn R, Hochadel M, Grau A, Senges J. Stent-supported angioplasty versus endarterectomy for carotid artery stenosis: evidence from current randomized trials. *Z Kardiol.* 2005 Dec;94(12):836-43.
93. Zaidat OO, Fitzsimmons BF, Woodward BK, et al, VISSIT Trial Investigators. Effect of a balloon-expandable intracranial stent vs medical therapy on risk of stroke in patients with symptomatic intracranial stenosis: the VISSIT randomized clinical trial. *JAMA.* 2015 Mar 24-31;313(12):1240-8.
94. Zhang SM, Liu LX, Ren PW, Xie XD, Miao J. Effectiveness, Safety and Risk Factors of Woven EndoBridge Device in the Treatment of Wide-Neck Intracranial Aneurysms: Systematic Review and Meta-Analysis. *World Neurosurg.* 2020;136:e1-e23.
95. Zhou Y, Wang L, Zhang JR, Zhou K, et al. Angioplasty and stenting for severe symptomatic atherosclerotic stenosis of intracranial vertebrobasilar artery. *J Clin Neurosci.* 2019 Mar 5. pii: S0967-5868(18)32019-8.

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2021 Cigna.