



## Drug Coverage Policy

Effective Date..... 11/01/2024  
Coverage Policy Number ..... IP0083  
Policy Title..... Rukobia

# Human Immunodeficiency Virus – Rukobia

- Rukobia™ (fostemsavir extended-release tablets – ViiV/GlaxoSmithKline)

### **INSTRUCTIONS FOR USE**

*The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.*

## Cigna Healthcare Coverage Policy

### **OVERVIEW**

*Rukobia is a human immunodeficiency virus-1 (HIV-1) gp120-directed attachment inhibitor.<sup>1</sup> It is indicated in combination with other antiretroviral(s) [ARVs] for the treatment of HIV-1 infection in heavily treatment-experienced adults with **multidrug-resistant HIV-1 infection** failing their current ARV regimen due to resistance, intolerance, or safety considerations.*

### **Clinical Efficacy**

The efficacy of Rukobia was established in one ongoing, Phase III, multicenter, 96-week pivotal study in heavily treatment-experienced adults with HIV-1 infection failing their current ARV regimen (BRIGHTE; n = 371).<sup>2,5</sup> Eligible patients were  $\geq 18$  years of age and had failure of their current ARV regimen (baseline HIV-1 RNA  $\geq 400$  copies/mL), with no viable ARV combination therapy due to exhaustion of a least four of six ARV classes (i.e., nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, integrase inhibitors, protease inhibitors, CCR5 antagonists, and entry inhibitors). Exhaustion was defined as the elimination of all ARVs within a given class as a fully active option to pair with Rukobia because of resistance, previous adverse events, or unwillingness to use Fuzeon<sup>®</sup> (enfuvirtide subcutaneous injection). Patients were assigned to one of two cohorts, "randomized" or "non-randomized," based on the number of fully active available ARVs with which to construct an optimized background regimen. Patients in the randomized cohort had one or two available fully active ARVs while those in the non-randomized cohort had no available fully active ARVs. There were 15 patients who received Trogarzo<sup>®</sup> (ibalizumab-uiyk intravenous injection) in combination with Rukobia in the original study (after approval of Rukobia, in 2018, 6 patients in the randomized cohort added Trogarzo to their optimized background therapy). At Week 48 (all patients on Rukobia + optimized background therapy), 54% of patients achieved viral suppression (HIV-1 RNA  $< 40$  copies/mL) and CD4 T-cell count increased to a mean of 139 cells/mm<sup>3</sup> (median baseline 99 cells/mm<sup>3</sup>). At Week 96, viral suppression was maintained or improved (60% of patients in the randomized cohort and 37% of patients in the non-randomized cohort).<sup>5</sup> Virologic response rates (in the intent-to-treat population) at Week 240 decreased to 45% in the randomized cohort and to 22% in the non-randomized cohort; however, this included 24 patients (n = 19 in the randomized cohort and n = 5 in the non-randomized cohort) who did not have virologic data due to Coronavirus-19-related disruptions to care. At Week 240, in the observed analysis, HIV-1 RNA was  $< 40$  copies/mL in 82% of patients in the randomized cohort and in 66% of patients in the non-randomized cohort and was  $< 400$  copies/mL in 95% and 80% of patients in the randomized and non-randomized cohorts, respectively. Mean CD4 T-cell count increased through Week 192 in the randomized cohort and remained similar at Week 240; in the non-randomized cohort, CD4 T-cell count increased through Week 240. A safety analysis reported additional findings.<sup>7</sup> At Week 102, 58% of patients with CD4 T-cell count  $< 200$  cells/mm<sup>3</sup> at baseline had on-treatment CD4 T-cell count reported at Week 192, and 75% of these patients had a CD4 T-cell count  $\geq 200$  cells/mm<sup>2</sup>. In addition, acquired immunodeficiency syndrome-defining events were lower among patients with CD4 T-cell count  $\geq 200$  cells/mm<sup>3</sup> vs.  $< 200$  cells/mm<sup>3</sup>.

## Guidelines

According to the Department of Health and Human Services Guidelines (February 27, 2024) for the use of antiviral agents in adults and adolescents with HIV infection, treatment-experienced patients with ongoing detectable viremia who lack sufficient treatment options to construct a fully suppressive regimen may be candidates for Trogarzo, Rukobia, or Sunlenca<sup>®</sup> (lenacapavir tablets/subcutaneous injection).<sup>3</sup> Patients who continue to have detectable viremia and who lack sufficient treatment options to construct a fully suppressive regimen may also be candidates for research studies or expanded access programs, or they may qualify for single-patient access to an investigational new drug as specified in FDA regulations. The goal of therapy is viral resuppression, if possible; otherwise, to keep the viral load as low as possible and CD4 T-cell count as high as possible. CD4 T-cell count is recommended to be monitored at entry into care, when switching or modifying ARVs, and then every 3, 6, or 12 months depending on CD4 T-cell count and the duration of viral suppression. The CD4 count response to ARV therapy varies widely, but a poor CD4 response in a patient with viral suppression is rarely an indication for modifying a treatment regimen.

The International Antiviral Society-USA recommendations (2022) for the treatment and prevention of HIV in adults recognize Rukobia in the setting of integrase strand-transfer inhibitor (INSTI) resistance. If INSTI resistance is relatively limited and a new antiviral regimen is to include an

INSTI, the regimen should also include at least one and preferably two other fully active drugs, optimally from drug classes not previously used which may include among other agents, Rukobia.<sup>4</sup>

Consensus recommendations for the use of novel ARVs in individuals with HIV who are heavily treatment-experienced and/or have multidrug-resistant HIV-1 endorsed by the American Academy of HIV Medicine and the American College of Clinical Pharmacology (2024) recommend adding Rukobia to an optimized background regimen that includes at least one other active drug in patients with multidrug resistant HIV-1 who are heavily treatment-experienced and are unable to achieve or maintain viral suppression on their current ARV regimen.<sup>8</sup> If an active drug cannot be included in the optimized background regimen, then the regimen should include partially active agents (preferably several). Prior treatment with Trogarzo should not impact candidacy for Rukobia.

## Medical Necessity Criteria

**Rukobia is considered medically necessary when the following criteria are met:**

### FDA-Approved Indication

- 1. Human Immunodeficiency Virus (HIV)-1 Infection.** Approve for the duration noted if the patient meets ONE of the following (A or B):
  - A) Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, and v):
    - i.** Patient is  $\geq 18$  years of age; AND
    - ii.** According to the prescriber, the patient is failing a current antiretroviral regimen for HIV; AND
    - iii.** According to the prescriber, the patient has exhausted at least FOUR of the following antiretroviral classes defined as elimination of all antiretrovirals within a given class due to demonstrated or projected resistance to the agent(s) in that class OR due to significant intolerance (FOUR of a, b, c, d, e, or f):
      - a)** Nucleoside reverse transcriptase inhibitor; OR  
Note: Examples of nucleoside reverse transcriptase inhibitors include abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir disoproxil fumarate, tenofovir alafenamide, zidovudine.
      - b)** Non-nucleoside reverse transcriptase inhibitor; OR  
Note: Examples of non-nucleoside reverse transcriptase inhibitor include delavirdine, efavirenz, etravirine, nevirapine, nevirapine XR, rilpivirine.
      - c)** Protease inhibitor; OR  
Note: Examples of protease inhibitors include atazanavir, darunavir, fosamprenavir, indinavir, nelfinavir, ritonavir, saquinavir, tipranavir.
      - d)** Fusion inhibitor; OR  
Note: Examples of fusion inhibitors include Fuzeon (enfuvirtide subcutaneous injection).
      - e)** Integrase strand transfer inhibitor; OR  
Note: Examples of integrase strand-transfer inhibitors include raltegravir, dolutegravir, elvitegravir.
      - f)** CCR5 antagonist; AND  
Note: Examples of CCR5 antagonists include Selzentry (maraviroc tablets).
    - iv.** The medication will be taken in combination with an optimized antiviral background regimen including one or more other antiretroviral agents; AND
    - v.** The medication is prescribed by or in consultation with a physician who specializes in the treatment of HIV infection.

- B) Patient is Currently Receiving Rukobia.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
- i. The medication will continue to be taken in combination with an optimized antiretroviral background regimen including one or more other antiretroviral agents; AND
  - ii. Patient has responded to a Rukobia-containing regimen, as determined by the prescriber. Note: Examples of a response are HIV RNA < 40 cells/mm<sup>3</sup>, HIV-1 RNA ≥ 0.5 log<sub>10</sub> reduction from baseline in viral load, improvement, or stabilization of CD4 T-cell count.

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

## Conditions Not Covered

Any other use is considered experimental, investigational, or unproven (criteria will be updated as new published data are available).

## References

1. Rukobia™ extended-release tablets [prescribing information]. Research Triangle Park, NC: ViiV/GlaxoSmithKline; February 2024.
2. Kozal M, Aberg J, Pialoux G, et al. Fostemsavir in adults with multidrug-resistant HIV-1 infection. *N Engl J Med.* 2020;382(13):1232-1243.
3. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in adults and adolescents with HIV. Department of Health and Human Services. Last Updated: February 24, 2024. Available at: [clinicalinfo.hiv.gov/sites/default/files/guidelines/documents/adult-adolescent-arv/guidelines-adult-adolescent-arv.pdf](https://clinicalinfo.hiv.gov/sites/default/files/guidelines/documents/adult-adolescent-arv/guidelines-adult-adolescent-arv.pdf). Accessed on: July 10, 2024.
4. Ghandi RT, Bedimo R, Hoy JF, et al. Antiviral drugs for treatment and prevention of HIV infection in adults. 2022 recommendations of the International Antiviral Society-USA panel. *JAMA.* 2023;329(1):63-84.
5. Lataillade M, Lalezari J, Kozal M, et al. Safety and efficacy of the HIV-1 attachment inhibitor prodrug fostemsavir in heavily treatment-experienced individuals: week 96 results of the phase 3 BRIGHTE study. *Lancet HIV.* 2020; 7(11): e740-e751.
6. Aberg JA, Shepard B, Wang M, et al. Week 240 efficacy and safety of fostemsavir plus optimized background therapy in heavily treatment-experienced adults with HIV-1. *Infect Dis Ther.* 2023; 12:2321-2335.
7. Ilibre JM, Aberg JA, Walmsley S, et al. Long-term safety, and impact of immune recovery in heavily treatment-experienced adults receiving fostemsavir for up to 5 years in the phase 3 BRIGHTE study. *Front Immunol.* 2024; 15:1394664.
8. Cluck DB, Chastain DB, Murray M, et al. Consensus recommendations for the use of novel antiretrovirals in persons with HIV who are heavily treatment-experienced and/or have multidrug-resistant HIV-1: Endorsed by the American Academy of HIV Medicine, American College of Clinical Pharmacy. *Pharmacotherapy.* 2024;44;360.

## Revision Details

Type of Revision	Summary of Changes	Date
------------------	--------------------	------

Annual Revision	<p><b>Policy Title:</b>  <b>Updated from</b> “Fostemsavir” <b>to</b> “Human Immunodeficiency Virus – Rukobia.”</p> <p><b>Human Immunodeficiency Virus-1 Infection.</b>  <b>Updated from</b> “Human Immunodeficiency Virus Infection” <b>to</b> “Human Immunodeficiency Virus-1 Infection”.</p> <p><u>Initial Therapy</u>  <b>Updated</b> approval duration <b>from</b> “12 months” <b>to</b> “6 months”.</p> <p><b>Updated from</b> “History of multi-drug resistant Human Immunodeficiency Virus” <b>to</b> “According to the prescriber, the patient is failing a current antiretroviral regimen for HIV”.</p> <p><b>Added</b> criterion, “According to the prescriber, the patient has exhausted at least FOUR of the following antiretroviral classes defined as elimination of all antiretrovirals within a given class due to demonstrated or projected resistance to the agent(s) in that class OR due to significant intolerance (FOUR of a, b, c, d, e, <u>or</u> f):</p> <p>a) Nucleoside reverse transcriptase inhibitor; OR  Note: Examples of nucleoside reverse transcriptase inhibitors include abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir disoproxil fumarate, tenofovir alafenamide, zidovudine.</p> <p>b) Non-nucleoside reverse transcriptase inhibitor; OR  Note: Examples of non-nucleoside reverse transcriptase inhibitor include delavirdine, efavirenz, etravirine, nevirapine, nevirapine XR, rilpivirine.</p> <p>c) Protease inhibitor; OR  Note: Examples of protease inhibitors include atazanavir, darunavir, fosamprenavir, indinavir, nelfinavir, ritonavir, saquinavir, tipranavir.</p> <p>d) Fusion inhibitor; OR  Note: Examples of fusion inhibitors include Fuzeon (enfuvirtide subcutaneous injection).</p> <p>e) Integrase strand transfer inhibitor; OR  Note: Examples of integrase strand-transfer inhibitors include raltegravir, dolutegravir, elvitegravir.</p> <p>f) CCR5 antagonist; AND  Note: Examples of CCR5 antagonists include Selzentry (maraviroc tablets)”.</p> <p><u>Patient is Currently Receiving Rukobia.</u>  <b>Updated from</b> “<u>Reauthorization Criteria.</u> Fostemsavir extended-release (Rukobia) tablets are considered medically necessary for continued use when initial criteria are met AND there is documentation of beneficial response” <b>to</b>  <u>Patient is Currently Receiving Rukobia.</u> Approve for 1 year if the patient meets BOTH of the following (i and ii):  i. The medication will continue to be taken in combination with an optimized antiviral background regimen including one or more other antiretroviral agents; AND</p>	11/01/2024
-----------------	--	------------

	<p>ii. Patient has responded to a Rukobia-containing regimen, as determined by the prescriber. Note: Examples of a response are HIV RNA &lt; 40 cells/mm3, HIV-1 RNA ≥ 0.5 log10 reduction from baseline in viral load, improvement, or stabilization of CD4 T-cell count.”</p>	
--	---	--

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.