



Drug Coverage Policy

Effective Date.....06/01/2024

Coverage Policy Number.....IP0158

Policy Title..... Zepatier

Hepatitis C – Zepatier

- Zepatier® (grazoprevir/elbasvir tablets – Merck)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Cigna Healthcare Coverage Policy

OVERVIEW

Zepatier, an oral fixed-dose combination tablet containing grazoprevir, a second generation protease inhibitor and elbasvir, an NS5A inhibitor, is indicated with or without ribavirin for the treatment of genotypes 1 and 4 **chronic hepatitis C virus** (HCV) in adults and pediatric patients ≥ 12 years of age or weighing at least 30 kg.¹

Safety

Zepatier is contraindicated in patients with Child-Pugh B or C liver disease (decompensated cirrhosis). Zepatier is also contraindicated with inhibitors of organic anion transporting polypeptides 1B1/3 that are known or expected to significantly increase grazoprevir plasma concentrations, strong inducers of cytochrome P450 3A, and efavirenz.

Dosing

The duration of treatment is outlined below (Table 1) and is dependent on the patient population. Prior to initiating Zepatier in patients with genotype 1a infection, testing for the NS5A resistance associated polymorphism is recommended to guide treatment duration. In patients with genotype 1a and this polymorphism present at baseline, 12 weeks of treatment with Zepatier resulted in lower rates of sustained viral response 12 weeks after treatment completion relative to patients with genotype 1a without the presence of this baseline polymorphism.

Table 1. Recommended Zepatier Dosage Regimens for the Treatment of Genotype 1 or 4 Chronic HCV.¹

Genotype	Treatment History	Baseline NS5A Polymorphism	Treatment Regimen	Treatment Duration
1a	TN/PR-experienced* without NS5A polymorphisms [†]	No [†]	Zepatier	12 weeks
1a	TN/PR-experienced* <u>with</u> baseline NS5A polymorphisms [†]	Yes [†]	Zepatier + ribavirin	16 weeks
1a [§] or 1b	PR + HCV PI-experienced ^β	NA	Zepatier + ribavirin	12 weeks
1b	TN/TE*	NA	Zepatier	12 weeks
4	TN	NA	Zepatier	12 weeks
4	PR-experienced*	NA	Zepatier + ribavirin [†]	16 weeks

HCV – Hepatitis C virus; TN – Treatment naïve; PR – Pegylated interferon/ribavirin; * Patients who have failed treatment with PR; [†] NS5A resistance-associated polymorphisms at amino acid positions 28, 30, 31, or 93;[§] The optimal Zepatier-based treatment regimen and duration of therapy for PR + HCV protease inhibitor-experienced genotype 1a-infected patients with one or more baseline NS5A resistance-associated polymorphisms at positions 28, 30, 31, and 93 has not been established; PI – Protease inhibitor; ^β Patients who have failed treatment with PR + and NS3/4A PI (i.e., Victrelis® [boceprevir capsules], Incivek® [telaprevir tablets], or Olysio® [simeprevir capsules]); TE – Treatment-experienced; NA – Not applicable.

Guidelines

According to the American Association for the Study of Liver Diseases (AASLD) [December 2023], NS5A resistance-associated substitution RAS testing is recommended for genotype 1a-infected, treatment-naïve or -experienced patients being considered for Zepatier.² If present, a different regimen should be considered. Zepatier is recognized as an alternative regimen in treatment-naïve patients with Genotype 1a with or without compensated cirrhosis, and a recommended treatment option in patients with genotype 1b or 4 chronic HCV with or without compensated cirrhosis in guidelines. It is also recognized as an alternative regimen in treatment-naïve and non-direct-acting antiviral-experienced kidney transplant patients with genotype 1 or 4 with or without compensated cirrhosis. The pediatric portion of the guidelines have not been updated since the approval of the lower age indication approved with Zepatier.

Medical Necessity Criteria

Zepatier is considered medically necessary when the following criteria are met:

FDA-Approved Indications

1. **Chronic Hepatitis C Virus (HCV) Genotype 1a.** Approve for the duration noted if the patient meets the following (A, B, C, and D):
 - A) Patient meets ONE of the following (i or ii):
 - i. Patient is \geq 12 years of age; OR
 - ii. Patient weighs \geq 30 kg; AND
 - B) Patient meets ONE of the following (i or ii):
 - i. Approve for 12 weeks if the patient meets ONE of the following (a or b):
 - a) Patient meets both of the following [(1) and (2)]:
 - (1) Patient is treatment-naïve, OR patient has previously been treated with pegylated interferon + ribavirin *only*; AND
 - (2) Patient does NOT have a baseline NS5A polymorphism at ONE (or more) of the following the amino acid positions: 28, 30, 31, or 93; OR
 - b) Patient meets both of the following [(1) and (2)]:
 - (1) Patient has previously been treated with pegylated interferon + ribavirin and an HCV protease inhibitor; AND
 - (2) The medication will be prescribed in combination with ribavirin; OR
 - ii. Approve for 16 weeks if the patient meets the following (a, b, and c):
 - a) Patient meets one of the following [(1) or (2)]:
 - (1) Patient is treatment-naïve; OR
 - (2) Patient has previously been treated with pegylated interferon + ribavirin *only*; AND
 - b) Patient has a baseline NS5A polymorphism at ONE (or more) of the following amino acid positions: 28, 30, 31, or 93; AND
 - c) The medication will be prescribed in combination with ribavirin; AND
 - C) The medication is prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician; AND
 - D) Preferred product criteria is met for the product(s) as listed in the below table(s)
2. **Chronic Hepatitis C Virus (HCV) Genotype 1b.** Approve for 12 weeks if the patient meets the following (A, B, C, and D):
 - A) Patient meets ONE of the following (i or ii):
 - i. Patient is \geq 12 years of age; OR
 - ii. Patient weighs \geq 30 kg; AND
 - B) Patient meets ONE of the following (i or ii):
 - i. Patient meets one of the following (a or b):
 - a) Patient is treatment-naïve; OR
 - b) Patient has previously been treated with pegylated interferon + ribavirin *only*; OR
 - ii. Patient meets the following (a and b):
 - a) Patient has previously been treated with pegylated interferon + ribavirin + an HCV protease inhibitor; AND
 - b) The medication will be prescribed in combination with ribavirin; AND
 - C) The medication is prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician; AND
 - D) Preferred product criteria is met for the product(s) as listed in the below table(s)
3. **Chronic Hepatitis C Virus (HCV) Genotype 4.** Approve for the duration noted if the patient meets the following (A, B, C, and D):
 - A) Patient meets ONE of the following (i or ii):
 - i. Patient is \geq 12 years of age; OR
 - ii. Patient weighs \geq 30 kg; AND
 - B) Patient meets ONE of the following (i or ii):
 - i. Patient is treatment-naïve: Approve for 12 weeks; OR
 - ii. Approve for 16 weeks if the patient meets both of the following (a and b):

- a) Patient has previously been treated with pegylated interferon and ribavirin; AND
- b) The medication will be prescribed in combination with ribavirin; AND
- c) The medication is prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician; AND
- d) Preferred product criteria is met for the product(s) as listed in the below table(s)

Other Uses with Supportive Evidence

4. Patient is Currently Receiving Zepatier. Approve for an indication or condition addressed as an approval in the Recommended Authorization Criteria section (FDA-Approved Indications). Approve the duration described above to complete a course of therapy (e.g., a patient who should receive 12 weeks and has received 3 weeks should be approved for 9 weeks to complete their 12-week course).

Employer Plans:

Product	Criteria
Zepatier (grazoprevir/elbasvir)	Failure, contraindication, or intolerance to ONE of the following: <ol style="list-style-type: none"> 1. Epclusa (sofosbuvir/velpatasvir) [may require prior authorization] 2. Harvoni (ledipasvir/sofosbuvir) [may require prior authorization] 3. Vosevi (sofosbuvir/velpatasvir/voxilaprevir) [may require prior authorization]

Individual and Family Plans:

Product	Criteria
Zepatier (grazoprevir/elbasvir)	Failure, contraindication, or intolerance to ONE of the following: <ol style="list-style-type: none"> 1. Epclusa (sofosbuvir/velpatasvir) [may require prior authorization] 2. sofosbuvir/velpatasvir [may require prior authorization] 3. Harvoni (ledipasvir/sofosbuvir) [may require prior authorization] 4. ledipasvir/sofosbuvir [may require prior authorization]

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

Conditions Not Covered

Any other use is considered experimental, investigational, or unproven, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Hepatitis C Virus (HCV), Child-Pugh Class B or Child-Pugh Class C Liver Disease (Moderate or Severe Hepatic Impairment).** Zepatier is contraindicated in patients with moderate or severe hepatic impairment (Child-Pugh Class B or C).¹

2. **Hepatitis C Virus (HCV) [Any Genotype], Combination with Any Other Direct-Acting Antivirals (Not Including Ribavirin).** Zepatier provides a complete antiviral regimen for patients with genotype 1 and 4 chronic HCV.
3. **Pediatric Patients (Age < 12 Years or < 30 kg).** The safety and efficacy of Zepatier have not been established in pediatric patients < 12 years of age or < 30 kg.¹ Guidelines recommend Harvoni (ledipasvir/sofosbuvir tablets) in pediatric patients with genotypes 1 or 4 chronic HCV.²
4. **Retreatment with Zepatier in Patients Who Have Previously Received Zepatier.** Zepatier is not recommended. This includes retreatment in prior null responders, prior partial responders, prior relapse patients, and patients who have not completed a course of therapy due to an adverse reaction or for other reasons.

References

1. Zepatier® tablets [prescribing information]. Whitehouse Station, NJ: Merck; December 2021.
2. American Association for the Study of Liver Diseases and the Infectious Diseases Society of America. Testing, managing, and treating hepatitis C. Updated December 19, 2023. Available at: <http://www.hcvguidelines.org>. Accessed on February 1, 2024.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	<p>Conditions Not Recommended for Approval: Life Expectancy Less Than 12 Months Due to Non-Liver Related Comorbidities. This condition was removed.</p> <p>Medical Necessity Criteria: Does not have decompensated cirrhosis (Child-Pugh B or C). This criterion was removed.</p>	06/01/2024

The policy effective date is in force until updated or retired.

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.