



## Drug Coverage Policy

Effective Date.....12/01/2024  
Coverage Policy Number.....IP0259  
Policy Title..... Multiple Sclerosis –  
Fingolimod

# Multiple Sclerosis – Fingolimod

Gilenya® (fingolimod capsules - Novartis, generic)

### INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

## Cigna Healthcare Coverage Policy

### Overview

Fingolimod, a sphingosine 1-phosphate receptor modulator, is indicated for the treatment of patients with relapsing forms of **multiple sclerosis** (MS), to include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease in patients  $\geq 10$  years of age.<sup>1</sup>

### Disease Overview

MS is a chronic, inflammatory, demyelinating, autoimmune disease of the central nervous system (CNS) that impacts almost 1,000,000 people in the US.<sup>2-4</sup> The condition is marked by inflammation and demyelination, as well as degenerative alterations. Patients usually experience relapses and remissions in their neurological symptoms. For most patients, the onset of MS symptoms occurs when patients are 20 to 40 years of age; however, children can get MS and new onset disease can occur in older adults. The MS disease course is heterogeneous but has some patterns. Approximately 85% to 90% of patients have a relapsing pattern at onset. However, this transitions over time in patients who are untreated to a worsening with very few or no relapses or magnetic resonance imaging (MRI) activity (secondary progressive MS). Around 10% to 15% of patients have a steady progression of symptoms over time (primary progressive MS), marked by some clinical manifestations or by MRI activity. Primary progressive MS is generally diagnosed in patients on the upper level of the typical age range (e.g., almost 40 years of age) and the distribution is equivalent among the two genders. Advances in the understanding of the MS disease process, as well as in MRI technology, spurred updated disease course descriptions in 2013,<sup>5</sup> as well as in 2017.<sup>6</sup> The revised disease courses are clinically isolated syndrome, relapsing remitting MS, primary progressive MS, and secondary progressive MS.<sup>2-6</sup> Clinically isolated syndrome is now more recognized among the course descriptions of MS. It is the first clinical presentation of MS that displays characteristics of inflammatory demyelination that may possibly be MS but has yet to fulfill diagnostic criteria. It is notable that the other MS designations can be further characterized considering whether patients have active disease (or not active), as well as if disease is worsening or stable. Disability in MS is commonly graded on the deterioration of mobility per the Expanded Disability Status Scale (EDSS) an ordinal scale that ranges from 0 to 10, with higher scores indicating greater disability.

### **Guidelines**

In September 2019, a consensus paper was updated by the MS Coalition that discusses the use of disease-modifying therapies in MS.<sup>2</sup> Many options from various disease classes, involving different mechanisms of action and modes of administration, have shown benefits in patients with MS.<sup>2</sup> The American Academy of Neurology has practice guidelines regarding disease-modifying therapies for adults with MS.<sup>7</sup> The guidelines cites fingolimod as one of the agents to consider for patients with MS who have highly active disease.

### **Safety**

The initiation of fingolimod leads to decreases in heart rate.<sup>1</sup> After the first dose of fingolimod, the heart rate decreases are noted within an hour and generally are greatest at 6 hours, although the effects can be observed 24 hours after the first dose in some patients. The first dose of fingolimod should be given in a setting with resources to appropriately manage symptomatic bradycardia. Observe patients for 6 hours after the first fingolimod dose for signs and symptoms of bradycardia. Patients with prolonged QTc interval at baseline or during the observation period, or taking medications with known risks of torsades de pointes, should be observed overnight with continuous electrocardiographic (ECG) monitoring. When restarting fingolimod after discontinuation for more than 14 days after the first treatment month, perform first-dose monitoring. There are several contraindications for use which mainly include patients with background cardiovascular disease. Fingolimod is associated with serious toxicities such as decreased heart rate and/or atrioventricular condition after the first dose; an increased risk of infections; macular edema; pulmonary toxicity; and elevated liver enzymes. Cases of progressive multifocal leukoencephalopathy (PML) have occurred in patients who were given fingolimod in the postmarketing setting.

## **Medical Necessity Criteria**

**Documentation:** Documentation is required where noted in the criteria. Documentation may include, but not limited to, chart notes, laboratory tests, claims records, and/or other information.

[Click here for Individual and Family Plan Criteria](#)

**For Employer Group Benefit Plans:**

**Gilenya is considered medically necessary when the following are met:**

**FDA-Approved Indication**

**1. Multiple Sclerosis.** Approve for the duration noted below if the patient meets one of the following (A or B):

**A) Initial Therapy.** Approve for 1 year if the patient meets the following (i, ii, iii, and iv):

**i.** Documentation the patient has a relapsing form of multiple sclerosis; AND  
Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.

**ii.** Patient is  $\geq 10$  years of age; AND

**iii.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; AND

**iv.** Preferred product criteria is met for the product(s) as listed in the below table(s);OR

**B) Patient is Currently Receiving Fingolimod for  $\geq 1$  Year.** Approve for 1 year if the patient meets the following (i, ii, iii and iv):

**i.** Documentation the patient has a relapsing form of multiple sclerosis; AND  
Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.

**ii.** Patient is  $\geq 10$  years of age; AND

**iii.** Patient meets one of the following (a or b):

**a)** Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR

Note: Examples include stabilization or reduced worsening in disease activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2 lesions]; stabilization or reduced worsening on the Expanded Disability State Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the fatigue symptom and impact questionnaire-relapsing multiple sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or 12-Item MS Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume loss.

**b)** Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND

**iv.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.

**For Individual and Family Plans:**

**Fingolimod products (Gilenya and fingolimod capsules) are considered medically necessary when the following are met:**

**FDA-Approved Indication**

**1. Multiple Sclerosis.** Approve for the duration noted below if the patient meets one of the following (A or B):

- A) Initial Therapy.** Approve for 1 year if the patient meets the following (i, ii, iii, and iv):
- i.** Documentation the patient has a relapsing form of multiple sclerosis; AND  
Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
  - ii.** Patient is  $\geq 10$  years of age; AND
  - iii.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; AND
  - iv.** Preferred product criteria is met for the product(s) as listed in the below table(s);OR
- B) Patient is Currently Receiving Fingolimod for  $\geq 1$  Year.** Approve for 1 year if the patient meets the following (i, ii, iii and iv):
- i.** Documentation the patient has a relapsing form of multiple sclerosis; AND  
Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
  - ii.** Patient is  $\geq 10$  years of age; AND
  - iii.** Patient meets one of the following (a or b):
    - c)** Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR  
Note: Examples include stabilization or reduced worsening in disease activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2 lesions]; stabilization or reduced worsening on the Expanded Disability State Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the fatigue symptom and impact questionnaire-relapsing multiple sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or 12-Item MS Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume loss.
    - d)** Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND
  - iv.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.

**Employer Plans:**

<b>Product</b>	<b>Criteria</b>
<b>Gilenya 0.5 mg</b> (fingolimod 0.5 mg capsule)	Documented trial of the bioequivalent generic product <b>fingolimod capsules</b> AND cannot take due to a formulation difference in the inactive ingredient(s) [for example, difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which, per the prescriber, would result in a significant allergy or serious adverse reaction

**Individual and Family Plans:**

<b>Product</b>	<b>Criteria</b>
<b>Gilenya 0.5 mg</b> (fingolimod 0.5 mg capsule)	Documented trial of the bioequivalent generic product <b>fingolimod capsules</b> [may require prior authorization] AND cannot take due to a formulation difference in the inactive ingredient(s) [for example, difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which, per the prescriber, would result in a significant allergy or serious adverse reaction

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based

literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

## Conditions Not Covered

Any other use is considered experimental, investigational, or unproven, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Concurrent Use with Other Disease-Modifying Agents Used for Multiple Sclerosis.** These agents are not indicated for use in combination (See [Appendix](#) for examples). Additional data are required to determine if use of disease-modifying multiple sclerosis agents in combination is safe and provides added efficacy.
- 2. Non-Relapsing Forms of Multiple Sclerosis.** In the INFORMS trial, fingolimod did not slow disease progression in patients with primary progressive multiple sclerosis.<sup>8</sup>  
Note: An example of a non-relapsing form of multiple sclerosis is primary progressive multiple sclerosis.

## References

1. Gilenya<sup>®</sup> capsules [prescribing information]. East Hanover, NJ: Novartis; September 2023.
2. A Consensus Paper by the Multiple Sclerosis Coalition. The use of disease-modifying therapies in multiple sclerosis. September 2019. Available at: [http://www.nationalmssociety.org/getmedia/5ca284d3-fc7c-4ba5-b005-ab537d495c3c/DMT\\_Consensus\\_MS\\_Coalition\\_color](http://www.nationalmssociety.org/getmedia/5ca284d3-fc7c-4ba5-b005-ab537d495c3c/DMT_Consensus_MS_Coalition_color). Accessed on November 4, 2023.
3. McGinley MP, Goldschmidt C, Rae-Grant AD. Diagnosis and treatment of multiple sclerosis. A review. *JAMA*. 2021;325(8):765-779.
4. No authors listed. Drugs for multiple sclerosis. *Med Lett Drugs Ther*. 2021;63(1620):42-48.
5. Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. *Neurology*. 2014;83:278-286.
6. Thompson AJ, Banwell BL, Barkhof F, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol*. 2018;17(2):162-173.
7. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: disease-modifying therapies for adults with multiple sclerosis. Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90:777-788.
8. Lublin F, Miller DH, Freedman MS, et al, on behalf of the INFORMS Study Investigators. Oral fingolimod in primary progressive multiple sclerosis (INFORMS): a phase 3, randomized, double-blind, placebo-controlled trial. *Lancet*. 2016;387:1075-1084.

### Appendix

Medication	Mode of Administration
Aubagio <sup>®</sup> (teriflunomide tablets, generic)	Oral
Avonex <sup>®</sup> (interferon beta-1a intramuscular injection)	Injection (self-administered)
Bafiertam <sup>®</sup> (monomethyl fumarate delayed-release capsules)	Oral
Betaseron <sup>®</sup> (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Briumvi <sup>®</sup> (ublituximab-xiij intravenous infusion)	Intravenous infusion

Copaxone® (glatiramer acetate subcutaneous injection, generic)	Injection (self-administered)
Extavia® (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Gilenya® (fingolimod capsules, generic)	Oral
Glatopa® (glatiramer acetate subcutaneous injection)	Injection (self-administered)
Kesimpta® (ofatumumab subcutaneous injection)	Injection (self-administered)
Lemtrada® (alemtuzumab intravenous infusion)	Intravenous infusion
Mavenclad® (cladribine tablets)	Oral
Mayzent® (siponimod tablets)	Oral
Ocrevus® (ocrelizumab intravenous infusion)	Intravenous infusion
Ocrevus Zunovo™ (ocrelizumab and hyaluronidase-ocsq subcutaneous injection)	Subcutaneous Injection (not self-administered)
Plegridy® (peginterferon beta-1a subcutaneous or intramuscular injection)	Injection (self-administered)
Ponvory® (ponesimod tablets)	Oral
Rebif® (interferon beta-1a subcutaneous injection)	Injection (self-administered)
Tascenso ODT® (fingolimod orally disintegrating tablets)	Oral
Tecfidera® (dimethyl fumarate delayed-release capsules, generic)	Oral
Tyruko® (natalizumab-sztn intravenous infusion)	Intravenous infusion
Tysabri® (natalizumab intravenous infusion)	Intravenous infusion
Vumerity® (diroximel fumarate delayed-release capsules)	Oral
Zeposia® (ozanimod capsules)	Oral

## Revision Details

Type of Revision	Summary of Changes	Date
Selected Revision	<p><b>Added</b> a definition for documentation.</p> <p><b>Added</b> a requirement for a patient to be <math>\geq 10</math> years of age.</p> <p><b>Added</b> a specialist prescribing requirement.</p> <p><b>Added</b> criteria for patient Currently Receiving Fingolimod for <math>\geq 1</math> Year.</p> <p><b>Updated</b> the preferred product multi-source brand language to current standards.</p> <p>Ocrevus Zunovo was added to the Appendix.</p>	12/01/2024

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life

Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.