Drug Coverage Policy

Effective Date……9/15/2026
Coverage Policy Number……IP0265
Policy Title……(Injectable – Beta Interferon) - Rebif

Multiple Sclerosis (Injectable – Beta Interferon) – Rebif

- Rebif® (interferon beta-1a subcutaneous injection - EMD Serono)

INSTRUCTIONS FOR USE
The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview
Rebif is indicated for the treatment of patients with relapsing forms of **multiple sclerosis** (MS) to include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease in adults.[1]

Guidelines
In September 2019, a consensus paper was updated by the MS Coalition that discusses the use of disease-modifying therapies in MS.[2] Many options from various disease classes, involving different mechanisms of action and modes of administration, have shown benefits in patients with MS. In addition, practice guidelines for disease-modifying therapies for adults with MS were published by the American Academy of Neurology in 2018.[3] It is noted that many disease-modifying therapies are available and have demonstrated efficacy for the prevention of clinical relapses and new magnetic resonance imaging (MRI)-detected lesions. A lack of response is difficult to define, as most patients with MS are not free of all disease activity. It is recommended clinicians evaluate various factors, such as clinical relapses, new MRI-detected lesions, or increased disability on examination, over a 1-year period of using a disease-modifying therapy.

Coverage Policy

Policy Statement
Prior Authorization is required for benefit coverage of Rebif. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Rebif as well as the monitoring required for adverse events and long-term efficacy, approval requires Rebif to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Rebif is considered medically necessary when the following are met:

FDA-Approved Indication

1. **Multiple Sclerosis.** Approve for 1 year if the patient meets ONE of the following (A or B):
   - **A)** Initial Therapy. Approve if the patient meets ALL of the following (i, ii and iii):
     - **i.** Patient has a relapsing form of multiple sclerosis; AND
       Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
     - **ii.** The medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; AND
     - **iii.** Preferred product criteria is met for the product(s) as listed in the below table(s) [Individual and Family Plans]; OR
   - **B)** Patient is Currently Receiving Rebif for ≥ 1 Year. Approve if the patient meets the ALL of the following (i, ii, and iii):
     Note: A patient who has received < 1 year of therapy or who is restarting therapy with Rebif should be considered under criterion 1A (Multiple Sclerosis, Initial Therapy).
     - **i.** Patient has a relapsing form of multiple sclerosis; AND
       Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
     - **ii.** Patient meets ONE of the following (a or b):
       - **a)** Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR
         Note: Examples include stabilization or reduced worsening in disease activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2

lesions]; stabilization or reduced worsening on the Expanded Disability State Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the Fatigue Symptom and Impact Questionnaire-Relapsing Multiple Sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or 12-Item Multiple Sclerosis Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume loss.

b) Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND

iii. The medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.

Individual and Family Plans:

Product	Criteria
Rebif (Interferon beta-1a)	Patient meets ONE of the following (1 or 2): **1.** Patient has tried and, according to the prescriber, the patient has had inadequate efficacy or significant intolerance to Avonex [requires prior authorization]; OR **2.** Patient is currently receiving Rebif

Conditions Not Covered

Rebif for any other use is considered not medically necessary including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

1. **Concurrent Use with Other Disease-Modifying Agents used for Multiple Sclerosis.** These agents are not indicated for use in combination (See Appendix for examples). Additional data are required to determine if use of disease-modifying multiple sclerosis agents in combination is safe and provides added efficacy.

2. **Non-Relapsing Forms of Multiple Sclerosis.** The efficacy of Rebif has not been established in patients with non-relapsing forms of multiple sclerosis.[1]
   Note: An example of a non-relapsing form of multiple sclerosis is primary progressive multiple sclerosis.

References

1. Rebif® subcutaneous injection [prescribing information]. Rockland, MA: EMD Serono; July 2023.
2. A Consensus Paper by the Multiple Sclerosis Coalition. The use of disease-modifying therapies in multiple sclerosis. September 2019.
3. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: disease-modifying therapies for adults with multiple sclerosis. Report of the guideline development, dissemination, and implementation subcommittee of the American Academy of Neurology. *Neurology*. 2018;90:777-788.

Revision Details

Summary of Changes	Review Date	Effective Date
Added a definition for documentation. **Added** a specialist prescribing requirement. **Added** criteria for patient Currently Receiving Rebif for ≥ 1 Year. **Removed** Employer Plans preferred product requirements. **Updated** Individual and Family Plans preferred product requirements. Ocrevus Zunovo was added to the Appendix.	10/24/2024	12/01/2024
The Policy name was changed to add "Injectable – Beta Interferon". **Added** a policy statement. **Removed** documentation requirements. **Updated** the Individual and Family Plans preferred product criteria. **Updated** the conditions not covered statement. **Removed** Extavia from the Appendix.	9/11/2025	11/01/2025
Multiple Sclerosis: Under the requirements for a patient who has received therapy for ≥ 1 year, a Note was added clarifying that a patient who has received < 1 year of therapy or who is restarting therapy should be reviewed under the Initial Therapy pathway. **Appendix:** It was noted that a generic for Mavenclad (cladribine tablets) is now available.	8/13/2026	9/15/2026

The policy effective date is in force until updated or retired.

APPENDIX

Medication	Mode of Administration
Aubagio® (teriflunomide tablets, generic)	Oral
Avonex® (interferon beta-1a intramuscular injection)	Injection (self-administered)
Bafiertam® (monomethyl fumarate delayed-release capsules)	Oral
Betaseron® (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Briumvi® (ublituximab-xiiy intravenous infusion)	Intravenous infusion
Copaxone® (glatiramer acetate subcutaneous injection, generic)	Injection (self-administered)
Gilenya® (fingolimod capsules, generic)	Oral
Glatopa® (glatiramer acetate subcutaneous injection)	Injection (self-administered)
Kesimpta® (ofatumumab subcutaneous injection)	Injection (self-administered)
Lemtrada® (alemtuzumab intravenous infusion)	Intravenous infusion
Mavenclad® (cladribine tablets, generic)	Oral
Mayzent® (siponimod tablets)	Oral
Ocrevus® (ocrelizumab intravenous infusion)	Intravenous infusion
Ocrevus Zunovo™ (ocrelizumab and hyaluronidase-ocsq subcutaneous injection)	Subcutaneous Injection (not self-administered)
Plegridy® (peginterferon beta-1a subcutaneous or intramuscular injection)	Injection (self-administered)
Ponvory® (ponesimod tablets)	Oral
Rebif® (interferon beta-1a subcutaneous injection)	Injection (self-administered)
Tascenso ODT® (fingolimod orally disintegrating tablets)	Oral
Tecfidera® (dimethyl fumarate delayed-release capsules, generic)	Oral
Tyruko® (natalizumab-sztn intravenous infusion)	Intravenous infusion
Tysabri® (natalizumab intravenous infusion)	Intravenous infusion
Vumerity® (diroximel fumarate delayed-release capsules)	Oral
Zeposia® (ozanimod capsules)	Oral

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2026 The Cigna Group.