



Drug Coverage Policy

Effective Date6/15/2025

Coverage Policy Number.....IP0499

Policy Title.....TOBI Podhaler

Antibiotics (Inhaled) – TOBI Podhaler

- TOBI® Podhaler (tobramycin inhalation powder – Novartis)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

OVERVIEW

TOBI Podhaler, an aminoglycoside antibiotic, is indicated for the management of **cystic fibrosis** (CF) patients with *Pseudomonas aeruginosa*.¹ Safety and efficacy have not been demonstrated in patients < 6 years of age, patients with forced expiratory volume in 1 second (FEV₁) < 25% or > 80% predicted, or patients colonized with *Burkholderia cepacia*.

Guidelines

The Cystic Fibrosis Foundation (CFF) Pulmonary Therapeutics Committee (2013) provides recommendations for the use of chronic medications in the management of CF lung disease.² In patients ≥ 6 years of age with CF and moderate-to-severe lung disease with *P. aeruginosa* persistently present in cultures of the airways, chronic use of inhaled tobramycin is strongly recommended to improve lung function and quality of life, and reduce exacerbations. For mild disease, the Committee recommends chronic use of inhaled tobramycin for patients ≥ 6 years of age with CF and *P. aeruginosa* persistently present in cultures of the airways, to reduce exacerbations.

The CFF published a systematic review of the literature regarding eradication of initial *P. aeruginosa* infections to develop guidelines for effective prevention (2014).³ The recommendations pertaining to inhaled antibiotics are as follows: 1) Inhaled antibiotic therapy is recommended for the treatment of initial or new growth of *P. aeruginosa* (the favored antibiotic regimen is tobramycin [300 mg twice daily] for 28 days); and 2) Prophylactic antipseudomonal antibiotics to prevent the acquisition of *P. aeruginosa* are not recommended.

The American Thoracic Society (ATS) published a clinical review (2013) of non-cystic fibrosis bronchiectasis on their webpage.⁴ The review lists nebulized antibiotics (e.g., colistin, gentamicin, tobramycin) as treatment options for the eradication or suppression of *P. aeruginosa*. The European Respiratory Society (ERS) have published guidelines (2017) for the management of adult bronchiectasis and recommends patients with a new isolate of *P. aeruginosa* be offered eradication antibiotic treatment which includes nebulized antibiotics (e.g., colistin, gentamicin, tobramycin).⁵ Neither the ATS nor the ERS guidelines include TOBI Podhaler as a treatment option for bronchiectasis and no clinical trials have been published with TOBI Podhaler for treatment of non-cystic fibrosis bronchiectasis.

Medical Necessity Criteria

Policy Statement

Prior Authorization is recommended for prescription benefit coverage of TOBI Podhaler. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with TOBI Podhaler as well as the monitoring required for adverse events and long-term efficacy, approval requires TOBI Podhaler to be prescribed by a physician who has consulted with or who specializes in the condition.

TOBI Podhaler is considered medically necessary when the following criteria are met:

FDA-Approved Indication

- 1. Cystic Fibrosis.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
 - A)** Patient is ≥ 6 years of age; AND
 - B)** Patient has *Pseudomonas aeruginosa* in culture of the airway; AND
Note: Examples of culture of the airway include sputum culture, oropharyngeal culture, bronchoalveolar lavage culture.
 - C)** The medication is prescribed by or in consultation with a pulmonologist or a physician who specializes in the treatment of cystic fibrosis; AND
 - D)** Preferred product criteria are met for the product(s) as listed in the below table

Other Uses with Supportive Evidence

- 2. Continuation of TOBI Podhaler (not Cystic Fibrosis).** Approve for 1 month if the patient meets BOTH of the following (A and B)
- A)** Patient was started on TOBI Podhaler and is continuing a course of therapy; AND
- B)** Preferred product criteria are met for the product(s) as listed in the below table

Individual and Family Plans:

Product	Criteria
TOBI Podhaler (tobramycin inhalation powder)	ONE of the following: <ol style="list-style-type: none"> Intolerance to ONE of the following: <ol style="list-style-type: none"> tobramycin 300 mg/5 mL inhalation solution (generic for TOBI) tobramycin pak 300mg/5ml inhalation solution Currently taking TOBI Podhaler

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

Conditions Not Covered

Any other use is considered experimental, investigational, or unproven (criteria will be updated as new published data are available).

References

- TOBI® Podhaler inhalation powder [prescribing information]. East Hanover, NJ: Novartis; February 2023.
- Mogayzel PJ, Naureckas ET, Robinson KA, et al. Cystic Fibrosis Pulmonary Guidelines. Chronic Medications for Maintenance of Lung Health. *Am J Respir Crit Care Med*. 2013; 187:680-689.
- Mogayzel PJ, Naureckas ET, Robinson KA, et al; and the Cystic Fibrosis Foundation Pulmonary Clinical Practice Guidelines Committee. Pharmacologic approaches to prevention and eradication of initial *Pseudomonas aeruginosa* infection. *Ann Am Thorac Soc*. 2014;11(10):1640-1650.
- McShane PJ, Naureckas ET, Tino G, Strek ME. Non-cystic fibrosis bronchiectasis. *Am J Respir Crit Care Med*. 2013; 188:647-656.
- Polverino E, Goeminne PC, McDonnell, et al. European Respiratory Society guidelines for the management of adult bronchiectasis. *Eur Respir J*. 2017; 50:1700629.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	Policy Name Change: Updated Policy Name from "Tobramycin Inhalation Powder" to "Antibiotics (Inhaled) – TOBI Podhaler." Added preferred product requirement criteria for Individual and Family Plans. Continuation of TOBI Podhaler: Changed approval duration from 12 months to 1 month.	06/15/2024

	Removed the requirement that the medication be prescribed by or in consultation with a pulmonologist.	
Annual Revision	No criteria change.	6/15/2025

The policy effective date is in force until updated or retired.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.