



## Drug and Biologic Coverage Policy

Effective Date ..... 8/1/2022  
Next Review Date... 8/1/2023  
Coverage Policy Number ..... 1206

# Vascular Endothelial Growth Factor (VEGF) Inhibitors for Ocular Use

### Table of Contents

Coverage Policy.....	1
FDA Approved Indications.....	5
Recommended Dosing.....	6
General Background.....	8
Coding/ Billing Information.....	10
References.....	10

### Related Coverage Resources

#### INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

### Coverage Policy

Vascular Endothelial Growth Factor (VEGF) Inhibitors for Ocular Use includes the following products:

- **afibercept** for intravitreal injection (**Eylea®**)
- **brolocizumab-dbli** for intravitreal injection (**Beovu®**)
- **faricimab-svoa** intravitreal injection (**Vabysmo™**)
- **pegaptanib sodium** for intravitreal injection (**Macugen®**)
- **ranibizumab** for intravitreal injection (**Lucentis®**)
- **ranibizumab-nuna** for intravitreal injection (**Byooviz™**)

**Aflibercept for intravitreal injection (Eylea) is considered medically necessary for the treatment of ONE of the following:**

- Diabetic Macular Edema (DME)
- Diabetic Retinopathy (DR)
- Macular Edema following retinal vein occlusion (RVO)
- Neovascular (wet) Age-Related Macular Degeneration

- Other Neovascular Diseases of the Eye (for example, neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions)

**Brolucizumab-dblI intravitreal injection (Beovu) is considered medically necessary for the treatment of the following:**

- Neovascular (wet) age-related macular degeneration (AMD)

**Faricimab-svoa intravitreal injection (Vabysmo) is considered medically necessary for the treatment of the following:**

- Diabetic Macular Edema (DME)
- Neovascular (wet) age-related macular degeneration (AMD)

**Pegaptanib sodium for intravitreal injection (Macugen) is considered medically necessary for the treatment of ONE of the following:**

- Neovascular (wet) age-related macular degeneration (AMD)
- Other Neovascular Diseases of the Eye (for example, neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions)

**Ranibizumab for intravitreal injection (Lucentis) and ranibizumab-nuna for intravitreal injection (Byooviz) are considered medically necessary for the treatment of ONE of the following:**

- Diabetic Macular Edema (DME)
- Diabetic Retinopathy (DR)
- Macular Edema following Retinal Vein Occlusion (RVO)
- Neovascular (wet) Age-Related Macular Degeneration (AMD)
- Myopic Choroidal Neovascularization (mCNV)
- Ocular Histoplasmosis Syndrome
- Other Neovascular Diseases of the Eye (for example, neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions)

**Coverage for Vascular Endothelial Growth Factor (VEGF) Inhibitors for Ocular Use varies across plans and may require the use of preferred products in addition to the medical necessity criteria listed above. Refer to the customer's benefit plan document for coverage details.**

When coverage requires the use of preferred products, there is documentation of **ONE** of the following:

- A. The individual has had inadequate efficacy to the number of covered alternatives according to the table below

**OR**

- B. The individual has a contraindication according to FDA label, significant intolerance, or is not a candidate\* for the covered alternatives according to the table below

*\*Note: Not a candidate due to being subject to a warning per the prescribing information (labeling), having a disease characteristic, individual clinical factor[s], other attributes/conditions, or is unable to administer and requires this dosage formulation)*

**Employer Group Non-Covered Products and Preferred Covered Alternatives by Drug List:**

Non-Covered Product	Standard / Performance	Value / Advantage	Cigna Total Savings	Legacy
Beovu (brolucizumab-dblI) for intravitreal injection	<b>ONE</b> of the following (1 <u>or</u> 2): 1. Individual is currently receiving Beovu 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B): A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab			

Non-Covered Product	Standard / Performance	Value / Advantage	Cigna Total Savings	Legacy
				B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern
<b>Byooviz</b> (ranibizumab-nuna) for intravitreal injection				<b>ONE</b> of the following (1, 2 <u>or</u> 3): 1. Individual is currently receiving Byooviz 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B): A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern 3. Individual has diabetic retinopathy (without diabetic macular edema)
<b>Eylea</b> (afibercept) for intravitreal injection				<b>ONE</b> of the following (1, 2, 3, 4 <u>or</u> 5): 1. Individual is currently receiving Eylea 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B): A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern 3. Individual has diabetic macular edema with a baseline visual acuity worse than 20/40 4. Individual has diabetic macular edema with significant retinal thickening 5. Individual has diabetic retinopathy (without diabetic macular edema)
<b>Lucentis</b> (ranibizumab) for intravitreal injection				<b>ONE</b> of the following (1, 2 <u>or</u> 3): 1. Individual is currently receiving Lucentis 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B): A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern 3. Individual has diabetic retinopathy (without diabetic macular edema)
<b>Vabysmo</b> (faricimab-svoa) for intravitreal injection				<b>ONE</b> of the following (1 <u>or</u> 2): 1. Individual is currently receiving Vabysmo 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B): A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern

**Individual and Family Plan Non-Covered Products and Covered Alternative(s):**

Non-Covered Product	Criteria
<b>Beovu</b> (brolocizumab-dblil) for intravitreal injection	<b>ONE</b> of the following (1 <u>or</u> 2): 1. Individual is currently receiving Beovu 2. Individual meets <b>ONE</b> of the following (A <u>or</u> B):

Non-Covered Product	Criteria
	<p>A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab</p> <p>B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern</p>
<p><b>Byooviz</b> (ranibizumab-nuna) for intravitreal injection</p>	<p><b>ONE</b> of the following (1, 2 <u>or</u> 3):</p> <ol style="list-style-type: none"> <li>1. Individual is currently receiving Byooviz</li> <li>2. Individual meets <b>ONE</b> of the following (A <u>or</u> B):               <ol style="list-style-type: none"> <li>A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab</li> <li>B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern</li> </ol> </li> <li>3. Individual has diabetic retinopathy (without diabetic macular edema)</li> </ol>
<p><b>Eylea</b> (aflibercept) for intravitreal injection</p>	<p><b>ONE</b> of the following (1, 2, 3, 4 <u>or</u> 5):</p> <ol style="list-style-type: none"> <li>1. Individual is currently receiving Eylea</li> <li>2. Individual meets <b>ONE</b> of the following (A <u>or</u> B):               <ol style="list-style-type: none"> <li>A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab</li> <li>B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern</li> </ol> </li> <li>3. Individual has diabetic macular edema with a baseline visual acuity worse than 20/40</li> <li>4. Individual has diabetic macular edema with significant retinal thickening</li> <li>5. Individual has diabetic retinopathy (without diabetic macular edema)</li> </ol>
<p><b>Lucentis</b> (ranibizumab) for intravitreal injection</p>	<p><b>ONE</b> of the following (1, 2 <u>or</u> 3):</p> <ol style="list-style-type: none"> <li>1. Individual is currently receiving Lucentis</li> <li>2. Individual meets <b>ONE</b> of the following (A <u>or</u> B):               <ol style="list-style-type: none"> <li>A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab</li> <li>B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern</li> </ol> </li> <li>3. Individual has diabetic retinopathy (without diabetic macular edema)</li> </ol>
<p><b>Vabysmo</b> (faricimab-svoa) for intravitreal injection</p>	<p><b>ONE</b> of the following (1 <u>or</u> 2):</p> <ol style="list-style-type: none"> <li>1. Individual is currently receiving Vabysmo</li> <li>2. Individual meets <b>ONE</b> of the following (A <u>or</u> B):               <ol style="list-style-type: none"> <li>A. Documentation that the individual has had an inadequate response, has a contraindication per FDA label, or experienced a significant intolerance with repackaged bevacizumab</li> <li>B. If, in the professional opinion of the prescriber, the safety of using the repackaged bevacizumab or the supplier of the repackaged bevacizumab is of significant concern</li> </ol> </li> </ol>

**Initial authorization is up to 12 months.**

**Vascular Endothelial Growth Factor (VEGF) Inhibitors for Ocular Use are considered medically necessary for continued use when the following are met:**

- Individual continues to meet the initial criteria
- Attestation of beneficial clinical response

**Reauthorization for up to 12 months**

**When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.**

**Vascular Endothelial Growth Factor (VEGF) Inhibitors for Ocular Use are considered experimental, investigational or unproven for ANY other use including the following:**

- Best disease
- Coats' disease
- central serous chorioretinopathy
- radiation maculopathy
- refractory noninfectious uveitic cystoid macular edema

**The concomitant use of greater than one VEGF Inhibitor in the same eye is not covered as it is considered experimental, investigational or unproven.**

**Note: Receipt of sample product does not satisfy any criteria requirements for coverage.**

**FDA Approved Indications**

**FDA Approved Indication**

Brand Name	Approved Indication
<b>Eylea</b>	Eylea is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> <li>• Macular Edema Following Retinal Vein Occlusion (RVO)</li> <li>• Diabetic Macular Edema (DME)</li> <li>• Diabetic Retinopathy (DR)</li> </ul>
<b>Beovu</b>	Beovu is indicated for the treatment of patients: <ul style="list-style-type: none"> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> </ul>
<b>Byooviz</b>	Byooviz is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> <li>• Macular Edema Following Retinal Vein Occlusion (RVO)</li> <li>• Myopic Choroidal Neovascularization (mCNV)</li> </ul>
<b>Lucentis</b>	Lucentis is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> <li>• Macular Edema Following Retinal Vein Occlusion (RVO)</li> <li>• Diabetic Macular Edema (DME)</li> <li>• Diabetic Retinopathy (DR)</li> <li>• Myopic Choroidal Neovascularization (mCNV)</li> </ul>
<b>Macugen</b>	Macugen is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> </ul>
<b>Vabysmo</b>	Vabysmo is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Diabetic Macular Edema (DME)</li> <li>• Neovascular (Wet) Age-Related Macular Degeneration (AMD)</li> </ul>

## Recommended Dosing

### FDA Recommended Dosing

Brand Name	Recommended Dosing
Eylea	<p><b>Neovascular (Wet) Age-Related Macular Degeneration (AMD)</b>            The recommended dose for Eylea is 2 mg (0.05 mL or 50 microliters) administered by intravitreal injection every 4 weeks (approximately every 28 days, monthly) for the first 12 weeks (3 months), followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks (2 months). Although Eylea may be dosed as frequently as 2 mg every 4 weeks (approximately every 25 days, monthly), additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first 12 weeks (3 months). Although not as effective as the recommended every 8 week dosing regimen, patients may also be treated with one dose every 12 weeks after one year of effective therapy. Patients should be assessed regularly.</p> <p><b>Macular Edema Following Central Retinal Vein Occlusion (CRVO)</b>            The recommended dose for Eylea is 2 mg (0.05 mL or 50 microliters) administered by intravitreal injection once every 4 weeks (approximately every 25 days, monthly)</p> <p><b>Diabetic Macular Edema (DME)</b>            The recommended dose for Eylea is 2 mg (0.05 mL or 50 microliters) administered by intravitreal injection every 4 weeks (approximately every 28 days, monthly) for the first 5 injections, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks (2 months). Although Eylea may be dosed as frequently as 2 mg every 4 weeks (approximately every 25 days, monthly), additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first 20 weeks (5 months).</p> <p><b>Diabetic Retinopathy (DR)</b>            The recommended dose for Eylea is 2 mg (0.05 mL) administered by intravitreal injection every 4 weeks (approximately every 28 days, monthly) for the first 5 injections followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks (2 months). Although Eylea may be dosed as frequently as 2 mg every 4 weeks (approximately every 25 days, monthly), additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first 20 weeks (5 months).</p>
Beovu	<p>Beovu is administered by intravitreal injection. The recommended dose for Beovu is 6 mg (0.05 mL of 120 mg/mL solution) monthly (approximately every 25-31 days) for the first three doses, followed by one dose of 6 mg (0.05 mL) every 8-12 weeks.</p>
Byooviz	<p>Byooviz is administered by intravitreal injection. The recommended dose for Byooviz is 0.5 mg (0.05 mL of 10 mg/mL solution) monthly (approximately every 28 days).            Although not as effective, patients may be treated with 3 monthly doses followed by less frequent dosing with regular assessment. In the 9 months after three initial monthly doses, less frequent dosing with 4-5 doses on average is expected to maintain visual acuity while monthly dosing may be expected to result in an additional average 1-2 letter gain. Patients should be assessed regularly.            Although not as effective, patients may also be treated with one dose every 3 months after 4 monthly doses. Compared with continued monthly dosing, dosing every 3 months over the next 9 months will lead to an approximate 5-letter (1-line) loss of visual acuity benefit, on average. Patients should be assessed regularly.</p>

Brand Name	Recommended Dosing
<p><b>Lucentis</b></p>	<p><b>Neovascular (Wet) Age-Related Macular Degeneration (AMD)</b>  Lucentis 0.5 mg (0.05 mL of 10 mg/mL solution) is recommended to be administered by intravitreal injection once a month (approximately 28 days). Although not as effective, patients may be treated with 3 monthly doses followed by less frequent dosing with regular assessment. In the 9 months after three initial monthly doses, less frequent dosing with 4-5 doses on average is expected to maintain visual acuity while monthly dosing may be expected to result in an additional average 1-2 letter gain. Patients should be assessed regularly. Although not as effective, patients may also be treated with one dose every 3 months after 4 monthly doses. Compared with continued monthly dosing, dosing every 3 months over the next 9 months will lead to an approximate 5-letter (1-line) loss of visual acuity benefit, on average. Patients should be assessed regularly.</p> <p><b>Macular Edema Following Retinal Vein Occlusion (RVO)</b>  Lucentis 0.5 mg (0.05 mL of 10 mg/mL solution) is recommended to be administered by intravitreal injection once a month (approximately 28 days). In Studies RVO-1 and RVO-2, patients received monthly injections of Lucentis for six months. In spite of being guided by optical coherence tomography and visual acuity re-treatment criteria, patients who were then not treated at Month 6 experienced on average, a loss of visual acuity at Month 7, whereas patients who were treated at Month 6 did not.</p> <p><b>Diabetic Macular Edema (DME) and Diabetic Retinopathy (DR):</b>  Lucentis 0.3 mg (0.05 mL of 6mg/ ml solution) is recommended to be administered by intravitreal injection once a month (approximately 28 days).</p> <p><b>Myopic Choroidal Neovascularization (mCNV)</b>  Lucentis 0.5 mg (0.05 mL of 10 mg/mL solution) is recommended to be initially administered by intravitreal injection once a month (approximately 28 days) for up to 3 months. Patients may be retreated if needed.</p>
<p><b>Macugen</b></p>	<p>Macugen 0.3 mg should be administered once every six weeks by intravitreal injection into the eye to be treated.</p>
<p><b>Vabysmo</b></p>	<p><b>Neovascular (wet) Age-Related Macular Degeneration (nAMD)</b>  The recommended dose for VABYSMO is 6 mg (0.05 mL of 120 mg/mL solution) administered by intravitreal injection every 4 weeks (approximately every 28 ± 7 days, monthly) for the first 4 doses, followed by optical coherence tomography and visual acuity evaluations 8 and 12 weeks later to inform whether to give a 6 mg dose via intravitreal injection on one of the following three regimens: 1) Weeks 28 and 44; 2) Weeks 24, 36 and 48; or 3) Weeks 20, 28, 36 and 44. Although additional efficacy was not demonstrated in most patients when VABYSMO was dosed every 4 weeks compared to every 8 weeks, some patients may need every 4 week (monthly) dosing after the first 4 doses. Patients should be assessed regularly.</p> <p><b>Diabetic Macular Edema (DME)</b>  VABYSMO is recommended to be dosed by following one of these two dose regimens: 1) 6 mg (0.05 mL of 120 mg/mL solution) administered by intravitreal injection every 4 weeks (approximately every 28 days ± 7 days, monthly) for at least 4 doses. If after at least 4 doses, resolution of edema based on the central subfield thickness (CST) of the macula as measured by optical coherence tomography is achieved, then the interval of dosing may be modified by extensions of up to 4 week interval increments or reductions of up to 8 week interval increments based on CST and visual acuity evaluations through week 52; or 2) 6 mg dose of VABYSMO can be administered every 4 weeks for the first 6 doses, followed by 6 mg dose via intravitreal injection at intervals of every 8 weeks (2 months) over the next 28 weeks. Although additional efficacy was not demonstrated in most patients when VABYSMO was dosed every 4 weeks compared to every 8 weeks, some patients may need every 4 week (monthly) dosing after the first 4 doses. Patients should be assessed regularly.</p>



## Drug Availability

Brand Name	Drug Availability
<b>Eylea</b>	Single-use vial designed to provide 0.05 mL of 40 mg/mL solution for intravitreal injection.
<b>Beovu</b>	Single-use vial designed to provide 0.05 mL of 120 mg/mL solution for intravitreal injection.
<b>Byooviz</b>	Single-use vial designed to provide 0.05 mL of 10 mg/mL solution for intravitreal injection.
<b>Lucentis</b>	Single-use, 2-mL glass vial designed to provide 0.05 mL for intravitreal injection: <ul style="list-style-type: none"><li>• 6 mg/mL solution</li><li>• 10 mg/mL solution</li></ul> Single-use prefilled syringe designed to provide 0.05 mL for intravitreal injections: <ul style="list-style-type: none"><li>• 10 mg/mL solution</li></ul>
<b>Macugen</b>	Single-use glass syringe pre-filled with 0.3 mg of Macugen in a nominal 90 µL solution for intravitreal injection.
<b>Vabysmo</b>	Single-dose vial designed to provide 0.05 mL of 120 mg/mL solution for intravitreal injection.

## General Background

### Pharmacology

#### EYLEA

Aflibercept is a soluble decoy receptor, binding PIGF and all isoforms of VEGF-A with high affinity. Aflibercept interferes with the activation of the VEGF receptors, thereby preventing the cascade of events leading to neovascularization and vascular leakage.

#### LUCENTIS

Ranibizumab binds to the receptor binding site of active forms of VEGF-A, including the biologically active, cleaved form of this molecule, VEGF. VEGF-A has been shown to cause neovascularization and leakage in models of ocular angiogenesis and vascular occlusion, and is thought to contribute to the progression of neovascular AMD and macular edema following RVO. The binding of ranibizumab to VEGF-A prevents the interaction of VEGF-A with its receptors (VEGFR1 and VEGFR2) on the surface of endothelial cells, reducing endothelial cell proliferation, vascular leakage, and new blood vessel formation.

#### MACUGEN

Pegaptanib is a synthetic oligonucleotide aptamer with high affinity for VEGF. It delays choroidal neovascularization by binding VEGF, preventing it from binding to its receptor present on vascular endothelial cells of the subretinal choroid.

#### BEVACIZUMAB

Bevacizumab intravenous infusion (Avastin®, biosimilars) is another agent that is repackaged and administered by intravitreal injection for off-label use in AMD, diabetic macular edema (including patients with diabetic retinopathy), and other neovascular conditions of the eye. (AAO, 2019)

### Professional Societies/Organizations

#### American Academy of Ophthalmology (AAO)

The current preferred practice patterns published by the AAO indicate that anti-VEGF medications are first-line treatment options to treat and stabilize the majority of neovascular age-related macular degeneration (AMD) cases. The committee states that anti-VEGF therapy is usually well tolerated with few serious adverse events, such as infectious endophthalmitis or retinal detachment. If either of these adverse events is suspected, a timely evaluation is warranted. (AAO, 2019)

In 2015, citing a positive risk to benefit ratio associated with anti-VEGF treatment, the AAO preferred practice pattern regarding retinal vein occlusion states these medications should be the initial treatment for macular edema related to BRVO. The AAO further mentions several trials that have shown the efficacy of the anti-VEGFs in CRVO related macular edema, but does not call out the anti-VEGFs as initial therapy in this situation. The



organization describes this class of medications as the safest treatments for macular edema associated with BRVO and CRVO. (AAO, 2019)

The 2016 update of the AAO guidelines for diabetic retinopathy state that intravitreal anti-VEGF therapy is recommended as first line treatment for center involving macular edema, followed by laser treatment as necessary. In addition, the organization remarks that intravitreal anti-VEGF therapy is an alternative treatment in individuals with high-risk proliferative diabetic retinopathy without macular edema, citing data from one trial assessing the non-inferiority of intravitreal ranibizumab compared to panretinal photocoagulation for visual acuity outcomes in individuals with proliferative diabetic retinopathy. (AAO, 2019)

## **Clinical Efficacy**

### **Neovascular (Wet) Age-Related Macular Degeneration (AMD)**

Visual improvement was similar between bevacizumab and ranibizumab, and between aflibercept and ranibizumab in patients with neovascular AMD. Mean best corrected visual acuity (BCVA) letters increased from baseline for aflibercept (+6.6 to +10.9), bevacizumab (+7.4 to +8.5), and ranibizumab (+7.9 to +9.4), but decreased for pegaptanib (-7.3). Mean foveal thickness decreased from baseline for aflibercept (-116 to -157 micron), bevacizumab (-62 to -112 micron) and ranibizumab (-116 to -139 micron). Foveal thickness may decrease more with ranibizumab than bevacizumab. Foveal thickness reductions with pegaptanib were not addressed in any direct comparative trials. (Berg, 2015; Heier, 2012; Schmid-Kubista, 2011; Schmidt-Erfurth, 2014; Scholler, 2014; Solomon, 2014; Yuzawa, 2015)

### **Diabetic Macular Edema (DME)**

Intravitreal bevacizumab or ranibizumab cause similar improvements in BCVA from baseline, while foveal thickness may decrease more with ranibizumab than bevacizumab. Intravitreal aflibercept may cause greater improvement in BCVA and foveal thickness than bevacizumab or ranibizumab, especially in patients with baseline BCVA worse than 20/50. In a meta-analysis that evaluated indirect comparisons, BCVA improved significantly less with pegaptanib than aflibercept, bevacizumab, or ranibizumab; foveal thickness was not reported. Mean BCVA letters increased from baseline for aflibercept (+13.3), bevacizumab (+2.5 to +9.7), pegaptanib (+5.1, based on pooled data), and ranibizumab (+3 to +11.2). Foveal thickness decreased for aflibercept (-169 micron), bevacizumab (-101 to -141.5 micron), and ranibizumab (-136 to -150.5 micron). (Ekinci, 2014; Nepomuceno, 2013; Ollendorf, 2013; Wells, 2015)

### **Other Covered Use**

#### **Ocular Histoplasmosis Syndrome (OHS)**

The effect of ranibizumab on choroidal neovascularization, secondary to causes other than age-related macular degeneration, was studied in a randomized, controlled trial, and included 9 patients with ocular histoplasmosis syndrome. Thirty patients were randomized to either monthly intravitreal ranibuzimab or 3 monthly intravitreal injections followed by PRN injections of ranibizumab. Results revealed no difference between the groups in visual acuity or central retinal thickness. In addition, a retrospective chart review of 52 patients treated with either bevacizumab or ranibizumab showed significant improvement in visual acuity from baseline in patients with choroidal neovascularization secondary to OHS. (Heier, 2011; Nielsen, 2012)

#### **Other Neovascular Diseases of the Eye (for example, neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions)**

Overproduction of VEGF may lead to other eye conditions, including neovascular glaucoma, retinopathy of prematurity, and other retinal and choroidal neovascular conditions affecting the eye, the VEGF inhibitors also have the potential to be used off-label and reduce vision loss associated with other eye conditions related to increased VEGF production. (Barakat, 2009, Tolentino, 2011) The use of anti-VEGF agents have been shown to stop the angiogenic process and maintain visual acuity and improve vision in patients with certain neovascular ophthalmic conditions; therefore, research is rapidly evolving on the use of VEGF inhibitors in other neovascular ophthalmic conditions which threaten vision. (Kinnunen, 2012, Horsley, 2010) Anti-VEGF therapy has the potential to be used off-label in other neovascular conditions affecting the eye and may prevent or slow visual impairment. (Baraket, 2009, Kinnunen 2012, Horsley, 2010)

#### **Choroidal Neovascularization (CNV) secondary to angioid streaks (AS) or pseudoxanthoma elasticum**

Treatment with anti-vascular endothelial growth factor (VEGF) such as ranibizumab has demonstrated efficacy in the treatment of choroidal neovascularization (CNV) secondary to angioid streaks (AS) in small studies with a short follow-up duration. (Shah, 2012; Vadala, 2010) A long-term retrospective study reinforced the short-term outcomes. (Tilleul, 2016) All of these studies were in patients with angioid streaks-associated CNV in causes other than pseudoxanthoma elasticum (PXE). In angioid streaks-associated choroidal neovascularization (CNV) secondary to pseudoxanthoma elasticum (PXE) specifically, positive visual outcomes over a long term with anti-VEGF therapy has been reported in few case reports and smaller studies (Finger, 2011; Myung, 2010; Yilmaz, 2014; Zebardast, 2012) In France, ranibizumab was granted temporary reimbursement from October 2011 to October 2014, for the treatment of patients with choroidal neovascularization secondary to pseudoxanthoma elasticum. The PIXEL study was conducted to evaluate the long-term effectiveness and safety of ranibizumab 0.5 mg in 72 patients (98 eyes) with CNV secondary to PXE. The main objective was to assess changes in best-corrected visual acuity over time. The mean visual acuity (VA) was 64.6 letters at the first ranibizumab injection, which was maintained at the 1-year follow-up (64.7 letters). VA remained stable until the 4-year follow-up. (Mimoun, 2017)

### Experimental, Investigational, Unproven Uses

Macugen has been studied for use in diabetic macular edema, however, there is insufficient published data in terms of safety and efficacy to support its use for this indication. (Ollendorf, 2013)

There is insufficient evidence in the peer-reviewed published scientific literature to support the safety and efficacy of the VEGF inhibitors in Best disease (Chaudhary, 2013), Coats' disease (Chaudhary, 2013), central serous chorioretinopathy (Salehi, 2015), radiation maculopathy (Seibel, 2016), refractory noninfectious uveitic cystoid macular edema (Lasave, 2009) and uveitis with childhood neovascularization (Chaudhary, 2013).

In addition, there are no published, randomized controlled trials evaluating the efficacy and safety of the concurrent use of intravitreal VEGF inhibitors.

## Coding/ Billing Information

- Note:** 1) This list of codes may not be all-inclusive.  
 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

### Covered when medically necessary:

HCPCS Codes	Description
J0178	Injection, aflibercept, 1 mg
J2503	Injection, pegaptanib sodium, 0.3 mg
J2778	Injection, ranibizumab, 0.1 mg
J3590	Unclassified biologics

## References

1. American Academy of Ophthalmology Retina/Vitreous Panel. Preferred Practice Pattern® Guidelines. Age-related macular degeneration. San Francisco, CA: American Academy of Ophthalmology; 2019. Available at: <https://www.aao.org/preferred-practice-pattern/age-related-macular-degeneration-ppp>. Accessed on May 27, 2021.
2. American Academy of Ophthalmology Retina/Vitreous Panel. Preferred Practice Pattern® Guidelines. Diabetic retinopathy. San Francisco, CA: American Academy of Ophthalmology; 2019. Available at: <https://www.aao.org/preferred-practice-pattern/diabetic-retinopathy-ppp>. Accessed on May 27, 2021.
3. Barakat MR, Kaiser PK. VEGF inhibitors for the treatment of neovascular age-related macular degeneration. Expert Opin Investig Drugs. 2009;18(5):637-646
4. Beovu® intravitreal injection [prescribing information]. Hanover, NJ: Novartis; June 2020.

5. Berg K, Pedersen T, Sandvik L, et al. Comparison of ranibizumab and bevacizumab for neovascular age-related macular degeneration according to LUCAS treat-and-extend protocol. *Ophthalmology*. Jan 2015;122(1):146-152
6. Byooviz™ intravitreal injection [prescribing information]. Cambridge, MA: Biogen Inc.; June 2022.
7. Casparis H, Wolfensberger TJ, Becker M, et al. Incidence of presumed endophthalmitis after intravitreal injection performed in the operating room: a retrospective multicenter study. *Retina*. Jan 2014;34(1):12-17.
8. Curtis L, Hammill B, Schulman K, et al. Risks of mortality, myocardial infarction, bleeding, and stroke associated with therapies for age-related macular degeneration. *Arch Ophthalmol*. Oct 2010;128(10):1273-1279.
9. Chaudhary KM, Mititelu M, Lieberman RM. An evidence-based review of cascular endothelial growth factor inhibition in pediatric retinal diseases: part 2. Coats' disease, best disease, and uveitis with childhood neovascularization. *J Pediatr Ophthalmol Strabismus*. 2013 Jan-Feb;50(1):11-9.
10. Diabetic Retinopathy Clinical Research Network, Wells J, Glassman A, et al. Supplementary appendix to: Aflibercept, bevacizumab, or ranibizumab for diabetic macular edema. *N Engl J Med*. Mar 26 2015;372(13):1193-1203.
11. Ekinci M, Ceylan E, Çakıcı Ö, Tanyıldız B, Olcaysu O, Çağatay HH. Treatment of macular edema in diabetic retinopathy: comparison of the efficacy of intravitreal bevacizumab and ranibizumab injections. *Expert Review of Ophthalmology*. 2014; 9(2):139-143.
12. Eyetech Inc. Macugen (pegaptanib sodium injection) package insert. Palm Beach Gardens, Florida: Eyetech, Inc. Oct 2011.
13. Finger RP, Charbel Issa P, Hendig D, et al. Monthly Ranibizumab for Choroidal Neovascularizations Secondary to Angioid Streaks in Pseudoxanthoma Elasticum: A One-Year Prospective Study. *Am J Ophthalmol*. 2011 Oct;152(4):695-703.
14. Genentech Inc. Lucentis (ranibizumab injection) Intravitreal Injection [product information]. South San Francisco, CA: Genentech, Inc. April 2017.
15. Heier J, Brown D, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology*. Dec 2012; 119(12):2537-2548.
16. Heier JS et al. Ranibizumab for choroidal neovascularization secondary to causes other than age-related macular degeneration: a phase 1 clinical trial. *Ophthalmology* 2011; 118: 111-8.
17. Horsley MB, Kahook MY. Anti-VEGF therapy for glaucoma. *Curr Opin Ophthalmol*. 2010; 21(2):112-117.
18. Inoue M, Kobayakawa S, Sotozono C, et al. Evaluation of the incidence of endophthalmitis after intravitreal injection of anti-vascular endothelial growth factor. *Ophthalmologica*. 2011; 226(3):145-150.
19. Johnson D, Jagan L, Kurji A, et al. Emergency department visits after intravitreal bevacizumab and ranibizumab injections in diabetic patients. *Can J Ophthalmol*. Dec 2014; 49(6):e146-148.
20. Kinnunen K, Ylä-Herttua S. Vascular endothelial growth factors in retinal and choroidal neovascular diseases. *Ann Med*. 2012; 44(1):1-17.
21. Lasave AF, Zeballos DG, El-Haig WM, et al. Short-term results of a single intravitreal bevacizumab (Avastin) injection versus a single intravitreal triamcinolone acetonide (Kenacort) injection for the management of refractory noninfectious uveitic cystoid macular edema. *Ocul Immunol Inflamm* 2009; 17:423-430.
22. McEvoy GK, ed. 2019. AHFS Drug Information® - 60th Ed. Bethesda, MD. American Society of Health-System Pharmacists.
23. Mimoun G, Ebran JM, Grenet T, et al. Ranibizumab for choroidal neovascularization secondary to pseudoxanthoma elasticum: 4-year results from the PIXEL study in France. *Graefes Arch Clin Exp Ophthalmol*. 2017 Aug; 255(8): 1651-1660.
24. Moja L, Lucenteforte E, Kwag Koren H, et al. Systemic safety of bevacizumab versus ranibizumab for neovascular age-related macular degeneration. *Cochrane Database Syst Rev*. 2014;9:CD011230.
25. Myung JS, Bhatnagar P, Spaide RF, et al. Long-term outcomes of intravitreal anti-vascular endothelial growth factor therapy for the management of choroidal neovascularization in pseudoxanthoma elasticum. *Retina*. 2010 May;30(5):748-55.
26. Nepomuceno A, Takaki E, Paes de Almeida F, et al. A prospective randomized trial of intravitreal bevacizumab versus ranibizumab for the management of diabetic macular edema. *Am J Ophthalmol*. Sep 2013;156(3):502-510 e502
27. Ollendorf DA, Colby JA, Pearson SD. Comparative effectiveness of anti-VEGF agents for diabetic macular edema. *Int J Technol Assess Health Care*. Oct 2013;29(4):392-40.
28. Regeneron Pharmaceuticals Inc. Eylea (aflibercept) Injection for intravitreal injection product information. Tarrytown, NY: Regeneron Pharmaceuticals, Inc. May 2019.

29. Salehi M, Wenick AS, Law HA, et al. Interventions for central serous chorioretinopathy: a network meta-analysis. *Cochrane Database Syst Rev*. 2015 Dec 22;(12):CD011841.
30. Schmid-Kubista K, Krebs I, Ansari-Shahrezaei S, et al. Comparing treatment of neovascular age-related macular degeneration with sequential intravitreal Avastin and Macugen versus intravitreal mono-therapy--a pilot study. *Curr Eye Res*. Oct 2011;36(10):958-963.
31. Schmidt-Erfurth U, Kaiser PK, Korobelnik JF, et al. Intravitreal aflibercept injection for neovascular age-related macular degeneration: ninety-six-week results of the VIEW studies. *Ophthalmology*. Jan 2014;121(1):193-201.
32. Scholler A, Richter-Mueksch S, Weingessel B, et al. Differences of frequency in administration of ranibizumab and bevacizumab in patients with neovascular AMD. *Wien Klin Wochenschr*. Jun 2014;126(11-12):355-359.
33. Seibel I, Hager A, Riechardt AI, et al. Antiangiogenic or Corticosteroid Treatment in Patients With Radiation Maculopathy After Proton Beam Therapy for Uveal Melanoma. *Am J Ophthalmol*. 2016 Aug;168:31-39.
34. Shah M and Amoaku. Intravitreal ranibizumab for the treatment of choroidal neovascularisation secondary to angioid streaks. *WMK. Eye (Lond)*. 2012 Sep; 26(9): 1194–1198.
35. Solomon S, Lindsley K, Vedula Satyanarayana S, et al. Anti-vascular endothelial growth factor for neovascular age-related macular degeneration. *Cochrane Database Syst Rev*. Aug 29 2014;8:CD005139.
36. Tilleul J, Mimoun G, Querques G, et al. INTRAVITREAL RANIBIZUMAB FOR CHOROIDAL NEOVASCULARIZATION IN ANGIOID STREAKS: Four-Year Follow-up. *Retina*. 2016 Mar;36(3):483-91.
37. Tolentino M. Systemic and ocular safety of intravitreal anti-VEGF therapies for ocular neovascular disease. *Surv Ophthalmol*. 2011; 56(2):95-113.
38. Vabysmo™ intravitreal injection [prescribing information]. South San Francisco, CA: Genentech/Roche; January 2022.
39. Vadala M, Pece A, Cipolla S, Monteleone C, et al. Angioid streak-related choroidal neovascularization treated by intravitreal ranibizumab. *Retina*. 2010 Jun;30(6):903-7.
40. Writing Committee for the Diabetic Retinopathy Clinical Research Network. Panretinal photocoagulation vs intravitreal ranibizumab for proliferative diabetic retinopathy: a randomized clinical trial. *JAMA* 2015;314:2137-46.
41. Yilmaz I, Ozkaya A, Alkin Z, et al. Intravitreal ranibizumab for bilateral choroidal neovascularisation in a patient with angioid streaks. *BMJ Case Rep*. 2014 Jul 29;2014.
42. Yuzawa M, Fujita K, Wittrup-Jensen K, et al. Improvement in vision-related function with intravitreal aflibercept: data from phase 3 studies in wet age-related macular degeneration. *Ophthalmology*. Mar 2015;122(3):571-578.
43. Zebardast N, Adelman RA. Intravitreal Ranibizumab for Treatment of Choroidal Neovascularization Secondary to Angioid Streaks in Pseudoxanthoma Elasticum - Five-year Follow-up. *Semin Ophthalmol*. 2012 May-Jul;27(3-4):61-4.

---

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2022 Cigna.