

Diagnosis (check all that apply to your patient):

- acquired hemophilia A
- acquired inhibitor titer to Factor VIII
- acquired inhibitors to factors XI or XII
- coagulation factor X deficiency
- congenital fibrinogen deficiency (factor I deficiency)-afibrinogenemia
- congenital fibrinogen deficiency (factor I deficiency)-hypofibrinogenemia
- congenital fibrinogen deficiency (factor I deficiency)-dysfibrinogenemia
- congenital factor VII (FVII) deficiency
- congenital factor XIII A-subunit deficiency
- congenital factor XIII B-subunit deficiency
- congenital Factor XIII deficiency
- factor II deficiency
- factor VIII deficiency (hemophilia A)
- factor IX deficiency (hemophilia B)
- factor X deficiency
- factor XIII deficiency
- Glanzmann's thrombasthenia with refractoriness to platelet transfusions
- hemophilia A
- hemophilia A (congenital factor VIII deficiency)
- hemophilia A with inhibitors
- hemophilia B
- hemophilia B with inhibitors
- hereditary antithrombin deficiency (antithrombin III deficiency, AT III deficiency)
- hereditary Factor X deficiency
- inhibitors to factors XI or XII
- severe von Willebrand disease (VWD)
- mild or moderate von Willebrand disease (VWD)
- Other (please specify):

Clinical Information

****FEIBA, NovoSeven RT, Obizur, SEVENFACT and Tretten: These drugs requires supportive documentation (chart notes, lab/test results, etc) be attached with this request****

(if AlphaNine SD, Alprolix, BeneFIX, Idelvion, Ixinity, Profilnine, Rebinyn, or Rixubis) Is this agent prescribed by (or in consultation with) a hemophilia specialist? Yes No

(if AlphaNine SD, Alprolix, BeneFIX, Idelvion, Ixinity, Profilnine, Rebinyn, or Rixubis AND has hemophilia B) Is this medication being used as on-demand treatment and control of bleeding episodes? Yes No

(if no) Is this medication being used for routine prophylaxis? Yes No

(if no) Is this medication being used for perioperative management? Yes No

(if no and requesting **AlphaNine SD, BeneFIX, Ixinity, Profilnine or Rixubis**) Is this medication being used for immune tolerance therapy (also known as immune tolerance induction)? Yes No

(if ATryn) Is ATryn being used for the prevention of perioperative or peripartum events? Yes No

(if Coagadex)

For which of the following is this drug being used?

- Peri-operative management of bleeding in individuals with mild or moderate hereditary Factor X deficiency
- Routine prophylaxis to reduce the frequency of bleeding episodes
- Treatment of bleeding episodes
- Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

Is the requested medication being prescribed by (or in consultation with) a hematologist? Yes No

(if Altuviio)

For which of the following is this drug being used?

- Peri-operative management of bleeding
- Routine prophylaxis
- On-demand treatment and control of bleeding episodes
- Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

Is this a request for initial therapy or is the patient currently receiving the requested medication (or they have in the past)? If patient has been taking samples, please pick 'initial therapy'.

- Initial therapy
 Currently receiving the requested medication (or they have in the past)

(if currently receiving therapy or have in the past) Does the patient have clinical manifestations suggesting the presence of Factor VIII inhibitors? Please Note: Inhibitors may be present if bleeding is not well controlled, there is decreased responsiveness to Factor VIII therapy, and/or if expected Factor VIII activity plasma levels are not achieved. Yes No

(if currently receiving therapy or have in the past) Has Factor VIII inhibitor testing been performed within the last 365 days? Yes No

(if currently receiving therapy or have in the past) Does the patient have a positive test for Factor VIII inhibitors greater than or equal to 0.6 Bethesda units/mL? Yes No

(if initial therapy) Has the patient received Factor VIII therapy in the past? Yes No

(if initial therapy) Has Factor VIII inhibitor testing been performed within the last 30 days? Yes No

(if initial therapy) Does the patient have a positive test for Factor VIII inhibitors greater than or equal to 1.0 Bethesda units/mL? Yes No

Is the requested medication being prescribed by, or in consultation with, a hemophilia specialist? Yes No

(if Corifact)

For which of the following is this drug being used?

- Peri-operative management of bleeding
 Routine prophylaxis to reduce the frequency of bleeding episodes
 Treatment of bleeding episodes
 Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

Is the requested medication being prescribed by (or in consultation with) a hematologist? Yes No

(if Tretten)

For which of the following is this drug being used?

- Peri-operative management of bleeding
 Routine prophylaxis to reduce the frequency of bleeding episodes
 Treatment of bleeding episodes
 Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

Is the requested medication being prescribed by (or in consultation with) a hematologist? Yes No

(if Vonvendi)

For which of the following is this drug being used?

- Peri-operative management of bleeding
 Routine prophylaxis to reduce the frequency of bleeding episodes in individuals with severe Type 3 von Willebrand disease
 Treatment of bleeding episodes
 Other

(if other) Please provide clinical rationale for the use of this drug in your patient.

Is the requested medication being prescribed by (or in consultation with) a hematologist? Yes No

(if Obizur)

(if acquired hemophilia) Has there been documentation provided of autoimmune inhibitory antibodies to human factor VIII? Yes No

For which of the following is this drug being used?

- treatment of current active bleed
 prevention of excessive bleeding during and/or following surgery
 routine prophylaxis
 as needed dosing for future bleeds
 Other

(if surgery) What is the date of surgery?

(if as needed dosing) What is the approximate number of bleeds requiring factor treatment per month?
(if other) Please provide clinical rationale for the use of this drug in your patient.

(if NovoSeven RT)

(if Glanzmann's thrombasthenia) Is the patient refractory to platelet transfusions?

Yes No

Is this medication prescribed by, or in consultation with, a hematologist?

Yes No

(if Feiba or Sevenfact)

Is the drug requested being prescribed by, or in consultation with, a hematologist?

Yes No

(if Hemophilia A with inhibitors) Does the patient have a positive inhibitor titer at least 5 Bethesda Units or greater?

Yes No

(if no to previous question) Does the patient have a history of an inhibitor with anamnestic response to Factor VIII replacement therapy, which precludes the use of Factor VIII replacement to treat bleeding episodes?

Yes No

(if no to previous question) Does the patient have a history of an inhibitor with refractory hemostatic response to increased Factor VIII dosing, which precludes the use of Factor VIII replacement to treat bleeding episodes?

Yes No

(if Hemophilia B with inhibitors) Does the patient have a positive inhibitor titer at least 5 Bethesda Units or greater? Yes No

(if no to previous question) Does the patient have a history of an inhibitor with anamnestic response to Factor IX replacement therapy, which precludes the use of Factor IX replacement to treat bleeding episodes?

Yes No

(if no to previous question) Does the patient have a history of an inhibitor with refractory hemostatic response to increased Factor IX dosing, which precludes the use of Factor IX replacement to treat bleeding episodes?

Yes No

Additional pertinent information: Please provide any additional pertinent clinical information, including: if the patient is currently on the requested drug (with dates of use) and how they have been receiving it (samples, out of pocket, etc).

Attestation: I attest the information provided is true and accurate to the best of my knowledge. I understand that the Health Plan or insurer its designees may perform a routine audit and request the medical information necessary to verify the accuracy of the information reported on this form.

Prescriber Signature: _____ **Date:** _____

Save Time! Submit Online at: www.covermymeds.com/main/prior-authorization-forms/cigna/ or via SureScripts in your EHR.

Our standard response time for prescription drug coverage requests is 5 business days. If your request is urgent, it is important that you call us to expedite the request. View our Prescription Drug List and Coverage Policies online at cigna.com.

V011525

"Cigna" is a registered service mark, and the "Tree of Life" logo is a service mark, of Cigna Intellectual Property, Inc., licensed for use by Cigna Corporation and its operating subsidiaries. All products and services are provided by or through such operating subsidiaries and not by Cigna Corporation. Such operating subsidiaries include, for example, Cigna Health and Life Insurance Company and Cigna Health Management, Inc. Address: Cigna Pharmacy Services, PO Box 42005, Phoenix AZ 85080-2005